

NORBERT SCHEU

ON THE COMPUTATION OF
STRUCTURE FUNCTIONS AND MASS SPECTRA
IN A RELATIVISTIC HAMILTONIAN FORMALISM:
A LATTICE POINT OF VIEW

Thèse
présentée
à la Faculté des Études Supérieures
de l'Université Laval
pour l'obtention
du grade de Philosophiæ Doctor (Ph.D.)

Département de physique
FACULTÉ DES SCIENCES ET DE GÉNIE
UNIVERSITÉ LAVAL
QUÉBEC

DÉCEMBRE, 1997

©Norbert Scheu, 1997

床前明月光
疑是地上霜
举头望明月
低头思故乡

李白

静
夜
思

Quiet thought at night

Bright moon light in front of my bed,
Maybe... frost on the ground.
I raise my head, behold the bright
moon,
I bow my head— home-sick.

[Lǐ Bái, Tāng Dynasty poet]

Abstract for HEP-TH

Modified version of Ph.D. thesis

Herein we propose a new numerical technique for solving field theories: the large momentum frame (LMF). This technique combines several advantages of lattice gauge theory with the simplicity of front form quantisation. We apply the LMF on QED(1+1) and on the $\phi^4(3+1)$ theory. We demonstrate both analytically and in practical examples (1) that the LMF does neither correspond to the *infinite* momentum frame (IMF) nor to the front-form (FF) (2) that the LMF is not equivalent to the IMF (3) that the IMF is unphysical since it violates the lattice scaling window and (4) that the FF is even more unphysical because FF propagators violate micro-causality, causality and the finiteness of the speed of light. We argue that distribution functions measured in deep inelastic scattering should be interpreted in the LMF (preferably in the Breit frame) rather than in the FF formalism. In particular, we argue that deep inelastic scattering probes *space*-like distribution functions.

Résumé I

A non-perturbative computation of HADRONIC STRUCTURE FUNCTIONS for deep inelastic lepton hadron scattering has not been achieved yet. In this thesis we investigate the viability of the Hamiltonian approach in order to compute hadronic structure functions. In the literature, the so-called FRONT FORM (FF) approach is favoured over the conventional the INSTANT FORM (IF, i.e. the conventional Hamiltonian approach) due to claims (a) that structure functions are related to LIGHT-LIKE CORRELATION FUNCTIONS and (b) that the front form is much simpler for numerical computations. We dispell both claims using general arguments as well as practical computations (in the case of the SCALAR MODEL and TWO-DIMENSIONAL QED) demonstrating (a) that structure functions are related to SPACE-LIKE CORRELATIONS and that (b) the IF is better suited for practical computations if appropriate approximations are introduced. Moreover, we show that the FF is UNPHYSICAL in general for reasons as follows: (1) the FF constitutes an INCOMPLETE QUANTISATION of field theories (2) the FF "predicts" an INFINITE SPEED OF LIGHT in one space dimension, a COMPLETE BREAKDOWN OF MICROCAUSALITY and the UBIQUITY OF TIME-TRAVEL. Additionally we demonstrate that the FF cannot be approached by so-called ε co-ordinates. We demonstrate that these co-ordinates are but the instant form in disguise. We argue that the FF cannot be considered to be an EFFECTIVE THEORY. Finally, we demonstrate that the so-called INFINITE MOMENTUM FRAME is neither physical nor equivalent to the FF.

Signé par

(Norbert Scheu)

(Helmut Kröger)

Résumé II

In this Ph.D. thesis we demonstrate that

1. a numerical diagonalisation of a lattice regularised Hamiltonian may become drastically simpler, for some field theories, if the lattice moves fast relative to the object which is to be described.

We propose a new numerical technique based on this simplification. We apply this technique to the massive Schwinger model and the scalar model

2. θ vacua and spectral flow arise naturally in our approach
3. structure functions are related to space-like correlation functions rather than to light-like correlation functions
4. the notion 'closeness to the light-cone' is irreconcilable with the theory of relativity
5. ε co-ordinates are completely equivalent to the instant form for $\varepsilon \neq 0$.
6. the conventional instant-form has to be used in order to obtain non-perturbative input for the parton model
7. the front-form is not equivalent to the instant form *in general*. We explain under which circumstances the front-form is able to come close to the accuracy of the instant form: Prominent examples are almost non-relativistic theories, scale-invariant theories without boson-boson couplings and some graphs in perturbation theory
8. microcausality and the finiteness of the speed of light are completely destroyed by light-like boundary conditions — however large the periodicity may be

9. even if the front-form describes the mass spectrum of a theory quite accurately, relativistic propagators are badly damaged by front form quantisation
10. the front form canNOT be considered to be an effective theory for the quantised instant form. The front-form is not even equivalent to the quantised infinite momentum frame. We remark on are a few, well-defined exceptions to this rule.
11. the UNQUANTISED front form is an effective theory for the UNQUANTISED infinite momentum frame, however.
12. beyond the mean field level, there is, in general, no physical information in zero mode constraints which arise in the front from. On the contrary, a correct implementation of zero-mode constraints adds further damage to the already damaged propagators
13. it is not true that computations in the front form are simpler than in the instant form
14. the FF is not needed in order to treat two-dimensional models in a simple, numerical way. For instance, the front form yields accurate mass spectra for the Schwinger model. So does the instant form, only with higher accuracy and less effort (only a finite lattice is needed)
15. the instant form effortlessly reproduces chiral perturbation theory on small (effective) lattices whereas an infinite lattice or ad hoc counterterms are necessary to this aim in the front form
16. the infinite momentum frame is less unphysical than the front form but unphysical nonetheless because it violates the the lattice scaling window

Signé par

(Norbert Scheu)

(Helmut Kröger)

Avant-propos

À CHRISTINA — QUAND MÊME ☺ ☺

Je tiens à souligner ma vive reconnaissance envers le Prof. Dr. Helmut Kröger —mon directeur de thèse— pour son appui dans ce projet. J'ai beaucoup profité de la collaboration avec Helmut, de sa porte toujours ouverte, des discussions avec lui et de sa connaissance profonde d'un grand nombre de domaines en physique. J'ai particulièrement apprécié sa compétence dans le domaine des phénomènes critiques et de la théorie de jauge sur réseau et son encouragement à poursuivre mes idées.

Merci beaucoup aussi aux "habitants" de la tour d'ivoire: à Baabak (et Raamak et le petit Mazdak), Bertrand, Frédéric, Ghislain, Gurgen, Gwendoline, Hamza (et Laurence), Jean-François (Audet), Jean-François (Addor), Luc, Marek, Michel, Nicolas, Patrick, Peter, Pierre, Robert (et Anne), Simon, Stéphane, Yorgo, Yves, et Ziad. J'ai beaucoup aimé la bonne et amicale atmosphère qui régnait dans cette salle et j'en garderai un très bon souvenir. Je voudrais aussi mentionner Alain, Ali, Denis et Danielle, Francine Caron et famille, Frants, Gilberto et Bartira, Jean-François, Lionel, Marius, Mike, Pierre, Raymond, M.Slobodrian, et Tim .

Je remercie Messieurs les Professeurs Amiot, Marleau et Potvin d'avoir accepté de corriger ma thèse.

Il me fait très plaisir de remercier Claudette, Colette, Diane, Francine et Lise, les secrétaires, car elles ont toujours été gentilles et toujours prêtes à aider.

$T\tilde{\eta} \alpha\gamma\nu\tilde{\omega}\tau\eta, \tau\tilde{\omega} \alpha\gamma\nu\tilde{\omega}\tau\omega$: merci aussi à tous ceux que je pourrais avoir oubliés dans l'ardeur de finir la rédaction de ma thèse.

Je me suis toujours senti bien à l'aise *icitte* ☺ au Canada... mise à part —bien sûr!— *certaines* coupures du budget universitaire ☺ et le temps

froid qu'il fait en hiver ☺ !

Acknowledgements

The author wants to express his appreciation for having been granted the AUFÉ fellowship from the DAAD (Deutscher Akademischer Austauschdienst) which has made this Ph.D. project possible. We are grateful for many discussions with Prof. Dr. Dieter Schütte and Prof. Dr. Xiàng-Qián Lù.

Contents

1	Introduction	1
1	The Computation of Structure Functions: A Non-Perturbative Problem	1
2	Lattice Gauge Theory	2
3	A Brief Review of Hamiltonian Methods	5
4	Advantages of the Lorentz-Contraction: Proposal and Test of a New Technique	6
5	Does the Hamiltonian exist?	8
6	Is Quantisation on Light-Like Quantisation Surfaces Viable? .	9
7	Organisation	12
8	Notation	13
9	Methodology	15
2	Structure Functions as Short-Distance Physics	17
1	The Hadronic Tensor and Structure Functions	21
2	The Hadronic Tensor: Formal Definition	22
3	Structure Functions and Distribution Functions	25
4	Parton Distribution Functions: Two Definitions	34
5	Special Case: Breit Frame	39
6	Distribution Functions in Other Frames	40

7	Beyond the Impulse Approximation	44
7.1	The Breit Frame and the Continuum Limit	48
3	Instant-Form and Front-Form: A Campaign For Real Time	49
1	The Problem of Front-Form Quantisation	49
2	Planes, Vectors and Frames	55
3	The Quantisation Hyper-Surface	58
4	The Boundary Vector	62
5	Kinematical Equivalence of Relativistic Frames	66
6	Defining Boost Invariance	69
7	The Front Form and ε Co-ordinates	71
7.1	A Sketch of the Problem	71
7.2	Kinematical Equivalence	74
7.3	Boosting the Lattice	76
7.4	Conclusion	77
8	The Infinite Momentum Frame	78
8.1	Operational definition of the IMF	78
8.2	Constraints and Boundary Conditions in the IMF . . .	80
8.3	How To Construct an Effective Hamiltonian	82
8.4	The Classically Effective Hamiltonian	84
8.5	The Quantum Effective Hamiltonian	86
8.6	Some Comments on Zero Modes in the FF	89
9	The Breakdown of Causality in the Front Form	90
9.1	The Causality Region	90
9.2	The Instant Form Perspective	94
9.3	Light-Like Boundary Conditions in the Instant Form .	96
10	The Limit of Infinite Light-Like Volume	96
11	Situations Where the FF May Be a Good Approximation . . .	99

11.1	The Non-relativistic Limit	99
11.2	Perturbation Theory With External Input	99
11.3	Theories Without Vertices Which Connect at Least Four Bosons	100
11.4	Accidental Cases	100
11.5	Phenomenology	100
12	Other Relativistic Forms	100
4	The Massless Schwinger Model	103
1	The Exact Solution to the Schwinger Model: A Brief Review	106
1.1	Terminology and Notation	106
1.2	Operators in Momentum Space	109
1.3	The Hamiltonian in Manton's basis	110
2	Translation into the Particle-Antiparticle Picture	117
3	Solutions to the Schwinger Model and the Influence of Ap- proximations	122
4	The Schwinger Model on an Infinitesimal Lattice	124
5	Theta Vacua and Cut-Off Regularisation	127
6	Conclusions for Numerical Computations	130
7	The Schwinger Model on a Large Lattice	131
8	The Vacuum Distribution Function: Axial Vacua	133
9	The Vacuum Distribution Function: Gauge Invariant Vacua	135
10	Epilogue: The Front Form and the Schwinger Model	136
5	The Massive Schwinger Model	139

1	Scaling Window and Region of Validity	143
2	Convergence and Covariance	146
3	Numerical Results	147
4	Distribution Functions	149
5	The Mass of the Vector Boson	155
6	The Mass of the Scalar Boson	162
7	The Modified Front Form	163
8	The Infinite Momentum Frame	169
9	Why the Schwinger Model is Special	173
10	The Influence of the θ Angle	174
6	The ϕ^4 Theory	179
7	Discussion	180
1	Relativistic Forms	180
2	ε Co-ordinates	184
3	The Massive Schwinger Model	185
4	The Massless Schwinger Model	187
5	Structure Functions	189

List of Tables

2.1	Overview of polarisation vectors	18
2.2	Overview of distribution functions	35
3.1	Relativistic forms	52
3.2	Three-dimensional hyper-planes and their corresponding normal 4-vectors	63
3.3	Conserved quantities in the presence of boundary conditions .	66
3.4	The Useful Stability Group	102
4.1	Vacua: a systematic collection	128

List of Figures

3.1	The IF co-ordinates x^0, x^3 and their corresponding FF co-ordinates $x^\pm \stackrel{def}{=} x^0 \pm x^3$	51
3.2	Illustration of charge non-conservation. The boundary conditions join A and B	64
3.3	Illustration of microcausality violation. The boundary conditions join A and B	64
3.4	The shrinking of the causality region $\mathcal{C}$ in a fixed frame	91
3.5	The shrinking of the causality region $\mathcal{C}_\varepsilon$ in ε co-ordinates . . .	91
4.1	Symbolic representation of the vacuum $ \Omega : \mathbf{A}_3, \mathcal{M}\rangle$ with $\mathcal{M} = 0$ and $\mathbf{A}_3 = 0$	113
4.2	Symbolic representation of a Schwinger boson	129
4.3	The meson cloud	133
4.4	Admissible BCs for the FF	137
5.1	Convergence of distribution functions in the chiral limit $m/g = 1/8$	148
5.2	Convergence of the distribution functions in the FF, $m/g = 1/8$	149
5.3	The number of fermions in a boson	150
5.4	The number of fermions in a boson (FF)	151
5.5	Convergence of distribution functions for $m/g = 32$	152

5.6	Overview of distribution functions on the entire range of fermion masses	153
5.7	Distribution functions (LMF) in the chiral limit $m/g \rightarrow 0$. . .	154
5.8	Distribution functions (FF) in the ultra-relativistic parameter region	155
5.9	The vector boson mass M_V . A comparison with chiral perturbation theory	156
5.10	The vector boson mass in the FF. A comparison with chiral perturbation theory	157
5.11	The binding energy of the vector boson	158
5.12	The binding energy of the vector boson in the FF	159
5.13	The binding energy of the scalar boson	162
5.14	The binding energy of the scalar boson in the FF	163
5.15	The vector boson mass in the modified FF. A comparison with chiral perturbation theory	164
5.16	The number of fermions in a boson (modified FF)	165
5.17	The binding energy of the vector boson in the modified FF . .	166
5.18	Distribution functions in the chiral limit $m/g \rightarrow 0$ (modified FF)	167
5.19	The binding energy of the scalar boson in the modified FF . .	168
5.20	The vector boson mass in the IMF. A comparison with chiral perturbation theory	169
5.21	The binding energy of the vector boson in the IMF	170
5.22	The binding energy of the scalar boson in the IMF.	171
5.23	The lowest-lying mass spectrum (for very small fermion masses)	175
5.24	The naïve mass spectrum (for very small fermion masses) . . .	176

Chapter 1

Introduction

1 The Computation of Structure Functions: A Non-Perturbative Problem

Ἦ ὅσων ὀψις ἀκοὴ μάθησις, ταῦτα ἐγὼ προτιμέω

'Tis things visible, audible, perceptible that I prefer.

[Heracleitos]. Citation found in a paper written by G. Parisi [1].

One of the most important problems in contemporary particle physics is the computation of the internal structure of hadrons from first principles (i.e. from QUANTUM CHROMO-DYNAMICS (QCD)). Information on the internal structure of hadrons can be obtained through scattering experiments in particle accelerators. This information —the scattering cross section— can be expressed in terms of frame-independent STRUCTURE FUNCTIONS (containing the actual structural information) and frame-dependent, kinematic factors independent of the internal structure of the scattered object. For a precise definition see [2] and Chapter ⟨2⟩.

In the last two decades, a wealth of data on the structure of the proton has been collected in collider experiments. The largest amount of data gathered so far stems from DEEP(LY) INELASTIC SCATTERING (DIS) of leptons off the proton (or off hadrons in general). This scattering-process is particularly important in order to understand how a hadron is built up in terms of quarks

and gluons, its elementary constituents, since leptons —being point-like at all experimentally accessible scales— constitute a clean probe of hadrons. In the framework of Feynman’s parton model, structure functions may be interpreted as linear combinations of quark DISTRIBUTION FUNCTIONS as long as the resolution Q of the experiment is sufficiently large when compared to the mass M_H of the hadron. A QUARK DISTRIBUTION FUNCTION is the density of quarks with flavour i carrying a fraction x_B of the total momentum $\mathbf{P}$ of the hadron. Unfortunately, the computation of nuclear structure functions from first principles (i.e. from QUANTUM CHROMO-DYNAMICS, QCD) has not been achieved yet. Perturbative QCD merely allows to predict the dependence of the structure functions on Q whereas genuinely non-perturbative methods are called for in order to compute the x_B dependence of the structure functions.

2 Lattice Gauge Theory

Now, it is generally accepted that the most powerful non-perturbative method in QCD is LATTICE GAUGE THEORY(LGT) (or EUCLIDEAN LATTICE GAUGE THEORY, ELGT) [3, 4]. ELGT is so far the only technique capable of computing the hadronic masses directly from QCD *without* phenomenological assumptions [5, 6]. ELGT is based on path-integral quantisation with imaginary time. In this framework, renormalisable relativistic field theories appear as theories of statistical mechanics close to a critical point, a fact which makes them accessible to powerful Monte-Carlo methods. To render numerical computations feasible and free of infinities, continuous space-time *must* be replaced by a finite number of space-time points. This approximation —referred to as (lattice) REGULARISATION in the literature— partially destroys the Poincaré symmetry of the QCD Lagrangian (e.g. rotational

invariance and boost invariance). The Lagrangian is now invariant under the *discrete* symmetry group of the lattice. One can show, however, that Poincaré invariance is restored in the so-called CONTINUUM LIMIT, i.e. in the limit where the correlation lengths ξ of Green functions (measured in units of the lattice spacing) diverge. In praxis, the Poincaré invariance of the Lagrangian is *approximately* restored already for correlation lengths that are only slightly larger than the lattice spacing. While external symmetries are thus automatically restored in the continuum limit, this does not hold for internal symmetries such as gauge invariance. The defining Lagrangian should therefore, ideally, be exactly invariant under internal symmetries (unless one is able to disentangle physical states and spurious states). This requires the gauge-group to be compact and gauge-theories to be regularised on a space-time lattice rather than on a momentum-lattice. This point was first realised by Wilson in the case of QCD (WILSON ACTION [7]) and, earlier, by Wegner in the case of a discrete gauge theory [8].

In the last years, important progress has been made in ELGT. The first moments of nucleon structure-functions, for instance, can now be computed for the first time [9–12] [13–19]. These moments, however, are computed using the so-called QUENCHED APPROXIMATION. They represent, roughly, the moments of valence structure functions rather than the moments of full structure functions including sea quarks. For the latter ones, fermions have to be accounted for dynamically [4,20]. Computations beyond the quenched approximation are much more difficult to perform (i.e. they require much more CPU time) but there is no reason for why their computation should not be achieved in the near future. A further problem of ELGT is that the direct computation of structure functions or distribution functions would require the computation of a four-point function. A four-point Green function,

however, is extremely difficult to compute with the present lattice technology [21]. Moments of distribution functions, in contrast, as well as $1/Q$ corrections thereof, can be computed via three-point Green functions which are well under control [21].

Minkowsky space-time (as opposed to Euclidean space-time based on imaginary time) can be replaced by a lattice, too. In Minkowsky space-time, the path-integral formalism is less practical than in Euclidean space-time and it is more advantageous to work in the framework of the TRANSFER MATRIX formalism [3, 4] since the knowledge of all eigenvectors and eigenvalues of the transfer matrix $\mathbb{T}$ is equivalent to a complete solution of the theory.

The TRANSFER MATRIX is the generator of the discrete group of finite lattice translations in temporal direction: in the limit of vanishing temporal lattice spacing $a_t \rightarrow 0$ the transfer matrix

$$\mathbb{T} = \exp(-ia_t H) \approx 1 - ia_t H \quad (1.1)$$

can be replaced by a generator of the Poincaré group, the HAMILTONIAN, and we end up with the familiar Hamiltonian formulation of quantum mechanics. Yet choosing the temporal lattice spacing a_t smaller than the spatial lattice spacing a is not as innocent as it may seem. It necessitates, in principle, the introduction of additional relevant operators and coupling constants¹ into the action which are excluded by the symmetries of a symmetric lattice. Fortunately, however, the exclusion of these operators seems to be justified as ELGT calculations on anisotropic lattices seem to indicate [22–25].

The Hamiltonian which corresponds to the Wilson Lagrangian of lattice gauge theory is the so-called KOGUT-SUSSKIND HAMILTONIAN, first derived in [26]. **The aim of this thesis** is to explore the **Hamiltonian approach** towards relativistic field theories from the point of view of lattice

¹This is referred to as RENORMALISATION OF THE SPEED OF LIGHT

(gauge) theory. The Hamiltonian approach provides us with the advantage of the intuitive particle picture which is somewhat obscured in an imaginary-time formalism such as Euclidean lattice field theory. Once wave functions are computed, the computation of distribution functions is straightforward. A second advantage is that it is relatively easy to compute scattering observables such as structure functions (or other S-Matrix elements), once the eigenfunctions of the Hamiltonian are found.

3 A Brief Review of Hamiltonian Methods

Over recent years several researchers have explored Hamiltonian methods. Prominent examples are the work of Lüscher [27] and van Baal [28, 29] who have discovered that much physics of the low-lying QCD -spectrum, at least for small lattices, can be described by zero-momentum dynamics plus a suitable treatment of the remaining degrees of freedom. H. Kröger et al. used the Hamiltonian formalism in order to compute S-matrix elements [30–34]. But Hamiltonian methods have not been mainstream in the domain of non-perturbative methods. One reason for this is that the particle number is not conserved in relativistic QFTs: any interaction in relativistic quantum mechanics is capable of producing particle-anti-particle pairs. Ultra-relativistic objects such as the proton are thus complex mixtures of few-body and many-body physics: Even the vacuum has a non-vanishing density of gluons, quarks and anti-quarks. Accordingly, the vacuum contains an infinite number of virtual particles in the THERMODYNAMIC LIMIT, i.e. the limit where the lattice size becomes infinite; even the fluctuation of the particle number diverges.

In the applications of the Kogut-Susskind Hamiltonian to QCD [26], several groups have developed clever ways to take into account a large number of degrees of freedom, e.g., via the t -expansion method by Horn and co-

workers [35]. The $\exp[S]$ method, coming from nuclear physics, is a very effective method in order to deal with the volume divergences in the virtual particle number. A real breakthrough in the application of the $\exp[S]$ method to the Kogut-Susskind Hamiltonian of pure QCD has been achieved recently by [36–40]: glueball masses and string tension have been correctly estimated. Further Hamiltonian approaches are front-form quantisation [41] (see $\langle 3 \rangle$), and —quite recently— a Hamiltonian renormalisation group approach [42–44]. For a more thorough review of Hamiltonian lattice gauge theory we refer the reader to [37, 45].

4 Advantages of the Lorentz-Contraction: Proposal and Test of a New Technique

In this thesis, we present a new method which drastically simplifies the numerical diagonalisation of a relativistic lattice Hamiltonian. The inspiration to our method comes from Feynman’s PARTON MODEL. The parton model necessitates a fast-moving hadron rather than a hadron at rest in order for the distribution functions to be related to the structure functions in a simple way. We are able to show that, surprisingly, a hadronic state which moves sufficiently fast *relative to the lattice* can be dramatically simpler when compared to a bound state at rest. Part, but not all, of this simplicity stems from the fact that a fast, Lorentz-contracted object fits into smaller lattices as we shall see. We shall refer to a frame in which the hadron moves with large but *finite* momentum relative to the lattice as a LARGE MOMENTUM FRAME (LMF). The LMF must not be confused with another, similar frame, the INFINITE MOMENTUM FRAME (IMF), i.e. a frame wherein all particle masses can be neglected compared to the energies of these particles. While the IMF is admissible for some elementary perturbative calculations, the IMF can-

not be used, in general, for the non-perturbative computation of distribution functions on a *finite* lattice: We demonstrate this in Chapter [3](#). We argue that the IMF on a finite lattice is unphysical in general since it is incompatible with the SCALING WINDOW [\[4\]](#) of LGT. The LMF, in contrast, allows the limit of infinite momentum (relative to the lattice) to be approached on a finite lattice without leaving the scaling window of LGT. In Chapter [5](#) we illustrate this by practical computations. We apply the LMF technique to two models: quantum electrodynamics in $1 + 1$ space-time dimensions — referred to as massive SCHWINGER MODEL or QED($1 + 1$)— and the scalar ϕ^4 model in $3 + 1$ dimensions. It turns out that it is much simpler, in these models, to describe a physical particle that moves sufficiently fast relative to the lattice than to describe a particle at rest. We demonstrate, both theoretically and with practical examples, that a physical particle cannot move arbitrarily fast on a finite lattice, implying, in particular, that the IMF is *unphysical* (with few exceptions in perturbation theory or purely fermionic systems). We also demonstrate that the parton distribution functions receive significant contributions from the vacuum and that there is only one reference frame, the Breit frame, in which the vacuum contributions cancel entirely.

5 Does the Hamiltonian exist?

Begriffe und Begriffssysteme erhalten die Berechtigung nur dadurch, daß sie zum Überschauen von Erlebniskomplexen dienen; eine andere Legitimation gibt es für sie nicht. Es ist deshalb nach meiner Überzeugung einer der verderblichsten Taten der Philosophen, daß sie gewisse begriffliche Grundlagen der Naturwissenschaft aus dem der Kontrolle zugänglichen Gebiete des Empirsch-Zweckmäßigen in die *unangreifbare Höhe des Denknöthwendigen* (Apriorischen) versetzt haben.

[Albert Einstein: Grundzüge der Relativitätstheorie.]

Historically, during the 60's and 70's, quantum field theory (QFT) in general and Hamiltonian field theory in particular were considered to be amateurish: they were neglected in favour of the boot-strap programme, the hope of finding the S-matrix for the forces of nature from principles such as duality, analyticity, crossing-symmetry and the like. Many findings of this time, such as Regge theory and dispersion-relations remain relevant independently of the underlying theory whereas the ambitious boot-strap programme itself failed: QFT prevailed. The reason for the widespread mistrust of QFT in its earlier stages was the dominant *philosophy* that Poincaré symmetry had to be treated as an *exact* symmetry in any sensible computation. Discretisation of space-time — the modern LGT approach — was not yet seriously considered. Taking *exact* Poincaré invariance of the Lagrangian as an axiom combined with other physically motivated axioms, it can be shown in the framework of AXIOMATIC FIELD THEORY [46] that relativistic QFT is not well-defined except for non-interacting theories. This problem is solved in the modern approach which interprets renormalisable QFTs as systems close to a critical point. In the framework of LGT, fields are defined on a finite lattice replacing continuous space-time; an infinite number of effective field degrees of freedom is replaced by a finite number of degrees of freedom.

Poincaré invariance of the Lagrangian arises *dynamically*, i.e. it is restored to arbitrary accuracy when approaching the critical point. Even though some quantities, such as bare parameters or field fluctuations, diverge in the infinite volume limit, this does not constitute a problem in the lattice approach as these quantities are finite on any finite lattice.

6 Is Quantisation on Light-Like Quantisation Surfaces Viable?

Before the modern QFT philosophy was fully developed, the Hamiltonian formalism had been re-introduced in the form of the so-called FRONT-FORM (FF) quantisation [47]. This approach seemed to lack the "problems" that afflicted the usual or INSTANT FORM (IF) Hamiltonian formalism. In particular, the vacuum in this formalism seemed to be trivial and the infinite field fluctuations seemed to be absent. Some researchers went even so far as to claim that the FF approach was well defined due to the absence of infrared-singular field fluctuations whereas the IF was not. The FF approach is partially successful when applied to some theories in $1 + 1$ dimensions: It describes observables such as distribution functions and mass spectra of e.g. QUANTUM ELECTRODYNAMICS in two dimensions (QED(1+1)) [48] or QCD(1+1) with little numerical effort compared to LGT [49, 50]. Our approach can be seen as a generalisation of the FF approach in the sense that the approximations on which the IMF approach is based are considerably less severe than the (implicit!) approximations the FF is built upon: whenever the FF convincingly describes physics, so does the LMF— usually with more ease and more accuracy. The contrary does not hold true (with the possible exception of coincidences). There is, nonetheless, still a discussion going on whether the FF is an exact method equivalent to path-integral quantisation the same way the

instant-form quantisation is equivalent to path-integral quantisation. This question is discussed (in a *perturbative* framework) in Ref. [51, 52]. It is known that the FF retains only half of the field degrees of freedom that are necessary in the IF quantisation, since half of the equations of motion are constraints (i.e. they do not involve the FF-time) [41]. However, it is often argued that the "missing" degrees of freedom are only necessary in the IF. The reduction of the number of degrees of freedom is then considered to be a major advantage of the FF since this simplifies the FF Hamiltonian enormously when compared to an IF Hamiltonian. **It is therefore important to demonstrate —once and for all— that the FF is indeed an approximation rather than a rigorous way of quantising relativistic field theories.** This is done in Sec.⟨3⟩. The fact that the FF and the IF are not equivalent has important consequences for the interpretation of DIS experiments since Feynman's parton model is often interpreted in terms of FF distribution functions (in addition to other possibilities such as the IMF and the Breit frame). Sometimes it is even claimed that the FF is the *only* way to interpret the parton model; in particular, it is often claimed that distribution functions must be interpreted in terms of light-like correlation functions, which — if it were true — would necessitate the FF approach in order to properly interpret DIS. We demonstrate, however, that structure functions are related to distribution functions obtained by conventional IF quantisation rather than distribution functions obtained by FF quantisation. In particular, we demonstrate that distribution functions can be related to light-like correlation functions only if an unphysical frame is introduced.

We have also provided some intuitive examples illustrating the nature of the approximations that go with a FF quantisation. In order to show that the FF and the IF approaches are not equivalent *in general* it suffices to demon-

strate their inequivalence for one special field theory. We have therefore chosen one of the simplest field theories, the ϕ^4 theory to make our point. In particular, we were able to show for this theory that the so-called left-movers, i.e. the degrees of freedom missing in the FF are in a subtle way responsible for the crucial property of microcausality. We show that —contrary to the LMF approximation proposed in this thesis—, ***microcausality does not hold in the FF*** even in a non-interacting scalar field theory. It is well-known that *causality* of time-ordered propagators is hampered if they are derived via FF quantisation, i.e. waves with positive energies are not necessarily moved forward in (real) time. One usually argues that (a) this is no serious reason for abandoning the FF approach and (b) this defect can be repaired [53–55]. Violation of micro-causality, however, cannot be discarded so easily. Either one refrains from using light-like periodic boundary conditions —in which case FF quantisation is not defined— or else one is faced with *observable* unphysical ”predictions” such as time-travel and an infinite speed of light in one spatial direction. We also show that the left-movers make a substantial self-energy contribution to the mass-spectrum of interacting bosons whereas interacting fermions do not receive this contribution. In Sec.⟨4⟩ we take another simple example related to a non-interacting field theory where the FF is unable to reproduce the results of the IF: The (massless) Schwinger-model. We show that calculation of the mass-spectrum of the Schwinger-model is as simple in the FF as in the IF if the unphysical axial gauge is chosen. In the limit of small fermion masses, the LMF method reproduces results from chiral perturbation theory with ease whereas the FF needs an infinite lattice in order to do so.

7 Organisation

This thesis is organised as follows:

In Chapter **⟨2⟩** we compute some DIS structure functions in the IMPULSE APPROXIMATION (IA). We argue that this approximation cannot be frame-independent if the vacuum is non-trivial and that the Breit frame is the most advantageous choice of frame in the sense that the impulse approximation is most accurate in this frame. We demonstrate that distribution functions must be interpreted in terms of space-like correlation functions: these correlation functions become light-like if and only if an infinite, unphysical boost is performed which collapses all space-like and light-like quantities onto a light-front.

In Chapter **⟨3⟩** we demonstrate the non-equivalence (in general) of the FF and the IF formalism and the strongly unphysical character of light-like boundary conditions. We trace the problems of the FF to the need to introduce boundary conditions which — as we are able to show — provoke an unacceptable, complete breakdown of microcausality. We introduce the notion of KINEMATICAL EQUIVALENCE OF QUANTISATION-FRAMES and show that the so-called ε co-ordinates do not legitimate the notion of "closeness to the light-cone". Quantisation in these co-ordinates is a mere re-parametrisation of the IF for $\varepsilon \neq 0$ and equivalent to the FF for $\varepsilon = 0$. Expressed differently: quantisation in $\varepsilon \neq 0$ co-ordinates is IF quantisation in a more clumsy form and the limit $\varepsilon \rightarrow 0$ is not continuous in general.

We demonstrate that the IMF and the FF violate elementary requirements of lattice (gauge) theory. The IMF is in general unphysical. The FF is even more unphysical in that it is an effective theory to the IMF *only* for the classical, unquantised theory. In quantised form, the FF cannot be considered to be an effective theory to the IMF. There are exceptions to this

rule as QED(1 + 1), for instance. We explain under which circumstances the FF can be almost equivalent to the IF.

In Chapter [4](#) we use the exactly known solution to massless QED in two space-time dimensions in order to study the impact of various approximations. In particular we justify the assumptions that went into our numerical computations in Chapter [5](#).

In Chapter [5](#) we apply our LMF method to QED in two space-time dimensions. We demonstrate that the masses of the vector boson and of the scalar boson are accurately described for the whole range of the fermionic mass. In particular our method is able to reproduce the linear fermion-mass-dependence of the mass spectrum on a *finite* lattice whereas an infinite lattice is necessary in the FF or in the IMF.

In Chapter [6](#) we include already published papers on the four-dimensional φ^4 model wherein we compare our method with Ref. [\[56, 57\]](#) showing that we are able to describe reasonably well the critical properties and the critical line of the scalar model.

In Chapter [7](#) we draw a few conclusions.

8 Notation

We are using natural units $\hbar = c = 1$ throughout this thesis ($\hbar$ is Planck's constant, c is the velocity of light). Please note that we properly distinguish co-variant four-vectors x_μ and contra-variant four-vectors x^μ since we are using non-orthogonal frames in Chapter [3](#). Our notation will therefore be close to [\[58\]](#). Accordingly, if x denotes a four-vector, the expression x^2 does in general *not* mean $x \cdot x$ unless it is unambiguously clear from the context. We shall use the abbreviations $x^\pm \stackrel{def}{=} x^0 \pm x^3$, $x_\pm = \frac{1}{2}x^\mp$ (the so-called LIGHT-FRONT CO-ORDINATES or LIGHT-CONE CO-ORDINATES).

In every chapter except for Chapter **3** we use a diagonal metric g with $g_{11} = g_{22} = g_{33} = -g_{00} = -1$. In these chapters, we can replace $(x^\mu)^2$ by x_μ^2 since then $x_\mu = \pm x^\mu$. Four-vectors are represented in terms of their coordinates in the form $x = (x^0, x^1, x^2, x^3) = (x^0, \mathbf{x})$. Sometimes we label a four-vector $x = (x^0, \mathbf{x})$ or a tensor with a point in order to underline that $\mathbf{x}$ consists of contra-variant components.

We also have to distinguish the distribution $\delta^{(3)}(\mathbf{k}-\mathbf{k}')$ from the Kronecker symbol $\delta_{x,y} = 1$ or 0 . When working in a finite box with length $2L$ in three-direction and $2L_\perp$ in one- and two-direction (where the momenta $\mathbf{k}$ become discrete) it is explicitly assumed that the δ distribution becomes a function defined as

$$\delta^{(3)}(\mathbf{k} - \mathbf{k}') = \delta_{\mathbf{k},\mathbf{k}'} \left(\frac{L}{\pi}\right) \left(\frac{L_\perp}{\pi}\right)^2 \quad (1.2)$$

and $\int d^3x$ abridges

$$\int_{-L_\perp}^{L_\perp} \int_{-L_\perp}^{L_\perp} \int_{-L}^L dx^1 dx^2 dx^3 \quad . \quad (1.3)$$

Only if the momentum lattice spacing $\Delta k \stackrel{def}{=} \frac{\pi}{L}$ is one do $\delta_{\mathbf{k},\mathbf{k}'}$ and $\delta^{(3)}(\mathbf{k}-\mathbf{k}')$ coincide. It is convenient to define other quantities such as annihilation operators similarly such that $b_{\mathbf{k}}^\dagger |0\rangle$ ² is normalised to

$$\langle 0 | b_{\mathbf{k}} b_{\mathbf{k}'}^\dagger | 0 \rangle = \delta_{\mathbf{k},\mathbf{k}'} \quad (1.4)$$

whereas $b(\mathbf{k})$ is defined such as

$$\langle 0 | b(\mathbf{k}) b^\dagger(\mathbf{k}') | 0 \rangle = \delta^{(3)}(\mathbf{k} - \mathbf{k}') \quad (1.5)$$

In the literature, the HELICITY $\mathcal{H} = \pm \frac{1}{2}$ of a proton is often defined to take on the values ± 1 . In order to facilitate comparisons with both notations, we shall write helicities with a bracket $(2\mathcal{H}) = \pm 1$.

² $|0\rangle$ is the perturbative vacuum

As this thesis uses the formalism of several, disparate branches of physics such as quantum field theory, scattering theory, lattice gauge theory, numerics, many-body theory, solid state physics, constraint quantisation, group theory and some notation from general relativity we are faced with the problem that one symbol may mean different things in these domains. In order to avoid changing familiar symbols we distinguish these symbols through four different fonts: *italic*, **sans serif**, *CALIGRAPHIC* and **BLACKBOARD** rather than through the introduction of entirely new symbols, a procedure which conserves the familiarity of symbols and reduces the ambiguity at a time. If it is not clear from the context that a quantity $\mathcal{O}$ is an OPERATOR rather than a number then this operator $\hat{\mathcal{O}}$ is identified as such with a hat. The transposition of a matrix T is denoted $T^\top$, its hermitian conjugate $T^\dagger$. NORMAL ORDERING of an operator $\hat{\mathcal{O}}$ is written as $:\hat{\mathcal{O}}:$. We shall be using the HEISENBERG PICTURE of quantum mechanics throughout this thesis. For the reader's convenience, most symbols that have been used are listed in a separate index at the end of this thesis. The index is alphabetically ordered except for operators (e.g. $\otimes$) which appear before the letter *a*. Greek letters are ordered according to the first letter of their Latin transcription, e.g. $\alpha = \textit{alpha}$ is treated as *a*, $\omega = \textit{omega}$ is treated as *o*.

9 Methodology

The numerical part was done using a combination of C++, UNIX and MATHEMATICA. A programme for the algebraical manipulation of quantised fields and their creation operators was designed and matrix elements of the Hamiltonian were algebraically computed using this programme and automatically translated into C++ code in the case of the three-dimensional scalar model or into MATHEMATICA code in the case of one-dimensional QED. The energy

spectra were obtained through a numerical diagonalisation of the thus computed matrices. Since small matrices already sufficed in order to reproduce very accurate results, the numerical diagonalisations could be effortlessly performed with MATHEMATICA routines.

Chapter 2

Structure Functions as Short-Distance Physics

Introduction

DEEP INELASTIC SCATTERING of leptons off a hadron provides us with information on the internal structure of hadrons. The incoming lepton with four-momentum k scatters off the hadron with four-momentum $P = (E, \mathbf{P})$, where $E = P^0 = \sqrt{M^2 + \mathbf{P}^2}$ denotes the energy and M the mass of the hadron. After the scattering process, the four-momentum k' of the lepton is measured. We shall only consider INCLUSIVE SCATTERING¹. We are not interested in the exact state $|P', X\rangle$ of the debris of the hadron *after* the scattering process. The inclusive differential scattering cross-section $d\sigma$ is proportional to the contraction $W^{\mu\nu}l_{\mu\nu}$ of the hadronic tensor $W^{\mu\nu}$ and the leptonic tensor $l^{\mu\nu}$ if first order perturbation theory is valid, $q \stackrel{def}{=} k - k'$ is the (space-like) momentum of the exchanged virtual photon. While the leptonic tensor $l^{\mu\nu}$ associated with the incoming lepton may be calculated

¹Inclusive scattering means that the final state of the proton is not measured.

	Polarisation	Pauli-Lubansky	Helicity	Spin/Transversity
Hadronic	$\overset{\uparrow}{S} = (\overset{\uparrow}{S}^0, \overset{\uparrow}{\mathbf{S}})$	$S = (S^0, \mathbf{S})$	$\mathcal{H}$	$\mathcal{T}$
Partonic	$\overset{\uparrow}{s} = (\overset{\uparrow}{s}^0, \overset{\uparrow}{\mathbf{s}})$	$s_{\text{P.L.}} = (s^0_{\text{P.L.}}, \mathbf{s}_{\text{P.L.}})$	s	—

Table 2.1: Overview of polarisation vectors

perturbatively, this is not the case for the hadronic tensor

$$W^{\mu\nu} = (2\pi)^6 E \int \frac{d^4 y}{(2\pi)^4} e^{+iq \cdot y} \langle PS | : [\hat{j}^\mu(y), \hat{j}^\nu(0)] : | PS \rangle \quad (2.1)$$

which contains the information on the internal structure of the hadron. In order to compute this tensor theoretically, knowledge of the hadronic wave function $|PS\rangle$ with normalisation

$$\langle PS | P'S \rangle = \delta^{(3)}(\mathbf{P} - \mathbf{P}') \quad (2.2)$$

is required. Here, $\hat{j}^\mu(x)$ stands for the hadronic CURRENT OPERATOR the definition of which will be detailed below. The four-vector S is the PAULI-LUBANSKY FOUR-VECTOR [58, 59] —a relativistic generalisation of the spin three-vector— with the properties

$$S \cdot P = S^0 P^0 - \mathbf{S} \cdot \mathbf{P} = 0 \text{ and } S \cdot S = -M^2 \quad (2.3)$$

which characterises the spin of the hadron as follows: One can always find a Lorentz frame in which $P = (M, 0, 0, 0)$ and $S = (0, \mathbf{S})$. In *this* frame, $M^{-1}\mathbf{S} = \mathbf{J}$ coincides with the TOTAL ANGULAR MOMENTUM THREE-VECTOR $\mathbf{J}$ which, in turn, coincides with the spin. Without loss of generality, we shall henceforth assume that the hadron moves RIGHT,

$$P = (P^0, P^1, P^2, P^3) = (E, \mathbf{P}) = (E, 0, 0, P^3) \quad M^2 \stackrel{def}{=} P \cdot P \quad (2.4)$$

i.e. the hadron moves in the positive 3-direction $P^3 > 0$. The direction of $\mathbf{P}$ defines the LONGITUDINAL direction throughout this thesis. If the hadron

is LONGITUDINALLY POLARISED with HELICITY

$$\mathcal{H} \stackrel{def}{=} \frac{\mathbf{J} \cdot \mathbf{P}}{|\mathbf{P}|} = \frac{1}{2} \frac{\mathbf{S} \cdot \mathbf{P}}{E|\mathbf{P}|} = \frac{1}{2} \frac{S^0}{|\mathbf{P}|} = \pm \frac{1}{2} \quad (2.5)$$

then $\mathbf{S}$ is collinear with the spin vector and with $\mathbf{P}$

$$S = (S^0, 0, 0, S^3) = (2\mathcal{H})(P^3, 0, 0, E) = (2\mathcal{H})M\overset{\uparrow}{S}_H \quad (2.6)$$

where $\overset{\uparrow}{S}_H = \frac{1}{M}(P^3, 0, 0, E)$ is the HELICITY POLARISATION AXIS. If the polarisation direction is perpendicular to the momentum of the hadron, one says that the hadron is TRANSVERSELY POLARISED. In this case, the Pauli-Lubansky four-vector reads

$$S = (0, S^1, 0, 0) = (2\mathcal{T})(0, M, 0, 0) = (2\mathcal{T})M\overset{\uparrow}{S}_T \quad (2.7)$$

if one (arbitrarily) chooses the 1-direction as polarisation axis. Here, $\overset{\uparrow}{S}_T = (0, M, 0, 0)$ is the TRANSVERSE POLARISATION AXIS and $\mathcal{T} = \pm \frac{1}{2}$ is the TRANSVERSE SPIN i.e. the quantised spin component in this direction. The reader might want to notice that some authors normalise $-S \cdot S$ to one. Furthermore, a covariant normalisation

$$\langle PS, \text{cov} | P' S, \text{cov} \rangle = 2E(2\pi)^3 \delta^{(3)}(\mathbf{P} - \mathbf{P}') \quad (2.8)$$

of the hadron state $|PS\rangle = |PS, \text{cov}\rangle / \sqrt{2E(2\pi)^3}$ is often used in the literature in which case the hadronic tensor reads

$$W^{\mu\nu} = \frac{1}{4\pi} \int d^4y e^{+iq \cdot y} \langle PS, \text{cov} | : [\hat{j}^\mu(y), \hat{j}^\nu(0)] : | PS, \text{cov} \rangle \quad . \quad (2.9)$$

The hadronic tensor W may be decomposed into a symmetric(S) part

$$W_S^{\mu\nu} \stackrel{def}{=} \frac{1}{2}(W^{\mu\nu} + W^{\nu\mu}) \quad (2.10)$$

independent of polarisation effects and an anti-symmetric(A) part

$$W_A^{\mu\nu} \stackrel{def}{=} \frac{1}{2}(W^{\mu\nu} - W^{\nu\mu}) \quad (2.11)$$

containing the polarisation effects. Both parts may, in turn, be written as a linear combination of Lorentz-scalar, dimensionless STRUCTURE FUNCTIONS [2, 60] $F_1(x_B, Q)$, $F_2(x_B, Q)$, $g_1(x_B, Q)$, $g_2(x_B, Q)$ which contain the structural information proper

$$\begin{aligned}
W_S^{\mu\nu} &= (-g^{\mu\nu} - \frac{q^\mu q^\nu}{Q^2}) F_1(x_B, Q) \\
&+ (P^\mu + \frac{P \cdot q}{Q^2} q^\mu)(P^\nu + \frac{P \cdot q}{Q^2} q^\nu) \frac{F_2(x_B, Q^2)}{P \cdot q} \\
&= (-g^{\mu\nu} - \frac{q^\mu q^\nu}{Q^2}) F_1(x_B, Q) \\
&+ (P^\mu + \frac{1}{2x_B} q^\mu)(P^\nu + \frac{1}{2x_B} q^\nu) \frac{2x_B F_2(x_B, Q^2)}{Q^2}
\end{aligned} \tag{2.12}$$

$$\begin{aligned}
W_A^{\mu\nu} &= -i\epsilon^{\mu\nu\alpha\beta} \frac{q_\alpha}{(P \cdot q)^2} \times \\
&\left[(g_1(x_B, Q^2) + g_2(x_B, Q^2)) (P \cdot q) S_\beta - g_2(x_B, Q^2)(q \cdot S) P_\beta \right]
\end{aligned} \tag{2.13}$$

and Lorentz-covariant kinematic tensors which do not depend on the structure of the hadron. The structure functions depend on the two invariants MOMENTUM TRANSFER (or scattering RESOLUTION)

$$Q \stackrel{def}{=} \sqrt{-q \cdot q} \tag{2.14}$$

and on the BJORKEN SCALING VARIABLE

$$x_B \stackrel{def}{=} \frac{Q^2}{2P \cdot q} \tag{2.15}$$

where $q \stackrel{def}{=} (P' - P)$. This is so because there are only two scalar quantities that can be formed from the kinematic four-vectors P and P' characterising the hadron. Hadronic tensors computed on a lattice, however, can be expected to depend additionally (weakly) on $\mathbf{P}$ because a lattice breaks Poincaré symmetry. Frame-dependent approximations also introduce $\mathbf{P}$ into the hadronic tensor. Of course, this dependence has to disappear in the

continuum limit— as long as the momentum $\mathbf{P}$ lies in a domain where the approximations are accurate. We have tacitly assumed here, that parity is an exact symmetry. In other words, we do not consider WEAK INTERACTIONS (which are not parity invariant). If we had taken weak interactions into account, more structure functions would have appeared [2]. Here, we shall only consider electromagnetic interactions between the hadron and the probing lepton.

1 The Hadronic Tensor and Structure Functions

In the BREIT FRAME, the hadronic tensor takes on a particularly simple form. In this frame, there is a particularly simple relation between structure functions and the hadronic tensor. The BREIT FRAME is defined as the frame where $\mathbf{q}$ is collinear with $\mathbf{P}$ and q is AT REST $q^0 \equiv 0$. Frames with $q^0 = 0$ are the *only* frames where Q corresponds to the resolution ability of the experiment. For in frames where $q^0 \neq 0$, the wave-length of the exchanged virtual photon is not $\frac{2\pi}{Q}$ but rather $\frac{2\pi}{|q^3|} = \frac{2\pi}{\sqrt{(q_0)^2 + Q^2}}$. In the Breit frame, the hadronic tensor reduces to

$$W(x_B, Q; \mathcal{H}) = \begin{pmatrix} W^{00} & 0 & 0 & 0 \\ 0 & W^{11} & W^{12} & 0 \\ 0 & W^{21} & W^{22} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (2.16)$$

with

$$W^{12}(x_B, Q^2; \mathcal{H}) = i2\mathcal{H}(g_1 - g_2 \frac{M^2}{(P^3)^2}) \approx i2\mathcal{H}g_1 \quad (2.17)$$

if the hadron is longitudinally polarised and to

$$W(x_B, Q; \mathcal{T}) = \begin{pmatrix} W^{00} & 0 & W^{02} & 0 \\ 0 & W^{11} & 0 & 0 \\ W^{20} & 0 & W^{22} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (2.18)$$

with

$$W^{02} = -W^{20} = -i2\mathcal{T} \frac{M}{P^3} (g_1 + g_2) \approx 0 \quad (2.19)$$

if the hadron is transversely polarised. The diagonal components read

$$W^{00} = \frac{1}{2}F_L = -F_1 + \frac{2x_B E^2}{Q^2} F_2 = -F_1 + \frac{E^2}{P^3 Q} F_2 \quad (2.20)$$

$$= -F_1 + (1 + (\frac{2x_B M}{Q})^2) \frac{1}{2x_B} F_2 \approx -F_1 + \frac{1}{2x_B} F_2$$

$$W^{11} = W^{22} = F_1 \quad (2.21)$$

independently of the polarisation. Equation **(2.21)** holds in any frame where both $\mathbf{P}$ and $\mathbf{q}$ point in 3-direction whereas $W^{33} = 0$ and Eq. **(2.20)** only hold in the Breit frame ($q^0 = 0$). For later purposes it is useful to define the total momentum P^3 and energy P^0 in the Breit frame as

$$P_B = \frac{Q}{2x_B} \text{ and } E_B = \sqrt{M^2 + P_B^2} \quad . \quad (2.22)$$

2 The Hadronic Tensor: Formal Definition

The current operators

$$\hat{j}^\mu(x^0, \mathbf{x}) = U(x^0) \hat{j}^\mu(0, \mathbf{x}) U^\dagger(x^0) \quad (2.23)$$

appearing in the definition **(2.1)** of the hadronic tensor are in general very complicated, interaction-dependent objects for $x^0 \neq 0$ as the TIME-EVOLUTION

OPERATOR

$$U(x^0) \stackrel{\text{def}}{=} e^{-iHx^0} \stackrel{\text{def}}{=} e^{-iH_0x^0 - iH_Ix^0} \quad (2.24)$$

depends on the interaction-part H_I of the lattice-regularised Hamiltonian of QCD (or another quantum field theory). H_0 stands for the kinetic energy. In what follows, we are not interested in the exact form of H and the problems involved in its definition. It suffices to know that H_{QCD} is well defined, when constructed on a lattice in configuration space [4, 26] using Wilson's compact lattice variables. As to a Hamiltonian on a *momentum* lattice, we remark that such a Hamiltonian would require the use of non-compact gauge fields A^μ which necessitate, in turn, complete gauge-fixing with all its complications such as the Gribov horizon etc. It may or may not be possible to write down such an object. In what follows, we shall use the momentum-space formalism usually employed in DIS for the practical reason that the corresponding expressions in configuration space would be by far more complicated and less intuitive.

Equation (2.24) may be written as a perturbative series. The conditions under which this is a good approximation will be discussed later. Feynman's parton model is based on the IMPULSE APPROXIMATION (IA) i.e. on zeroth order perturbation theory. In this case $U(x^0) \stackrel{\text{def}}{=} e^{-iHx^0}$ and the currents can be represented in terms of free fermion fields

$$\psi(x) = e^{-iH_0x^0} \psi(0, \mathbf{x}) e^{+iH_0x^0} = \sum_s \int d^R k (u_s(\mathbf{k}) e^{-ik \cdot x} b_s(\mathbf{k}) + v_s(\mathbf{k}) e^{+ik \cdot x} d_s^\dagger(\mathbf{k})) \quad (2.25)$$

obeying the anti-commutation relations

$$\{\psi(t, \mathbf{x}), \psi^\dagger(t, \mathbf{y})\} = \int d^{[R]} k \sum_s (u_s \otimes u_s^\dagger + v_s^\dagger \otimes v_s) = \mathbb{1} \delta^{(3)}(\mathbf{x} - \mathbf{y}) \quad . \quad (2.26)$$

The symbol $\mathbb{1}$ is the (4×4) unity matrix in spinor space. For convenience we have introduced relativistic (R) integration measures

$$d^R k \stackrel{def}{=} \frac{d^3 k}{\sqrt{(2\pi)^3 2\omega(\mathbf{k})}} \quad (2.27)$$

$$d^{[R]} k \stackrel{def}{=} \frac{d^3 k}{(2\pi)^3 2\omega(\mathbf{k})} \quad (2.28)$$

similar to Ref. [58]. The fermion fields are expressed in terms of the spinors u and v which are normalised to

$$\bar{u}_s u_{s'} = +2m\delta_{ss'} \quad (2.29)$$

$$\bar{v}_s v_{s'} = -2m\delta_{ss'}. \quad (2.30)$$

and in terms of the fermionic creation and annihilation operators obeying the standard anti-commutation relations

$$\{b_s(\mathbf{k}), b_{s'}^\dagger(\mathbf{k}')\} = \delta^{(3)}(\mathbf{k} - \mathbf{k}')\delta_{ss'} \quad (2.31)$$

$$\{d_s(\mathbf{k}), d_{s'}^\dagger(\mathbf{k}')\} = \delta^{(3)}(\mathbf{k} - \mathbf{k}')\delta_{ss'} \quad (2.32)$$

where $s = \pm\frac{1}{2}$ designates the PARTON HELICITY. The spinors obey the completeness relations

$$u_s(\mathbf{k}) \otimes \bar{u}_s(\mathbf{k}) = (\not{k} + m) \frac{1 + (2s)\gamma_5 \not{\hat{s}}}{2} \xrightarrow{m \rightarrow 0} \not{k} \frac{1 + (2s)\gamma_5}{2} \quad (2.33)$$

and

$$v_s(\mathbf{k}) \otimes \bar{v}_s(\mathbf{k}) = (\not{k} - m) \frac{1 + (2s)\gamma_5 \not{\hat{s}}}{2} \xrightarrow{m \rightarrow 0} \not{k} \frac{1 - (2s)\gamma_5}{2} \quad (2.34)$$

where

$$\hat{s}(\mathbf{k}) = \frac{\omega(\mathbf{k})}{m} \left(\frac{|\mathbf{k}|}{\omega(\mathbf{k})}, \frac{\mathbf{k}}{|\mathbf{k}|} \right) \quad (2.35)$$

defines, analogously to $\overset{\uparrow}{S}_H$, the HELICITY-DIRECTION (or SPIN QUANTISATION AXIS) associated with the fermion. The Pauli-Lubansky four-vector of the fermion

$$s_{\text{P.L.}} \stackrel{\text{def}}{=} (2s)m\overset{\uparrow}{s} \quad (2.36)$$

is defined such that it is normalised to $-m^2$. If transverse momenta can be neglected—which is assumed in the framework of the parton model—it would be equally convenient to choose a spin polarisation

$$\overset{\uparrow}{s}_{\text{spin}} = \frac{1}{m_{\perp}}(k^3, 0, 0, k^0) \quad \text{with} \quad m_{\perp} \stackrel{\text{def}}{=} \sqrt{m^2 + k_{\perp}^2} \quad (2.37)$$

instead of a helicity polarisation. There are two ways to compute the hadronic tensor in the impulse approximation. One may expand the field commutator in terms of bilinears—which is usually done [60]—or one may do everything on the level of the creation and annihilation operators, which we shall do herein since we consider it to be closer to intuition. We note—at this place—that the fermionic field $\psi(x)$ is not gauge invariant because a gauge-transformation $\psi \rightarrow \psi e^{i\alpha}$ changes its phase. If the fermions move in a gluonic background, we should replace the local fermion field $\psi(x)$ with the non-local field $\psi_A(y) \stackrel{\text{def}}{=} U_A(y)\psi(y)$ where $U_A(y)$ (with $U_A(0) = \mathbb{1}$) is a non-local string of gauge fields [60] connecting the points 0 and y .

3 Structure Functions and Distribution Functions

The current commutator $[j^{\mu}(y), j^{\nu}(0)]$ appearing in the hadronic tensor involves a product of four fermionic fields ψ , each of which is a sum of quark operators $b, b^{\dagger}$ and anti-quark operators $d^{\dagger}, d$. Consequently, the current-commutator consists of $2^4 = 16$ terms with all possible combinations of

particles and anti-particles. The hadronic tensor may thus be written in the form

$$W = \frac{(2\pi)^6 E}{(2\pi)^4} \int d^4 y \, d^R k \, d^R k' \, d^R l \, d^R l' \sum_{r=1}^{16} T_r^A T_r^B T_r^C \quad (2.38)$$

where $T_r^{A/B/C}$ are constants

$$T_r^A = T_r^{A\mu\nu}(s, s', \sigma, \sigma'; \mathbf{k}, \mathbf{k}', \mathbf{l}, \mathbf{l}') \quad (2.39)$$

whose dependence on Lorentz indices, momenta and spin we suppress in order to avoid awkward expressions. We shall write the T -symbols as 16-dimensional vectors (since the index r runs over 16 values)

$$T^B \stackrel{def}{=} \langle \hat{T}^B \rangle \stackrel{def}{=} \langle PS | \hat{T}^B | PS \rangle \quad , \quad (2.40)$$

$$T^A = \begin{pmatrix} \bar{u}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{u}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu u_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{u}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu u_{\sigma'}(\mathbf{l}') \\ \bar{v}_s(\mathbf{k})\gamma^\mu v_{s'}(\mathbf{k}') \cdot \bar{v}_\sigma(\mathbf{l})\gamma^\nu v_{\sigma'}(\mathbf{l}') \end{pmatrix} \quad , \quad (2.41)$$

$$\hat{T}^B = \begin{pmatrix} [b_s^\dagger(\mathbf{k})b_{s'}(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})b_{s'}(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})b_{s'}(\mathbf{k}'), d_\sigma(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})b_{s'}(\mathbf{k}'), d_\sigma(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ \\ [b_s^\dagger(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), d_\sigma(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [b_s^\dagger(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), d_\sigma(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ \\ [d_s(\mathbf{k})b_{s'}(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [d_s(\mathbf{k})b_{s'}(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ [d_s(\mathbf{k})b_{s'}(\mathbf{k}'), d_\sigma(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [d_s(\mathbf{k})b_{s'}(\mathbf{k}'), d_\sigma(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ \\ [d_s(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [d_s(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \\ [d_s(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), d_\sigma(\mathbf{l})b_{\sigma'}(\mathbf{l}')] \\ [d_s(\mathbf{k})d_{s'}^\dagger(\mathbf{k}'), d_\sigma(\mathbf{l})d_{\sigma'}^\dagger(\mathbf{l}')] \end{pmatrix}, \quad T^C = \begin{pmatrix} e^{i(k \cdot y - k' \cdot y + qy)} \\ e^{i(k \cdot y - k' \cdot y + qy)} \\ e^{i(k \cdot y - k' \cdot y + qy)} \\ e^{i(k \cdot y - k' \cdot y + qy)} \\ \\ e^{i(k \cdot y + k' \cdot y + qy)} \\ e^{i(k \cdot y + k' \cdot y + qy)} \\ e^{i(k \cdot y + k' \cdot y + qy)} \\ e^{i(k \cdot y + k' \cdot y + qy)} \\ \\ e^{i(-k \cdot y - k' \cdot y + qy)} \\ e^{i(-k \cdot y - k' \cdot y + qy)} \\ e^{i(-k \cdot y - k' \cdot y + qy)} \\ e^{i(-k \cdot y - k' \cdot y + qy)} \\ \\ e^{i(-k \cdot y + k' \cdot y + qy)} \\ e^{i(-k \cdot y + k' \cdot y + qy)} \\ e^{i(-k \cdot y + k' \cdot y + qy)} \\ e^{i(-k \cdot y + k' \cdot y + qy)} \end{pmatrix}, \quad (2.42)$$

Performing the integral $\int d^4y$ in (2.38) leaves us with a lower-dimensional integral

$$(2\pi)^6 E \int d^R k \, d^R k' \, d^R l \, d^R l' \sum_{s, s', \sigma, \sigma'}$$

over momentum conserving δ distributions

$$\tilde{T}^C \stackrel{def}{=} \int \frac{d^4 y}{(2\pi)^4} T^C = \begin{pmatrix} \delta^{(4)}(k - k' + q) \\ \delta^{(4)}(k - k' + q) \\ \delta^{(4)}(k - k' + q) \\ \delta^{(4)}(k - k' + q) \\ \delta^{(4)}(k + k' + q) \\ \delta^{(4)}(k + k' + q) \\ \delta^{(4)}(k + k' + q) \\ \delta^{(4)}(k + k' + q) \\ \delta^{(4)}(-k - k' + q) \\ \delta^{(4)}(-k - k' + q) \\ \delta^{(4)}(-k - k' + q) \\ \delta^{(4)}(-k - k' + q) \\ \delta^{(4)}(-k + k' + q) \\ \delta^{(4)}(-k + k' + q) \\ \delta^{(4)}(-k + k' + q) \\ \delta^{(4)}(-k + k' + q) \end{pmatrix}. \quad (2.43)$$

Expanding the commutators and sandwiching them with the vector $|PS\rangle$ yields

$$T^B = \begin{pmatrix} \langle b_s^\dagger(\mathbf{k})b_{\sigma'}(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & + \langle b_{s'}(\mathbf{k}')b_\sigma^\dagger(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\ \langle b_s^\dagger(\mathbf{k})d_{\sigma'}^\dagger(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & \\ \langle b_{s'}(\mathbf{k}')d_\sigma(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} & \\ 0 & \\ \langle d_{s'}^\dagger(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l})\delta_{s\sigma'} & \\ 0 & \\ \langle b_s^\dagger(\mathbf{k})b_{\sigma'}(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & + \langle d_{s'}^\dagger(\mathbf{k}')d_\sigma(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\ \langle b_s^\dagger(\mathbf{k})d_{\sigma'}^\dagger(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & \\ \langle d_s(\mathbf{k})b_{\sigma'}(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & \\ \langle d_s(\mathbf{k})d_{\sigma'}^\dagger(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & + \langle b_{s'}(\mathbf{k}'), b_\sigma^\dagger(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\ 0 & \\ \langle b_{s'}(\mathbf{k}'), d_\sigma(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} & \\ 0 & \\ \langle d_{s'}^\dagger(\mathbf{k}')b_\sigma^\dagger(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} & \\ \langle d_s(\mathbf{k})b_{\sigma'}(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & \\ \langle d_s(\mathbf{k})d_{\sigma'}^\dagger(\mathbf{l}') \rangle > \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} & + \langle d_{s'}^\dagger(\mathbf{k}')d_\sigma(\mathbf{l}) \rangle > \delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \end{pmatrix}$$

which, in turn, simplifies to

$$\begin{pmatrix}
[f_{s,\sigma'}(\mathbf{k}) - f_{\sigma,s'}(\mathbf{l})] & \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s'\sigma} \\
-\Xi_{\sigma',s}^*(\mathbf{l}') & \delta^{(3)}(\mathbf{k} + \mathbf{l}')\delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} \\
\Xi_{\sigma,s'}(\mathbf{l}) & \delta^{(3)}(\mathbf{k}' + \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\
0 & \\
\Xi_{s',\sigma}^*(\mathbf{k}') & \delta^{(3)}(\mathbf{k}' + \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l})\delta_{s\sigma'} \\
0 & \\
[f_{s,\sigma'}(\mathbf{k}) + \bar{f}_{s',\sigma}(\mathbf{k}')] & \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s'\sigma} \\
-\Xi_{\sigma',s}^*(\mathbf{l}') & \delta^{(3)}(\mathbf{k} + \mathbf{l}')\delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} \\
-\Xi_{s,\sigma'}(\mathbf{k}) & \delta^{(3)}(\mathbf{k} + \mathbf{l}')\delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} \\
[-\bar{f}_s(\mathbf{k}) - f_{s'}(\mathbf{k}')] & \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s'\sigma} \\
0 & \\
+\Xi_{\sigma,s'}(\mathbf{l}) & \delta^{(3)}(\mathbf{k}' + \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\
0 & \\
\Xi_{s',\sigma}^*(\mathbf{k}') & \delta^{(3)}(\mathbf{k}' + \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s\sigma'} \\
-\Xi_{s,\sigma'}(\mathbf{k}) & \delta^{(3)}(\mathbf{k} + \mathbf{l}')\delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta_{s'\sigma} \\
[\bar{f}_{\sigma's}(\mathbf{k}') - \bar{f}_{s'\sigma}(\mathbf{l}')] & \delta^{(3)}(\mathbf{k}' - \mathbf{l})\delta^{(3)}(\mathbf{k} - \mathbf{l}')\delta_{s'\sigma}
\end{pmatrix} \quad (2.44)$$

if the expectation values are replaced by what we shall call RAW DISTRIBUTION FUNCTIONS f_s

$$\langle PS | b_s^\dagger(\mathbf{k}) b_{\sigma'}(\mathbf{l}') | PS \rangle = f_{s,\sigma'}(\mathbf{k}, \mathbf{P}, S) \delta^{(3)}(\mathbf{k} - \mathbf{l}') \quad (2.45)$$

$$\langle PS | d_s^\dagger(\mathbf{k}) d_{\sigma'}(\mathbf{l}') | PS \rangle = \bar{f}_{s,\sigma'}(\mathbf{k}, \mathbf{P}, S) \delta^{(3)}(\mathbf{k} - \mathbf{l}') \quad (2.46)$$

$$f_s(\mathbf{k}, \mathbf{P}, S) \stackrel{def}{=} f_{s,s}(\mathbf{k}, \mathbf{P}, S) \quad (2.47)$$

$$\bar{f}_s(\mathbf{k}, \mathbf{P}, S) \stackrel{def}{=} \bar{f}_{s,s}(\mathbf{k}, \mathbf{P}, S) \quad (2.48)$$

and RAW PAIRING FUNCTIONS $\Xi_{s,\sigma}$

$$\langle PS | d_s(\mathbf{k}) b_{\sigma'}(\mathbf{l}') | PS \rangle = -\Xi_{s,\sigma'}(\mathbf{k}, \mathbf{P}, S) \delta^{(3)}(\mathbf{k} + \mathbf{l}') \quad (2.49)$$

$$\langle PS | b_s^\dagger(\mathbf{k}) d_{\sigma'}^\dagger(\mathbf{l}') | PS \rangle = -\Xi_{\sigma',s}^*(\mathbf{l}', \mathbf{P}, S) \delta^{(3)}(\mathbf{k} + \mathbf{l}') \quad (2.50)$$

$$\langle PS | b_s(\mathbf{k}) d_{\sigma'}(\mathbf{l}') | PS \rangle = +\Xi_{\sigma',s}(\mathbf{l}', \mathbf{P}, S) \delta^{(3)}(\mathbf{k} + \mathbf{l}') \quad (2.51)$$

$$\langle PS | d_s^\dagger(\mathbf{k}) b_{\sigma'}^\dagger(\mathbf{l}') | PS \rangle = +\Xi_{s,\sigma'}^*(\mathbf{k}, \mathbf{P}, S) \delta^{(3)}(\mathbf{k} + \mathbf{l}') \quad (2.52)$$

in order to distinguish them from what is called PARTON DISTRIBUTION functions in the literature.

The hadronic spin does not coincide with the total angular momentum $\mathbf{J}$ if the spin quantisation axis $\hat{\mathbf{S}}_H$ is not collinear to the hadron momentum $\mathbf{P}$ — in the case of transverse polarisation $\hat{S}_T$ for instance. In such cases, the spin is no longer kinematical. Up to now we did not use any particular photon momentum. Now we assume that we are in the Breit-frame, i.e. $q = (0, 0, 0, -Q)$. In this case, the delta-distribution (first four terms)

$$\begin{aligned} \delta(k' - k - q) &= \delta(\omega(\mathbf{k}') - \omega(\mathbf{k})) \delta(k^{3'} - k^3 + Q) \delta^{(2)}(\mathbf{k}^{\perp'} - \mathbf{k}^\perp) \\ &= \frac{\omega(\mathbf{k}_Q)}{Q} \delta(k^3 - Q/2) \delta(k^{3'} + Q/2) \delta^{(2)}(\mathbf{k}^{\perp'} - \mathbf{k}^\perp) \end{aligned} \quad (2.53)$$

can be expressed in terms of the vectors

$$\mathbf{k}_Q \stackrel{def}{=} (k^1, k^2, Q/2) \quad (2.54)$$

$$\mathbf{k}_{-Q} \stackrel{def}{=} (k^1, k^2, -Q/2) \quad (2.55)$$

since k and k' are on the energy shell due to the impulse approximation. In frames with $\mathbf{P} \parallel \mathbf{q}$ but $q^0 \neq 0$ we would have to replace k_Q^3 by

$$k_r^3 \stackrel{def}{=} -\frac{q^3}{2} - \frac{q^0}{Q} \sqrt{m^2 + (Q/2)^2} \approx -\frac{q^0 + q^3}{2} = -q_- \quad (2.56)$$

and k_{-Q}^3 by

$$k_l^3 \stackrel{def}{=} \frac{q^3}{2} - \frac{q^0}{Q} \sqrt{m^2 + (Q/2)^2} \approx -\frac{q^0 - q^3}{2} = -q_+ = -\frac{(Q/2)^2}{k_r^3} \quad (2.57)$$

and modify the weights $\omega(k_Q)/Q$ as well. Using these expressions, a six-fold integral of the form

$$\begin{aligned} &(2\pi)^6 \int d^R k d^R k' d^R l d^R l' \delta^4(k' - k + q) \delta^{(3)}(\mathbf{k} - \mathbf{l}') \delta^{(3)}(\mathbf{k}' - \mathbf{l}) T^{AB}(\mathbf{k}, \mathbf{k}', \mathbf{l}, \mathbf{l}') \\ &= \int \frac{d^2 k^\perp}{4Q\omega(\mathbf{k}_Q)} T^{AB}(\mathbf{k}_Q, \mathbf{k}_{-Q}, \mathbf{k}_{-Q}, \mathbf{k}_Q) \end{aligned} \quad (2.58)$$

is reduced to an integration over the transverse momenta $\mathbf{k}^\perp$ weighted with $\frac{Q}{\omega(\mathbf{k}_Q)}$. Here, T^{AB} abridges $T^A T^B$. Analogously, the delta-distribution (appearing in the last four terms)

$$\begin{aligned}\delta(k' - k + q) &= \delta(\omega(\mathbf{k}') - \omega(\mathbf{k})) \delta(k^{3'} - k^3 - Q) \delta^{(2)}(\mathbf{k}^{\perp'} - \mathbf{k}^\perp) \\ &= \frac{\omega(\mathbf{k}_Q)}{Q} \delta(k^3 + Q/2) \delta(k^{3'} - Q/2) \delta^{(2)}(\mathbf{k}^{\perp'} - \mathbf{k}^\perp)\end{aligned}\quad (2.59)$$

leads to

$$\begin{aligned}(2\pi)^6 \int d^R k d^R k' d^R l d^R l' \delta^4(k' - k + q) \delta^{(3)}(\mathbf{k} - \mathbf{l}') \delta^{(3)}(\mathbf{k}' - \mathbf{l}) T^{AB}(\mathbf{k}, \mathbf{k}', \mathbf{l}, \mathbf{l}') \\ = \int \frac{d^2 k^\perp}{4Q\omega(\mathbf{k}_Q)} T^{AB}(\mathbf{k}_{-Q}, \mathbf{k}_Q, \mathbf{k}_Q, \mathbf{k}_{-Q})\end{aligned}\quad (2.60)$$

which means that the rôles of k and k' are interchanged. Here we have used the fact that $\omega(k^3 + Q) = \omega(k^3)$ implies that $k^3 = -\frac{Q}{2}$. Delta distributions of the form (appearing in the middle)

$$\delta^{(4)}(k' + k + q) = \delta(\omega(\mathbf{k}') + \omega(\mathbf{k})) \delta(k^{3'} - k^3 - Q) \delta^{(2)}(\mathbf{k}^{\perp'} - \mathbf{k}^\perp) = 0 \quad (2.61)$$

vanish with a space-like four-vector q since $\omega(\mathbf{k}) = -\omega(\mathbf{k}')$ cannot be fulfilled. The same holds for the terms of the form

$$\delta^{(4)}(k' + k - q). \quad (2.62)$$

We end up with

$$\begin{aligned}
W^{\mu\nu} = & \sum_{s,s',\sigma,\sigma'} \int d^2k^\perp \frac{E}{4Q\omega(\mathbf{k}_Q)} \\
& (T_1^{A\mu\nu}(s, s', \sigma, \sigma'; \mathbf{k}_Q, \mathbf{k}_{-Q}, \mathbf{k}_{-Q}, \mathbf{k}_Q) [f_{s,\sigma'}(\mathbf{k}_Q) - f_{\sigma,s'}(\mathbf{k}_{-Q})] \\
& - T_2^{A\mu\nu}(s, s', \sigma, \sigma'; \mathbf{k}_Q, \mathbf{k}_{-Q}, \mathbf{k}_{-Q}, -\mathbf{k}_Q) \Xi_{\sigma',s}^*(-\mathbf{k}_Q) \\
& + T_3^{A\mu\nu}(s, s', -s', s, \sigma, \sigma'; \mathbf{k}_Q, \mathbf{k}_{-Q}, -\mathbf{k}_{-Q}, \mathbf{k}_Q) \Xi_{\sigma,s'}(-\mathbf{k}_{-Q}) \\
& + T_{14}^{A\mu\nu}(s, s', \sigma, \sigma'; \mathbf{k}_{-Q}, \mathbf{k}_Q, -\mathbf{k}_Q, \mathbf{k}_{-Q}) \Xi_{s',\sigma}^*(\mathbf{k}_Q) \\
& - T_{15}^{A\mu\nu}(s, s', s', -s, \sigma, \sigma'; \mathbf{k}_{-Q}, \mathbf{k}_Q, \mathbf{k}_Q, -\mathbf{k}_{-Q}) \Xi_{s,\sigma'}(\mathbf{k}_{-Q}) \\
& + T_{16}^{A\mu\nu}(s, s', \sigma, \sigma'; \mathbf{k}_{-Q}, \mathbf{k}_Q, \mathbf{k}_Q, \mathbf{k}_{-Q}) [\bar{f}_{\sigma',s}(\mathbf{k}_Q) - \bar{f}_{s',\sigma}(\mathbf{k}_{-Q})]) \quad .
\end{aligned} \tag{2.63}$$

The fluctuation functions Ξ do not appear if we repeat the same calculation in the FF (for a definition and references see Chapter [3](#)). They correspond to a reflection of particles "backward in time". The equivalent of these terms in ELGT has been described in [\[61\]](#). At first sight it would seem that the presence of these functions would spoil the interpretation of structure functions in terms of distribution functions alone. It would also seem to mean that a relation between distribution functions and structure function can only be established in the FF. Fortunately, however, the leptonic tensors associated with scattering backward in time are order $\mathcal{O}(m)$ in the limit $m \rightarrow 0$. The same holds for helicity-flip processes. If fermion masses and transverse momenta can be neglected, only the helicity-nonflip distribution functions $f_{\sigma\sigma}$ contribute to the hadronic tensor $W^{\mu\nu}$. In this limit, the hadronic tensor reads:

$$\begin{aligned}
W^{\mu\nu} \approx & \sum_s \int d^2k^\perp \frac{E}{4Q\omega(\mathbf{k}_Q)} (l_s^{\mu\nu} f_s(\mathbf{k}_Q) - l_s^{\mu\nu} f_s(\mathbf{k}_{-Q}) + \bar{l}_s^{\mu\nu} \bar{f}_s(\mathbf{k}_Q) - \bar{l}_s^{\mu\nu} \bar{f}_s(\mathbf{k}_{-Q})) \\
\approx & \sum_s \int d^2k^\perp \frac{E}{2Q^2} (f_s(\mathbf{k}_Q) l_s^{\mu\nu} + \bar{f}_s(\mathbf{k}_Q) \bar{l}_s^{\mu\nu}) + \mathcal{O}(m/Q)
\end{aligned} \tag{2.64}$$

where we have defined the LEPTONIC TENSOR

$$l_s^{\mu\nu}(\mathbf{k}, \mathbf{k}') \stackrel{def}{=} l_{\oplus}^{\mu\nu}(\mathbf{k}, \mathbf{k}') + (2s)l_{\ominus}^{\mu\nu}(\mathbf{k}, \mathbf{k}') \quad (2.65)$$

$$\bar{l}_s^{\mu\nu}(\mathbf{k}', \mathbf{k}) \stackrel{def}{=} l_{\oplus}^{\mu\nu}(\mathbf{k}', \mathbf{k}) - (2s)l_{\ominus}^{\mu\nu}(\mathbf{k}', \mathbf{k}) \quad (2.66)$$

$$l_{\oplus}^{\mu\nu}(\mathbf{k}, \mathbf{k}') \stackrel{def}{=} 4k^{\mu}k^{\nu} + 2k^{\mu}q^{\nu} + 2k^{\nu}q^{\mu} - Q^2g^{\mu\nu} \quad (2.67)$$

$$l_{\ominus}^{\mu\nu}(\mathbf{k}, \mathbf{k}') \stackrel{def}{=} 2im\epsilon^{\mu\nu\alpha\beta}q_{\alpha}^{\uparrow}s_{\beta} \quad (2.68)$$

and the PARTON DISTRIBUTION FUNCTIONS (as opposed to the *raw* distribution functions f or distribution functions proper)

$$\mathbf{f}_s(\mathbf{k}^{\perp}, +k^3; \mathbf{P}) \stackrel{def}{=} f_s(\mathbf{k}^{\perp}, +k^3; \mathbf{P}) - \bar{f}_s(\mathbf{k}^{\perp}, -k^3; \mathbf{P}) \quad (2.69)$$

$$\bar{\mathbf{f}}_s(\mathbf{k}^{\perp}, +k^3; \mathbf{P}) \stackrel{def}{=} \bar{f}_s(\mathbf{k}^{\perp}, +k^3; \mathbf{P}) - f_s(\mathbf{k}^{\perp}, -k^3; \mathbf{P}) = -\mathbf{f}_s(\mathbf{k}^{\perp}, -k^3; \mathbf{P}) \quad (2.70)$$

The particular form of (2.64) has an intuitive interpretation. In the impulse approximation, partons are considered to be free. A fermion with momentum $\mathbf{k}_Q$ is scattered to a different place $\mathbf{k}_{-Q}$ on the momentum lattice. Should the place $\mathbf{k}_{-Q}$ be already occupied by another parton of the same type, however, then the Pauli exclusion principle prevents it from being deposited there. The process of 'chasing' the occupant of $\mathbf{k}_{-Q}$ away would be a scattering event of higher order.

4 Parton Distribution Functions: Two Definitions

	distribution	unpolarised	polarised
parton distribution	$\mathbf{f}_s(\mathbf{k}; \mathbf{P})$	$\mathbf{f}(\mathbf{k}; \mathbf{P})$	—
raw distribution	$f_s(\mathbf{k}; \mathbf{P})$	$f(\mathbf{k}; \mathbf{P})$	—
integrated parton distribution	$\mathbf{q}_s(x_B, Q)$	$\mathbf{q}(x_B, Q)$	$\mathbf{g}(x_B, Q)$
integrated raw distribution	$\mathbf{q}_s(x_B, Q)$	$\mathbf{q}(x_B, Q)$	$g(x_B, Q)$

Table 2.2: Overview of distribution functions

We may define integrated PARTON DISTRIBUTION FUNCTIONS $\mathbf{q}$

$$\mathbf{q}_s(x_B, Q) \stackrel{def}{=} E_B \int d^2 k^\perp f(\mathbf{k}^\perp, \frac{Q}{2}, \frac{Q}{2x_B}) \approx \frac{Q}{2x_B} \int d^2 k^\perp f_s(\mathbf{k}^\perp, \frac{Q}{2}, \frac{Q}{2x_B}) \quad (2.71)$$

$$\mathbf{q}_s(x_B, Q) \stackrel{def}{=} E_B \int d^2 k^\perp \mathbf{f}(\mathbf{k}^\perp, \frac{Q}{2}, \frac{Q}{2x_B}) \approx \frac{Q}{2x_B} \int d^2 k^\perp \mathbf{f}_s(\mathbf{k}^\perp, \frac{Q}{2}, \frac{Q}{2x_B}) \quad (2.72)$$

which correspond to what is usually referred to as 'distribution functions' in the terminology of DIS. From these we finally form the UNPOLARISED PARTON DISTRIBUTION FUNCTION

$$\mathbf{q}(x_B, Q) \stackrel{def}{=} \mathbf{q}_{+\frac{1}{2}}(x_B, Q) + \mathbf{q}_{-\frac{1}{2}}(x_B, Q) \quad (2.73)$$

and the POLARISED PARTON DISTRIBUTION FUNCTION

$$\mathbf{g}(x_B, Q) \stackrel{def}{=} (2\mathcal{H}) \left[\mathbf{q}_{+\frac{1}{2}}(x_B, Q) - \mathbf{q}_{-\frac{1}{2}}(x_B, Q) \right] \quad (2.74)$$

(analogous definitions for the anti-quark distributions $\bar{\mathbf{q}}$). This definition implies the crossing relations [60] $\bar{\mathbf{q}}(x_B) = -\mathbf{q}(-x_B)$. An overview of the distribution functions introduced so far is given in **Tab. 2.2**. The parton distribution function $\mathbf{q}$ may be written as $\mathbf{q}(x_B, Q) = \mathbf{q}(x_B, Q|P_B S)$ where

$$\mathbf{q}(x_B, Q|PS) \stackrel{def}{=} \int_\ell \frac{dy^\mu}{2\pi} \tilde{\mathbf{q}}_\mu(y|PS) e^{iy \cdot q/2} \quad ; \quad \ell \stackrel{def}{=} \{qs ; s \in \mathbb{R}\} \quad (2.75)$$

$$\tilde{\mathbf{q}}^\mu(y|PS) \stackrel{def}{=} (2\pi)^3 E \langle PS | \bar{\psi}_A(Y) \gamma^\mu \psi_A(Y+y) | PS \rangle \quad (2.76)$$

$$= (2\pi)^3 E \langle PS | \bar{\psi}(Y) \gamma^\mu U_A(Y+y) \psi(Y+y) | PS \rangle \quad . \quad (2.77)$$

is a *space-like* line-integral. Y is an arbitrary four-vector which shall be chosen as $Y = 0$ from now on. This can be verified by inserting free fields into (2.75). The gauge string $U_A(y) \stackrel{def}{=} \exp(i \int_0^y dy'^\mu g A_\mu(y'))$ ensures the gauge-invariance of $\mathbf{q}$. It contains a *space-like* contour-integral over A linking the points 0 and y . In the Breit frame, the integration contour ℓ lies on the quantisation hyper-surface x^0 and U_A contains an integral over the A_3 component of the gauge-field. The integration contour appears to be light-like only in a frame that moves with the speed of light relative to the Breit frame— but this perception is wrong. The momentum transfer q is always space-like, even in the IMF. The properties "space-like" and "time-like" are Lorentz-invariant properties: they cannot be changed by boosts. In the axial gauge $A_3 = 0$, the gauge-string $U_G = \mathbb{1}$ is the unit-operator. For any space-like q , it is a space like AXIAL GAUGE $A \cdot q = 0$ which eliminates U_A —not the light-like gauge $A_- = 0$ sometimes called LIGHT-CONE GAUGE. This gauge coincides with the gauge $q \cdot A = 0$ only if the exchanged photon with momentum q is *real*, i.e. $q \cdot q = -Q^2 = 0$. Only in this case does the light-cone gauge constitute an advantage. The claim in the literature [2, 60] that structure functions have to be expressed in terms of light-like correlation functions is based on a *frame dependent* argument. This argument [60] which allegedly proves that DIS is dominated by light-like correlation functions is based on the assumptions that (a) the four-momentum P of the hadron is fixed and (b) the momentum transfer Q becomes *infinite* (as opposed to large but finite) while x_B is zero. Expression (2.75) allows to trace the implications

of such assumptions. The Lorentz-invariant RELATIVE VELOCITY²

$$v_B = 1/\sqrt{1 + \left(\frac{MQ}{P \cdot q}\right)^2} = 1/\sqrt{1 + \left(\frac{2Mx_B}{Q}\right)^2} \quad (2.78)$$

between P and q becomes 1 in the limit where the experimental resolution Q diverges. In this limit, an unphysical boost with boost-velocity $v = 1$ would be needed in order to relate the Breit frame to a frame with finite P . Hence, the limit $Q \rightarrow \infty$ renders the choice of frame irreversible. In this limit it appears as if q were light-like. Consequently, the integration contour ℓ along the q direction appears to be light-like, too. But q never becomes light-like: q remains space-like even for the somewhat grotesque choice of an infinite experimental resolution $Q = \infty$, q because $-q \cdot q = Q^2 = \infty$ is not zero, as required for a space-like four-vector. The integration contour ℓ becomes light-like for $Q = \infty$ but remains space-like for any finite experimental resolution Q . Instead of choosing a frame with fixed P we might as well choose a frame with fixed q/Q . In this case, it is the hadron which approaches the speed of light in the limit $Q \rightarrow \infty$. Physics only depends on the relative velocity v_B between P and q . The absolute velocity of P or of q is completely irrelevant. An argument which crucially depends on keeping P fixed instead of q , can not be trusted since it is a frame-dependent argument: Indeed, repeating the argument given in Ref. [60] in a frame where the orientation of q is kept fixed, leads to completely different conclusions.

Care must always be taken when a boost with boost-parameter $v = 1$ goes into an argument. These boosts are singular (and should therefore not really be called boosts) as they contract a four-dimensional universe onto a three-dimensional sub-space: the light-front. After the action of such a "boost",

² The reason for calling v_B the relative velocity is that the proton with 4-momentum P moves with the velocity v_B in the rest-frame of q defined as one of the frames where $q^0 = 0$. P and q define two Lorentz-invariants. One possible choice is x_B and Q ; an other choice is v_B and x_B . The last choice is obviously problematic in the Bjorken limit.

all four-vectors appear to be light-like, or "light-cone" dominated. Please note that the Fourier transform $\tilde{W}^{\mu\nu}(y)$ is indeed "light-cone" dominated in the sense that $|\tilde{W}^{\mu\nu}(y)|$ is largest close to the light-cone $y \cdot y = 0$. We do not deny this. What we claim is this: distribution functions are related to space-like line-integrals. There is no need to choose a frame where the integration contours become light-like and there is no need to choose $Q = \infty$ (instead of merely large $Q \gg M$) as there is no point in considering experiments with infinite experimental resolution. For finite resolution Q , however, it is not justified to treat the integration contour as light-like. An infinite boost with $v = 1$ would be needed in order to justify such a step.

We close this section with a few words on terminology. If a function $\tilde{W}(x)$ is almost zero outside the region $R_\epsilon = \{x | 0 \leq x^2 < \epsilon\}$, we would prefer to characterise $\tilde{W}(x - y)$ as being dominated by small, *time-like* distances τ since the relativistic measure of distance is $\tau \stackrel{def}{=} \sqrt{(x - y) \cdot (x - y)}$ — not $r = |\mathbf{x} - \mathbf{y}|$. Applied to DIS this means: the *interval of time* 2τ during which the scattering process takes place is *short*.

Unfortunately, another process is called short-distance dominated in present terminology. We shall argue that "small-(hyper-)volume-dominated process", would be more a more accurate expression — albeit more lengthly. This process, a process with time-like momentum transfer³ $q = (Q, 0, 0, 0)$ probes values of $\tilde{W}(x)$ in a small SPACE-TIME VOLUME, i.e. the major contribution to the Fourier transformed function $W(q)$ stems from a finite space-time region with $x \cdot x \approx 0$ and $|x^0| < 1/Q$. We recall that a hyper-volume is a Lorentz-invariant whereas a difference of two spatial components $|\mathbf{x} - \mathbf{y}|$ is not. The difference of the notions "small hyper-volume" and "small distance" is subtle but crucial.

³ for instance inclusive e^+e^- annihilation, see [60]

We are aware of the fact that these points may contradict some researchers we hold in deep respect and whose publications have taught us much (maybe not enough). If we are wrong or if we have misrepresented ideas, we would like to apologize in advance. The same applies to Chapter **3**.

5 Special Case: Breit Frame

The leptonic tensors in the Breit frame take on the form

$$l_s(k_Q, k_{-Q}) \approx Q^2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & +i(2s) & 0 \\ 0 & -i(2s) & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (2.79)$$

$$\bar{l}_s(k_{-Q}, k_Q) \approx Q^2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -i(2s) & 0 \\ 0 & +i(2s) & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (2.80)$$

if both transverse momenta $\mathbf{k}^\perp$ and fermion masses m are small when compared to the longitudinal momenta k^3 .

$$l^{00} = 4[m^2 + \mathbf{k}_\perp^2] = 4\omega^2(\mathbf{k}_Q) - Q^2 \approx 0 \quad (2.81)$$

$$l^{11} = 4(k^1)^2 + Q^2 \approx Q^2 \quad (2.82)$$

$$l^{22} = 4(k^2)^2 + Q^2 \approx Q^2 \quad (2.83)$$

Inserting the expression **(2.79)** into the hadronic tensor **(2.64)** allows us to compute the structure functions in terms of the parton distribution functions $\mathbf{q}$ and $\mathbf{g}$

$$F_2(x_B, Q) \approx 2x_B F_1(x_B, Q) \approx x_B \mathbf{q}(x_B, Q) + x_B \bar{\mathbf{q}}(x_B, Q) \quad (2.84)$$

$$g_1(x_B, Q) \approx \mathbf{g}(x_B, Q) + \bar{\mathbf{g}}(x_B, Q) \quad (2.85)$$

and

$$\frac{1}{2}g_2(x_B, Q) \approx 0 \quad (2.86)$$

If we had taken several quark flavours i into account we would have obtained

$$F_2(x_B, Q) \approx 2x_B F_1(x_B, Q) \approx x_B \sum_i e_i^2 [\mathbf{q}(x_B, Q; i) + \bar{\mathbf{q}}(x_B, Q; i)] \quad (2.87)$$

$$g_1(x_B, Q) \approx \frac{1}{2} \sum_i e_i^2 [\mathbf{g}(x_B, Q; i) + \bar{\mathbf{g}}(x_B, Q; i)] \quad (2.88)$$

where e_i is the charge of the quark with flavour i , $\mathbf{q}(x_B, Q; i)$ and $\mathbf{g}(x_B, Q; i)$ are its respective parton distribution functions.

6 Distribution Functions in Other Frames

If we use a frame with $\mathbf{P} \parallel \mathbf{q}$ but with $q^0 \neq 0$ then the relations between raw distribution functions and structure functions become slightly more complicated. In the limit where fermion masses and transverse momenta can be neglected, the structure function F_1 reads

$$\begin{aligned} F_1^{\text{IA}}(x_B, Q; P^3) &= \mathbf{q}(x_B, P^3) - \bar{\mathbf{q}}\left(-\frac{Q^2}{(2P^3)^2 x_B}, P^3\right) \\ &\quad + \bar{\mathbf{q}}(x_B, P^3) - \mathbf{q}\left(-\frac{Q^2}{(2P^3)^2 x_B}, P^3\right) \end{aligned} \quad (2.89)$$

which for $P^3 = P_B = \frac{Q}{2x_B}$ coincides with the expression we gave for the Breit frame as it should. We have written $F_1^{\text{IA}}(x_B, Q; P^3)$ instead of $F_1(x_B, Q)$ since this formula is only accurate to the extent that the impulse approximation can be trusted. The full structure function

$$F_1(x_B, Q) = F_1^{\text{IA}}(x_B, Q; P^3) + \triangle F_1^{\text{IA}}(x_B, Q; P^3) \quad (2.90)$$

should depend on the invariants x_B and Q only; it should not depend on the momentum of the hadron. The expression $F_1^{\text{IA}}(x_B, Q; P^3)$ obtained in the impulse approximation, however, does depend on P^3 . Yet it does so weakly. The same holds for higher-order corrections $\Delta F_1^{\text{IA}}(x_B, Q; P^3)$ to the impulse approximation. We recall a feature of non-perturbative physics which is fundamentally different to what the practitioner of perturbation theory is used to: creation operators $b_s^\dagger(\mathbf{k})$ are not irreducible representations of the interaction-dependent Lorentz group. They are irreducible representations of the Euclidean group $E(3)$ ⁴ only. The generators of the Euclidean group are KINEMATICAL, i.e. they do not depend on the interaction. A rotation transforms a creation operator into another creation operator. A boost, however, is DYNAMICAL: it contains interactions and, therefore, a boost transforms a quark creation operator into a complex mixture of quark operators and gluon operators. This is why—in the presence of *non-perturbative* interactions— *creation operators do depend on the quantisation surface*. A distribution function $f(\mathbf{k}; \mathbf{P})$ of virtual particles defined on a given quantisation surface and distribution functions of virtual particles defined on a different quantisation surface are essentially different. Distribution functions are not Lorentz-covariant. Contrary to the intuition gained in perturbation theory, the description of a hadron in terms of *free* quarks and gluons *does* depend on the quantisation surface whereas physical observables must not depend on the quantisation surface.

If the vacuum is not trivial, then only the full structure function F_1 can be independent of the hadron momentum $\mathbf{P}$ as we are going to argue now. An approximation may work better in one frame than it does in another frame especially if it is defined in terms of frame-dependent creation opera-

⁴ I.e. the stability group of the quantisation surface

tors. Indeed, it will turn out that $F_1^{\text{IA}}(x_B, Q; P^3)$ is almost boost invariant for momenta P^3 comparable to P_B whereas the dependence on P^3 becomes substantial for $P^3 \approx 0$ and for $P^3 \rightarrow \infty$. The break-down of the impulse approximation at $P^3 = 0$ is standard knowledge; the break-down of this approximation at infinite momenta, has not yet been realised in the literature.

The equation(2.89) has some salient consequences. In any interacting QFT there is pair-production yielding a non-trivial vacuum. This implies that the raw distribution function $f_s^{\text{vac}}(\mathbf{k})$ of the physical vacuum cannot vanish. Raw distribution functions $f_s(\mathbf{k}, \mathbf{P})$ associated with a physical particle can be decomposed into the parity-invariant vacuum part

$$f_s^{\text{vac}}(\mathbf{k}) \stackrel{\text{def}}{=} \langle \text{vac} | b_s^\dagger(\mathbf{k}) b_s(\mathbf{k}) | \text{vac} \rangle \quad (2.91)$$

and the residual part

$$\triangle f_s(\mathbf{k}, \mathbf{P}) \stackrel{\text{def}}{=} f_s(\mathbf{k}, \mathbf{P}) - f_s^{\text{vac}}(\mathbf{k}) \quad (2.92)$$

associated with the moving particle. Our point is that while there may well be a sufficiently large momentum $|\mathbf{P}|$ such that $\triangle f_s$ vanishes for left-movers, this can no longer be true for the raw distribution f_s itself. The raw distribution function f_s cannot vanish for left-movers— whatever the total momentum $\mathbf{P}$ may be. The presence of a physical particle cannot seriously modify the vacuum state since a particle may be localised inside a finite microscopic region whereas the the vacuum arises through the spontaneous creation and annihilation of virtual particles all over the universe (or all over the lattice). Parity symmetry of QCD implies that the vacuum state must contain the same number of left-movers and right-movers. The state of a fast physical particle which moves right, contains more fast right-movers than left-movers. The wave-function of a fast hadron consists of a long-distance part and a small-distance part. The *long-distance part* consists of virtual particles which

arise from the spontaneous creation of virtual particles all over the universe. The *long-distance part* of the hadronic wave function can be expected to be almost identical to the vacuum wave function. Therefore it contains the same number of left-movers and right-movers. The small distance part, in contrast, consists of virtual particles whose presence is due to the presence of the hadron itself. It is the partons associated with the short-distance part which are accessible in a DIS experiment. Expressed briefly: the hadron is embedded in the vacuum and a non-trivial vacuum always contains left-movers. We shall present an example of this phenomenon in Chapter [3](#).

In the Breit frame, this does not constitute any problem as the *parity invariant* vacuum distribution is subtracted away in Eq. [\(2.64\)](#) and only Δf_s survives. This mechanism need not even be invoked though, as the vacuum distribution can be expected to be concentrated in the region of long wave-lengths $|\mathbf{k}| \lesssim M$ around the origin $\mathbf{k} = 0$ momentum space. The impulse approximation is only valid for large $Q \gg M$ anyway and, therefore, the momentum $|\mathbf{k}_{-Q}| \approx Q/2$ of the scattered parton is not small. The scattered parton is therefore deposited outside the region $|\mathbf{k}| \lesssim M$ where the vacuum distribution is concentrated. In other words: $\mathbf{k}_{-Q}$ cannot probe the vacuum distribution for large Q . In Chapter [4](#) we shall compute the vacuum distribution function of QED(1+1) and demonstrate that it is indeed concentrated inside the region $\mathbf{k} < M$. Consequently, in reference frames that are close to the Breit frame, the vacuum does not influence the structure functions either. The IA becomes less reliable, however, if either the rest-frame or the IMF are approached. *Close to the rest-frame*, the momentum $P^3 x_B$ of the parton before scattering becomes so small that the vacuum is probed. *Close to the IMF*, the momentum $-\frac{Q^2}{4P^3 x_B}$ of the left-moving scattered parton starts to probe the vacuum if P^3 becomes larger than $\frac{Q^2}{4M}$ or

if x_B becomes too small. *Close to the Breit frame*, in contrast, the vacuum is only probed if Q (or x_B) is too small. Therefore, the structure function F_1^{IA} as calculated in the impulse approximation is approximately boost-invariant only for frames close to the Breit-frame. Close to the rest-frame and to the IMF, the structure functions depend severely on the frame (which they should not), a fact which signals the breakdown of the impulse approximation in these frames. Of course, this is no problem for the parton model as such: Observables must not depend on the frame, but an approximation is not required to be boost invariant: the quality of an approximation may depend on the velocity of the physical particle *relative* to the lattice. This simply means that the hadronic tensor has to be computed non-perturbatively in the IMF or the rest-frame where corrections to the parton model become more important.

One comment on the IMF is in order. In the IMF, the vacuum distribution can be eliminated by choosing a very small volume but the IMF is unphysical as discussed in Chapters [3](#) and [4](#).

Finally, we remark that the domain in momentum space

$$\frac{M}{x_B} \ll P^3 \ll \frac{Q^2}{4x_B M} \quad (2.93)$$

where F_1^{IA} is almost independent of P^3 increases in extension if Q is increased or if $1/x_B$ is decreased⁵.

7 Beyond the Impulse Approximation

So far we have been working in the framework of the NAÏVE PARTON MODEL: we did not take higher order *QCD* corrections into account. In this frame-

⁵There is a problem for $x_B \approx 1$ which only becomes apparent when calculating higher-order corrections to the impulse approximation. For more details the reader is referred to [[2](#), [60](#), [62](#)].

work, the domain(2.93) becomes infinitely large in the so-called BJORKEN LIMIT

Definition 1 (Bjorken limit: strong form) *The Bjorken limit is the limit where $Q \rightarrow \infty$ diverges and x_B remains constant.*

As in the last section, we argue that this limit is not useful at all in its *strong* form, $Q = \infty$, as the strong Bjorken limit corresponds to the limit of infinite experimental resolution. In praxis, it suffices to use the Bjorken limit in a weaker form

Definition 2 (Bjorken limit: weak form) *The (weak) Bjorken "limit" is attained if Q is sufficiently large and x_B sufficiently small such that both $M/Q < 1$ and $1/x_B < 1$ are small.*

Cf. [2,62] and references therein. It would be a serious mistake to claim that the impulse approximation becomes better—or even exact—in the Bjorken limit. Even though the effective QCD coupling $\alpha_s(Q)$ becomes smaller⁶ with increasing Q , higher order corrections to the impulse approximation become *more* important (not less) in this limit. In order to justify the parton model, we have but two choices: (1) choosing a resolution Q which is *large enough* compared to M such that $\alpha_s(Q)$ is sufficiently small to validate perturbation theory and yet *small enough* such that higher order QCD corrections are sufficiently suppressed. (2) using the RENORMALISATION GROUP EQUATIONS (RGE) which allows us to describe QCD corrections up to arbitrary orders.

The second choice leads to the RGE improved parton model. Let us discuss the first choice first. The IMPULSE APPROXIMATION—the zeroth order limit of perturbative QCD—can only be accurate *if* first order QCD is accurate as well. First order QCD is accurate if

⁶ One should realise that the effective QCD coupling is always larger than the effective QED coupling for *every* experimental resolution smaller than the great unification scale(!)

1. Q is larger than the typical scale Λ_{QCD} of QCD.
2. the parton masses m_i have approximately the same order of magnitude as Q (in order to avoid logarithmic collinear singularities) $m_i \sim Q$.
3. the momentum of the struck parton is smaller than the cut-off Λ before and after the scattering process
4. the cut-off Λ has the same order of magnitude as Q (in order to perturbatively relate the effective coupling $g_{\text{eff}}(Q)$ to the bare coupling $g(\Lambda)$)

If Q is too large when compared to the parton masses, higher order QCD corrections become too important to be neglected: the naïve impulse approximation breaks down. Fortunately, an arbitrary number of the most important perturbative higher-order corrections (ladder graphs) can be analytically resummed for $M \gg Q$ as explained in [2]. This resummation, which may also be seen as an iteration of first order perturbation theory, is described by the celebrated Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) equations [2, 62]. Improving the impulse approximation (i.e. zeroth order perturbation theory) by taking ladder diagrams into account, the structure function

$$F_1(x_B, Q|P^3) = \hat{F}_1(Q|y_B P^3) \circledast [\mathbf{q}(P^3) + \bar{\mathbf{q}}(P^3)] \\ \stackrel{\text{def}}{=} \int \frac{dy_B}{y_B} \hat{F}_1(x_B/y_B, Q|y_B P^3) [\mathbf{q}(y_B|P^3) + \bar{\mathbf{q}}(y_B|P^3)] \quad (2.94)$$

can be written as a CONVOLUTION $F_1 \circledast \mathbf{q}$ of the quark distribution functions and a function $\hat{F}_1(z_B)$ which may be interpreted as the structure function of a single quark⁷. It can be demonstrated via the OPERATOR PRODUCT EXPANSION [2] that the corrections to (2.94) are repressed by powers of M/Q (this

⁷ Beyond zeroth order perturbation theory, the gluon participates in the scattering process and acquires a structure-function $F_1^{\text{Gluon}}(z_B)$ of its own. Due to confinement, the expressions quark- and gluon- structure functions have to be taken *cum grano salis* [with a grain of salt], of course.

—as an aside— is the rationale for Definition (2)). These so-called HIGHER TWIST CORRECTIONS are due to non-zero parton masses, more general perturbative diagrams, non-perturbative final state/confining interactions etc. To the extent that (2.94) is valid, scattering to arbitrarily large order in ladder diagrams off *point-like quarks* may be replaced by zeroth order scattering off *quarks with internal structure* described by their structure function $\hat{F}_1$. This justifies our using the impulse approximation: *Apart from this modification, our results based on the impulse approximation remain intact.* This modification constitutes the highly successful RENORMALISATION GROUP IMPROVED PARTON MODEL. The NAÏVE PARTON MODEL based on the impulse approximation, in contrast, would correspond to $\hat{F}_1(z_B) = \delta(1 - z_B)$. If the parton distribution functions $q(x_B|P^3)$ are approximately boost-invariant, i.e. if the dependence on P^3 is only weak, then the NAÏVE PARTON MODEL predicts BJORKEN SCALING: The structure function $F_1(x_B, Q)$ becomes independent of Q . This is *not* what one observes experimentally. It is not true that the structure function F_1 loses its dependence on Q in the limit $Q \rightarrow \infty$ —contrary to what is often stated in the literature. The dependence on Q does not disappear; it is logarithmic. This means that F_1 may have to be measured over several orders of magnitude in Q in order to observe an appreciable Q dependence. Yet this Q dependence is extremely strong so as to render $F_2(x_B, Q)$ proportional to the delta function $\delta(x_B)$ [62] in the limit of infinite experimental resolution $Q \rightarrow \infty$. In other words: experiments at infinite resolution—if they were possible— would yield trivial, divergent results⁸. We are always interested in finite experimental resolu-

⁸ Divergences in field-theories are not unphysical as such: QCD predicts finite results for finite experiments. For infinite experiments, e.g. $Q = \infty$, it predicts infinite results—a fact which should not surprise us too much. Moreover, QCD must be seen as an effective theory of general theory which includes gravitation and other forces: the standard model no longer describes reality if Q is large enough to see *new physics*.

tion and therefore, the Bjorken limit should be defined as $Q \gg M$ rather than $Q = \infty$. For the sake of completeness we briefly mention that other problems arise for the IA if the parton masses m are too small when compared to the hadron mass M . In this case, $\hat{F}_1(z_B)$ never equals $\delta(1 - z_B)$ even if Q is very small. The basic picture, however, is salvaged by the fact that $\hat{F}_1$ can be written as a convolution of a m/M dependent function and a Q/M dependent function [2]. This justifies the use of perturbation theory (i.e. the DGLAP equations) in order to describe the Q dependence of the structure functions even in the limit $m/M \rightarrow 0$ where the m/M dependent convolution-factor of $\hat{F}_1$ is not perturbatively accessible. The reader interested in this subject who wants to go beyond the brief (and thus necessarily over-simplified) explanations in this section is referred to [1, 2, 62, 63].

7.1 The Breit Frame and the Continuum Limit

In the BREIT FRAME, the BJORKEN LIMIT corresponds to the CONTINUUM LIMIT as follows. For a sensible computation, the cut-off Λ has to be chosen larger than the total momentum P^3 . At the same time, Λ must not be much larger than Q in order for perturbation theory to be applicable. Therefore $P^3 = \frac{Q}{2x_B} \sim \Lambda$. Consequently, the (strong) Bjorken limit $x_B = \text{const}$, $Q \rightarrow \infty$ coincides with the continuum limit $\Lambda/M \rightarrow \infty$. This has an intuitive interpretation: Q corresponds to the experimental resolution and Λ is the lattice resolution. High experimental resolution can only be achieved if the lattice resolution is high, too.

Chapter 3

Instant-Form and Front-Form: A Campaign For Real Time

"It was during the course of my researches at the Campaign for Real Time that I..."

"The what?" said Arthur again.

[Douglas Adams: Life, the Universe and Everything]

1 The Problem of Front-Form Quantisation

In non-relativistic quantum-mechanics, there is only one possible quantisation surface: the INSTANT OF TIME $t = x^0 = 0$. In relativistic quantum-mechanics, however, there is a larger choice of quantisation surfaces since time is no absolute concept but depends on the observer. How can we tell whether a three-dimensional (hyper-) surface of four-dimensional space-time may serve as quantisation surface? This question must be discussed on the classical level first since the INITIAL SURFACE of the classical equations of motion corresponds to the QUANTISATION SURFACE of the quantised system. The Euler-Lagrange equations of classical, Newtonian mechanics are deterministic: For instance, the non-relativistic, *classical* Euler-Lagrange equations of motion are deterministic in the sense that the knowledge of every

observable at one instant of time suffices in order to predict the entire future or the past. Expressed in the terminology of partial differential equations this reads: the instant of time $t = 0$ is an INITIAL SURFACE of the equations of motion because initial data specified on this hyper-plane determine a unique solution to the equations of motion.

Classical relativistic physics is deterministic as well: the relativistic equations of motion allow the state of the universe to be calculated at any time from the information specified at one INSTANT OF TIME. Contrary to non-relativistic physics, however, there is a larger class of three-dimensional hyper-surfaces of the four-dimensional space-time which may be regarded as an instant of time. The non-uniqueness of the initial surface is a direct consequence of the finiteness of the speed of light. Every space-like hyper-plane may be regarded as an INSTANT OF TIME whose points are not causally connected and consequently, any *space-like* hyper-surface may serve as initialisation-surface (or as quantisation-surface in the quantum case).

Non-relativistically, in contrast, information may propagate arbitrarily fast. Consequently, only space-time events occurring at the same (*absolute*) time are causally independent; rephrased in geometrical terms this reads: an instant of time has only one possible orientation in space-time. In brief, the mathematical term INITIAL SURFACE is intimately related to the more intuitive terms INSTANT OF TIME and SET OF CAUSALLY INDEPENDENT POINTS. This observation will enable us to understand the problems treated in this chapter more easily.

Naturally, the question arises whether quantisation surfaces other than space-like hyper-planes would serve as well in the relativistic case. The first to ask this question was P.A.M. Dirac [47]. In 1950 he proposed to use light-like hyper-planes $x^+ = 0$ as quantisation-surfaces. Note that a light-

like hyper surface $x^+ = 0$ is not defined as an instant of REAL TIME x^0 : it is defined in terms of an abstract time x^+ — called LIGHT-CONE time which does not coincide with the proper time of *any* real observer. For this form of quantisation, Dirac coined the term FRONT FORM (FF) quantisation in order to distinguish it from the more familiar quantisation on a space-like hyper-plane which he referred to as INSTANT FORM (IF). Dirac's idea, soon forgotten, re-emerged again for several times under different names such as LIGHT-CONE, NULL-PLANE or LIGHT-FRONT quantisation. For a more thorough historical overview the reader is referred to [41].




Figure 3.1: The IF co-ordinates x^0, x^3 and their corresponding FF co-ordinates $x^\pm \stackrel{def}{=} x^0 \pm x^3$

Sometimes, the FF is referred to as INFINITE MOMENTUM FRAME (IMF) [64]. The IMF is basically a frame where every particle moves with the speed of light. For more precise a definition see section ⟨8⟩. The confusion of the IMF and the FF arises because the two approaches are very similar (when expressed in light-cone co-ordinates $x^\pm$). This similarity is deceptive, though, as as we shall show in ⟨8⟩. Dirac also found another possible quantisation surface — a space-time parabola — which he called POINT-FORM. Later

on, in 1977, H. Leutwyler and J. Stern [65] gave a group-theoretical characterisation of all possible quantisation-surfaces adding two additional forms to Dirac's catalogue. They pointed out, furthermore, that a light-like hyper-plane has the largest STABILITY GROUP of all hyper-planes, i.e. the largest number of generators of the Poincaré group which leave the respective hyper-

Relativistic form	Quantisation-surface	Algebra of the Stability group
Front form	$x^+ = e^{(+)} \cdot x = 0$	$\mathbf{P}_\perp, \mathbf{J}_{12}, \mathbf{J}_{03}; \mathbf{J}_{\perp-}, P_-$
Instant form	$x^0 = e^{(0)} \cdot x = 0$	$\mathbf{P}, \mathbf{J}_{mn}$
Point form	$x \cdot x = \kappa^2, x^0 > 0$	$\mathbf{J}_{\mu\nu}$
Line form	$(x^0)^2 - (\mathbf{x}^\perp)^2 = \kappa^2, x^0 > 0$	$P_3, \mathbf{J}_{12}, \mathbf{J}_{03}$
Extended front form	$x^+ x^- = \kappa^2, x^0 > 0$	$\mathbf{P}_\perp, \mathbf{J}_{12}, \mathbf{J}_{03}$

Table 3.1: Relativistic forms. The parameter κ is a real number.

plane invariant. The stability group of a light-like plane is spanned by seven generators, one generator more than the IF or the point-form respectively. The complete catalogue of relativistic forms is represented in **Tab. 3.1** A large stability group is usually advantageous because the generators of this group are KINEMATICAL, i.e. they do not depend on the interactions [65]. But although this may be true, a larger stability-group is *not necessarily* advantageous, as we shall show below and in Chapter⟨5⟩. There are useful generators of the Poincaré group which facilitate the diagonalisation of a Hamiltonian and useless generators that do not. For instance, momenta in the IF are useful, boosts in the IF form are not useful: the fact that the boost in the direction of the quantisation-plane naïvely appears to be kinematical in the FF is not as useful as are rotations and translations in the IF since this boost does not commute with the FF Hamiltonian.

In 1971 it was pointed out by Neville and Rohrlich [66] and later by Steinhardt [67] that initial-conditions specified on only one light-like hyper-surface do not suffice in order to unambiguously integrate the equations of motion of a scalar field theory. *Two* light-like initialisation-planes are necessary in order to properly determine the space-time evolution of a relativistic field. Quantisation on only *one* light-like quantisation plane is mathematically ill-

defined because relativistic equations of motion cannot be un-ambiguously solved if the fields are known on one light-like hyper-surface only. This becomes plausible already from the observation that one point on a light-like hyper-plane may influence another point on this hyper-plane via light-like signals. Consequently, a light-like surface is no instant of time in the sense that points on a light-like surface are not causally independent. One may argue, following Dirac, that signals always propagate slower than the speed of light if massive particles are described only. Dirac, however, considered many-body physics in ref. [47] and did *not* have QFTs in mind. In fact, he was trying to construct alternatives to QFTs. In this section we shall be concerned with field-theories where Diracs argument is not applicable since in a relativistic *field* theory, causal propagators always connect light-like directions regardless if massless particle fields are present or not.

This discovery has some uncomfortable consequences. Firstly, we note that it is difficult to define a FF Hamiltonian approach since two quantisation surfaces require two Hamiltonians generating temporal evolution with respect to *two* "time"-directions of space. Secondly, two quantisation-planes have a smaller stability-group as the naïve quantisation on one light-like plane would suggest, since the translation which leaves one of the two planes invariant necessarily moves the other plane, hence becomes dynamical. This additional dynamical translation P_- is the second Hamiltonian. The generators $J_{\perp-}$ become dynamical as well (thus we venture the opinion that the FF on two quantisation surfaces should be identified with the last form of the catalogue **Tab. 3.1** for $\kappa = 0$). In brief, correct quantisation on two light-like quantisation surfaces shrinks the stability group to a group with *four generators* —three less than the FF. TRANSITIVITY¹ of the stability group

¹Transitivity means that the stability group is able to connect any points on the quantisation surface. See [65].

is lost as well.

In 1994, Heinzl and Werner [68] were able to find a loop-hole avoiding the necessity of two light-like hyper-surfaces. They demonstrated in the case of the *free* scalar theory that one light-like hyper-plane suffices as an initial surface *if* periodic or anti-periodic light-like BOUNDARY CONDITIONS(BC) are chosen on the initialisation-plane. This seemed to provide a justification for FF quantisation on one hyper-plane as long as (anti-) periodic boundary-conditions are specified (as e.g. in DISCRETISED LIGHT-CONE QUANTISATION [41]). However, periodic *light-like* boundary conditions are unphysical as we shall demonstrate in this chapter.

Firstly, periodic BCs eliminate boosts as members of the FF stability-group since boosts change these boundary-conditions. We shall show that boosts are therefore no longer automatically kinematical transformations. In Chapter⟨5⟩ we demonstrate the mass-spectrum of QED((1 + 1) is boost-invariant to a much higher degree in the IF when compared to the FF. Again, as in the case of two initialisation planes, the number of the really kinematical operators becomes the same as in the IF.

Secondly, light-like periodic BCs destroy important symmetries such as parity and rotational invariance *even* in the infinite (light-like) volume limit. While space-like BCs are invariant under parity and time-reversal, light-like BCs are not.

Thirdly, and most importantly, we shall show that light-like boundary conditions completely destroy microcausality even for arbitrarily large light-like volumes. Consequently they are *unphysical*, contrary to space-like boundary conditions. An implication of this is that the speed of light becomes infinite.

It is important to note that the term "unphysical" does *not* mean that *no*

physical observable *at all* can be correctly described. Some observables such as mass spectra are sometimes less sensitive to the damage than propagators, especially for computations performed close to the non-relativistic limit where all relativistic forms must coincide. The damage due to light-like boundary conditions also depends on the theory under consideration. Some specific examples will be given.

2 Planes, Vectors and Frames

In the ensuing sections, we are concerned with general co-ordinate frames and their interpretation. We shall introduce three frames which have to be distinguished for a non-perturbative, theoretical description of a relativistic particle with 4-momentum P :

(F-1) An arbitrary FRAME OF REFERENCE related to the relativistic particle which may be taken to be the rest frame of the particle under study.

(F-2) The INERTIAL FRAME or QUANTISATION FRAME, i.e. the frame where the observer is at rest. This instant of time with respect to this frame defines the quantisation surface upon which initial data and charges are defined.

(F-3) The LATTICE REST FRAME.

Our intention is to study how different quantisation surfaces are related to each other. This has to be done with respect to the fixed but otherwise arbitrary FRAME OF REFERENCE (F-1). In particular, we are interested in the extreme case of co-ordinate transformations which transform a space-like lattice into a light-like lattice. Unfortunately, there is no non-singular Lorentz transformation which does this. Therefore, the lattice rest frame has to be transported by a GENERAL CO-ORDINATE TRANSFORMATION which does not leave the metric invariant. One of these co-ordinate transformations,

the ε co-ordinates [69–71], will be studied in section ⟨7⟩. For the ensuing discussion, it is helpful to briefly review some salient aspects of GENERAL CO-ORDINATE TRANSFORMATIONS. The reader not familiar with the concepts presented in this section is referred to standard text-books on mathematical methods or general relativity as e.g. Ref. [72]. A 4-vector

$$u = u^\mu e_{(\mu)} \quad (3.1)$$

is independent of the CO-ORDINATE FRAME $e_{(\mu)}$ whereas its CONTRA-VARIANT components u^μ or its CO-VARIANT components u_μ do depend on the frame. For any frame $e_{(\mu)}$ of co-variant vectors there is a DUAL FRAME of contra-variant vectors $e^{(\mu)}$ such that

$$e_{(\mu)} \cdot e^{(\nu)} = \delta_\mu^\nu \quad (3.2)$$

holds. An orthonormal co-ordinate system $e_{(\mu)}$ is a co-ordinate system with diagonal Minkowski metric $g_{\mu\nu} \stackrel{def}{=} e_{(\mu)} \cdot e_{(\nu)}$ where $g = \eta$, $\eta^{00} = 1, \eta^{11} = \eta^{22} = \eta^{33} = -1$. GENERAL CO-ORDINATES $\bar{x}^\mu = T^\mu_\nu x^\nu$ are obtained from the ortho-normal co-ordinates x^μ by means of the CO-ORDINATE TRANSFORMATION T^μ_ν . In non-orthogonal co-ordinates, the contra-variant 4-vector $\bar{e}^{(0)} = \bar{g}^{\mu\nu} \bar{e}_{(\nu)}$ need not coincide with the direction $\bar{e}_{(0)}$ of the time axis. What is the intuitive meaning of $\bar{e}_{(\mu)}$ and $\bar{e}^{(\mu)}$? Let us assume that x^μ is the REST FRAME OF THE OBSERVER. In an ORTHO-NORMAL FRAME, the rest frame of the observer may be characterised as one of the frames in which the observer's world-line $x(\tau)$ with $\tau \in \mathbb{R}$ assumes the simple form $x(\tau) = \tau e_{(0)} = \tau e^{(0)}$. In this frame $x^0 = x_0$ coincides with the observer's PROPER TIME $\tau = \sqrt{x \cdot x}$ and the equation $x^0 = x_0 = 0$ defines an instant of time. In general co-ordinates, several characterisations are possible: We define the general REST FRAME OF THE OBSERVER as the frame wherein the observer's world line reads $x(\tau) = \tau e^{(0)}$. In this frame, $\bar{x}_0$ coincides with the observer's PROPER

TIME and $\bar{x}^0 = 0$ defines an INSTANT OF TIME. We could have defined this frame equally well via $x(\tau) = \tau \bar{e}_{(0)}$. In that case, the respective rôles of $\bar{x}_0$ and $\bar{x}^0$ would be interchanged and $\bar{x}^0 = \bar{y}^0$ would no longer mean that the events x and y are simultaneous. The choice of $\bar{x}_0$ as proper time implies that the direction $\bar{e}_{(0)}$ of the time-axis is of minor importance whereas the dual time direction $e^{(0)}$ characterises the observer's world-line and represents, furthermore, the normal vector of the instant of time $x^0 = 0$ where a simultaneous measurement of initial conditions takes place². The co-variant vector $e_{(0)}$, however, is merely the direction of an abstract time-axis, a quantity which is physically irrelevant in a general frame. We shall see that $\bar{e}_{(0)}$ may even be space-like whereas $\bar{e}^{(0)}$ *must* be time-like in order to define a space-like hyper-surface. Henceforth, we shall use $\bar{e}^{(0)}$ synonymously with the instant of time $\bar{x}^0 = 0$ it defines. We summarise: time imbodyes two aspects: the METRIC ASPECT of the observer's proper time $x_0 = \tau$ and the ASPECT OF SIMULTANEITY $\bar{x}^0 = \bar{y}^0$. Only in orthogonal frames can these two aspects be described by the same time variable $x^0 = x_0$.

Another intricacy of general co-ordinate systems —which is easily glimpsed over by the practitioner used to the ortho-normal Minkowski metric— is the definition of a space-time volume element: while the volume element $d^4x = dx^0 dx^1 dx^2 dx^3$ is invariant under orthogonal co-ordinate transformations only (i.e. under Lorentz-transformations), its generalisation

$$d^4x / \sqrt{\det g} = d^4\bar{x} / \sqrt{\det \bar{g}} \quad (3.3)$$

is invariant under general, non-singular co-ordinate transformations. The metric of the light-cone co-ordinates $\bar{x}^0 = x^+$, $\bar{x}^3 = x^3$, for instance, is

$$\bar{g} = \begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix} \quad (3.4)$$

² In a quantised system, this will be the quantisation surface where the state-of-the-universe is prepared.

(transverse co-ordinates suppressed) and therefore

$$k \cdot k = k_+ g^{+-} k_- + k_- g^{-+} k_+ = 4k_+ k_- \quad . \quad (3.5)$$

3 The Quantisation Hyper-Surface

If a charge Q corresponding to a charge density $J^0(x)$ is measured at the instant of time $x^0 = 0$, it may be represented as

$$Q(e^{\text{quant}}) = Q(x^0 = 0) = \int d^3x J^0(0, \mathbf{x}) = \int d^4x J^0(x) \delta(x^0) \quad (3.6)$$

in an *orthonormal* co-ordinate system $e_{(\mu)}$. Note that this charge depends on the hyper-surface $e^{\text{quant}} = e^{(0)}$ upon which it is defined, a detail that will turn out to be crucial. The definition of Q is not linked to a specific co-ordinate system. We may equally well define the same charge

$$Q(e^{\text{quant}}) \stackrel{\text{def}}{=} \int d^4\bar{x} \frac{1}{\sqrt{|\det \bar{g}|}} J^0(x) \delta(e^{\text{quant}} \cdot x) \quad (3.7)$$

in any other general co-ordinate frame $\bar{e}_{(\mu)}$ even if it is not a QUANTISATION FRAME, i.e. even if $e^{\text{quant}} \neq \bar{e}^{(0)}$: what we have gained is that we have disentangled the hyper plane e^{quant} on which the charge is defined from a particular co-ordinate system.

The relativistic 4-momentum P^μ may be considered as a charge. For definiteness, let us now consider a scalar field-theory with Lagrangian

$$\mathcal{L} = \frac{1}{2} \partial^\mu \varphi(x) \partial_\mu \varphi(x) - \frac{m_0^2}{2} \varphi^2(x) - V(\varphi(x)) \quad (3.8)$$

where $V(\varphi)$ is a polynomial of the fields, e.g. $V(\varphi) = \frac{g_0}{4!} \varphi^4$. Invariance of $\mathcal{L}$ under spatio-temporal translations leads to the conservation of the ENERGY-MOMENTUM TENSOR

$$\Theta^{\mu\nu} = \partial^\mu \varphi \partial^\nu \varphi - g^{\mu\nu} \mathcal{L} \quad . \quad (3.9)$$

The charges associated with this current — the 4-momentum that is —

$$P_\mu^{(0)} \stackrel{\text{def}}{=} P_\mu(e^{(0)}) = \int dx^1 dx^2 dx^3 \Theta_\mu^0 \quad (3.10)$$

are conserved *if* the boundary conditions are properly chosen on the quantisation surface. Fixed BCs, for instance, must be excluded here since they would break translational invariance. The spatial momenta $\hat{P}_i(e^{(0)})$ which leave the quantisation surface $e^{(0)}$ invariant are KINEMATICAL, i.e. they do not depend on the interaction. The formal reason for this is that the components $\Theta_i^0 = \partial^0 \varphi \partial_i \varphi$ of the energy-momentum tensor **(3.9)** do not depend on the Lagrangian (because $g_i^0 = 0$).

The conserved charge $P^0(e^{\text{quant}})$ associated with temporal translations, however, is dynamical. It is identical to the Hamiltonian $H(e^{\text{quant}})$ obtained by canonical quantisation if the velocities $\partial_0 \varphi(x^0 = 0, \mathbf{x})$ appearing in Θ_0^0 are expressed in terms of their conjugate momenta

$$\pi(0, \mathbf{x}) = \frac{\partial \mathcal{L}}{\partial \partial_0 \varphi(0, \mathbf{x})} = \partial^0 \varphi(0, \mathbf{x}) \quad (3.11)$$

subjected to the usual commutation-relations

$$[\pi(0, \mathbf{x}), \varphi(0, \mathbf{y})] = -i\delta(\mathbf{x} - \mathbf{y}). \quad (3.12)$$

In the same way, the charges $P_i(e^{\text{quant}})$ with $i = 1, 2, 3$ can be identified with the generators translations in $e_{(i)}$ -direction. In summary, the charges $\hat{P}_i(e^{\text{quant}})$ are generators of space-time translations of the fields $\hat{\varphi}(x)$ in $e^{(\mu)}$ -direction: they fulfill the equations

$$[\hat{P}_\mu, \hat{\varphi}(x)] = i\partial_\mu \hat{\varphi}(x) \quad (3.13)$$

as is easily verified (cf. **[58]**).

On a light-like hyper-surface, CONSTRAINTS are present, i.e. field velocities $\partial_0 \varphi$ cannot be expressed in terms of field momenta $\hat{\pi}$. We shall treat this

case later on in this chapter. Later on we would like to study what happens if the quantisation surface e^{quant} changes relative to some fixed reference frame $e_{(\mu)}$. To this end we have to render Eq.(3.10)

$$\hat{P}_\mu(e^{\text{quant}}) = \int \frac{d^4x}{\sqrt{|\det g|}} \delta(x \cdot e^{\text{quant}}) e_\nu^{\text{quant}} \Theta_\mu^\nu \quad (3.14)$$

independent of the specific reference-frame used. Any non-singular co-ordinate frame with

$$e^{(0)} \parallel e^{\text{quant}} \quad (3.15)$$

may be called a QUANTISATION FRAME associated with the quantisation surface e^{quant} . Note that $\hat{P}_0(e^{(0)})$ is the generator of translations in the direction of the TIME AXIS $e^{(0)}$; if $e^{(0)}$ is not parallel to $e_{(0)}$, then the FLOW OF ARTIFICIAL TIME does not coincide with the observer's world line: $\hat{P}_0(e^{(0)})$ must then be interpreted as the sum of a translation inside the instant of time $\bar{x}^0 = 0$ and $\hat{P}^0(e^{(0)})$, the generator which moves the observer's proper time $\bar{x}_0$. In the FF, proper time $\bar{x}_0 = x_+$ coincides with the *spatial* co-ordinate $\bar{x}^3 = 2\bar{x}_0$. Setting $e^{\text{quant}} = e^{(0)}$ we obtain the IF Hamiltonian

$$P_0(e^{(0)}) = \int d^3x \left[\frac{1}{2}(\hat{\pi})^2 + \frac{1}{2}(\nabla\varphi)^2 + \frac{1}{2}m_0^2\varphi^2 + V(\varphi) \right]_{x^0=0} \quad (3.16)$$

with the commutation relations

$$[\hat{\pi}(0, \mathbf{x}), \varphi(0, \mathbf{y})] = -i\delta(\mathbf{x} - \mathbf{y}) \quad (3.17)$$

and

$$[\varphi(0, \mathbf{x}), \varphi(0, \mathbf{y})] = 0 \quad (3.18)$$

where $\nabla \stackrel{\text{def}}{=} -(\partial_1, \partial_2, \partial_3)$. Setting $e^{\text{quant}} = e^{(+)} = e^{(0)} + e^{(3)}$ we obtain the FF Hamiltonian

$$P_+(e^{(+)}) = \frac{1}{2} \int dx^- d^2x^\perp \left[\frac{1}{2}(\nabla_\perp\varphi)^2 + \frac{1}{2}m_0^2\varphi^2 + V(\varphi) \right]_{x^+=0} \quad (3.19)$$

where $\nabla_{\perp} \stackrel{def}{=} -(\partial_1, \partial_2)$. Using the Dirac-Bergmann constraint-quantisation procedure one can deduce [68, 73] the commutation relations

$$[\varphi(x^+ = 0, x^-, \mathbf{x}^{\perp}), \varphi(y^+ = 0, y^-, \mathbf{y}^{\perp})] = \frac{1}{2i\partial_-} \delta(x^- - y^-) \delta(\mathbf{x}^{\perp} - \mathbf{y}^{\perp}) \quad (3.20)$$

which quantise the theory *if* the field is periodic in x^- direction. Zero modes $\int dx^- \varphi(x^+, x^-)$ of $\varphi(x)$ are subjected to the constraint

$$(-m_0^2 + \nabla_{\perp}^2) \int dx^- \varphi = \int dx^- \frac{\partial}{\partial \varphi} V(\varphi) \quad (3.21)$$

which is an artifact of periodic BCs.

4 The Boundary Vector

Time travel is increasingly regarded as a menace. History is being polluted. The *Encyclopedia Galactica* has much to say on the theory and practice of time travel, most of which is incomprehensible to anyone who hasn't spent at least four lifetimes studying advanced hyper-mathematics, and since it was impossible to do this before time travel was invented, there is a certain amount of confusion as to how the idea was arrived at in the first place. One rationalization of this problem states that time travel was, by its very nature, discovered simultaneously at all periods of history, but this is clearly bunk. The trouble is that a lot of history is now quite clearly bunk as well.

[Douglas Adams: Life, the Universe and Everything]

We shall be concerned with systems on a finite lattice and, therefore, it is convenient to introduce another hyper-plane with normal vector e^{BC} on which the boundary conditions of the box are specified. Periodic boundary conditions may be written in the form

$$\varphi(x + \mathfrak{L}) = \varphi(x - \mathfrak{L}) \quad (3.22)$$

where $\mathfrak{L}$ is a 4-vector with components

$$\mathfrak{L} = (\mathfrak{L}^0, \mathfrak{L}^1, \mathfrak{L}^2, \mathfrak{L}^3) \quad (3.23)$$

In order to obtain a boundary box with finite volume, three boundary vectors $\mathfrak{L}_{(i)}$ ($i \in [1, 2, 3]$)

$$\mathfrak{L}_{(i)} \cdot e^{\text{BC}} = 0 \quad (3.24)$$

with invariant lengths $L_{(i)} \stackrel{\text{def}}{=} \sqrt{-\mathfrak{L}_{(i)} \cdot \mathfrak{L}_{(i)}}$ are necessary. We shall call $2L_{(i)}$ the (invariant) SIZE OF THE LATTICE. As we are primarily interested in boundary conditions in 3-direction we shall skip, henceforth, the index (3) which indicates the 3-direction

$$\mathfrak{L} \stackrel{\text{def}}{=} \mathfrak{L}_{(3)} \quad L \stackrel{\text{def}}{=} L_{(3)} . \quad (3.25)$$

Similarly, a cut-off Λ in momentum space corresponding to a lattice spacing $a = \frac{\pi}{\Lambda}$ may be characterised by the four vector $\mathbf{a} \stackrel{\text{def}}{=} a\mathfrak{L}/L$.

Three-dimensional hyper-plane	normal vector
Instant of time with respect to the co-ordinate frame $e_{(\mu)}$	$e^{(0)}$
Instant of time where charges are measured	e^{quant}
Boundary plane containing the boundary 4-vectors $\mathfrak{L}_{(i)}$	e^{BC}

Table 3.2: Three-dimensional hyper-planes and their corresponding normal 4-vectors

The three hyper-planes e^{quant} , $e^{(0)}$ and e^{BC} —as summarised in **Tab. 3.2**— may in general be different without leading to inconsistencies. For practical purposes, however, not identifying these planes would not be a good idea. It is clearly so exceedingly convenient to use a frame $e^{\text{quant}} = e^{(0)}$ whose instant of time $x \cdot e^{(0)} = 0$ coincides with the quantisation-surface $x \cdot e^{\text{quant}} = 0$ that in QFT text-books, this choice is always implicitly made. Even more expedient, from the practical point of view, is the identification

$$e^{\text{quant}} \equiv e^{\text{BC}} \quad (3.26)$$

of the quantisation hyper-plane and the hyper-plane on which boundary-conditions are chosen unless one is willing to unnecessarily deal with dynamical boundary terms and charges which are not conserved in time. This fact may be formulated as:

Theorem 1 (Charge non-conservation) *Let $Q(e^{(0)})$ be the charge associated with the conserved current $J^\mu(x)$, i.e. $\partial_\mu J^\mu(x) = 0$. All fields are periodic with the periodicity $\mathfrak{L} = (0, 0, 0, \mathfrak{L}^3)$. Then only $Q(e^{\text{BC}})$ is conserved whereas $Q(e^{\text{quant}})$ with $e^{\text{BC}} \neq e^{(0)}$ is not conserved in general.*

Proof: It suffices to give *one* example of a non-conserved charge. Take

$$J(x) = \begin{pmatrix} \rho(x) \\ v\rho(x) \end{pmatrix} \quad (3.27)$$




Figure 3.2: Illustration of charge non-conservation. The boundary conditions join A and B




Figure 3.3: Illustration of micro-causality violation. The boundary conditions join A and B

with

$$\rho(x^0, x^3) = \begin{cases} \delta(x^3 - vx^0 + \mathfrak{L}^3) & \text{if } 0 \leq x^0 < +\mathfrak{L}^3/v \\ \delta(x^3 - vx^0 - \mathfrak{L}^3) & \text{if } -\mathfrak{L}^3/v < x^0 < 0 \end{cases} \quad (3.28)$$

and $0 < v < 1$. The reader will easily convince himself (cf. **Fig. 3.2** and **Fig. 3.3** were $\mathfrak{L} = \overrightarrow{AO} = \overrightarrow{OB}$) that the charge jumps between $Q(\bar{e}^{(0)}) = 0$ and $Q(\bar{e}^{(0)}) = 1$ in the frame

$$\bar{x}^0 = \frac{x^0 - vx^3}{\sqrt{1 - v^2}} \quad \bar{x}^3 = \frac{x^3 - vx^0}{\sqrt{1 - v^2}} \quad (3.29)$$

where $\bar{J}(\bar{x})$ at rest. The deeper mathematical reason behind this is the fact that the conservation of charges follows from Gauss' theorem under the condition that the integration contour over the boundaries vanishes. The charge (3.7) is not properly defined if the boundary terms do not vanish. In order for a charge to be conserved one would have to include the boundary terms into its definition. In particular, the momenta $\hat{P}^\mu(e^{\text{quant}})$ constructed in eq. (3.14) would cease to be conserved charges if (3.26) did not hold and, consequently, they would no longer be identical to the generators of

translations. Periodicity breaks the Poincaré invariance of the Lagrangian density because it singles out the rest frame of $\mathfrak{L}$. This does not merely imply non-conservation of charge, it implies, furthermore, that microcausality is violated in every frame except for the lattice rest-frame with $e^{(0)} = e^{\text{quant}}$. This is demonstrated in **Fig. 3.3** which represents a charge that disappears at the boundary (A) and "re"appears in the past (B). The inhabitants of the $2L$ -periodic universe, however, would need (at least) the time L in order to find out that they are in a frame where charge is not conserved³. This is the same time it would take in order to find out that the universe is $2L$ -periodic. The finiteness of the speed of light justifies the replacement of a potentially infinite universe by a small model universe with *space-like* periodic BCs. Effects of periodic BCs are *unobservable* if the periodicity L is large enough. An experimentalist has access to a finite space-time region of the universe only.

The inhabitants of a universe with light-like BCs (i.e. $L = 0$), in contrast, are always in the "wrong" frame, i.e. in a frame where charge is not conserved since no observer can move with (almost) the speed of light. Worse still, they would find out immediately by sending a light-signal in three-direction: this signal would return almost instantly in the past. Another signal travelling with less than the speed of light could be used in order to transport information instantly to any place in the universe. This is the first of a series of *observable* unphysical predictions resulting from light-like BCs.

We conclude that there is no preferred time direction $e_{(0)}$ when using periodic boundary conditions⁴. The contra-variant 4-vector $e^{(0)}$, in contrast, should be parallel to e^{BC} in order for the naïve charges to be conserved.

³To be more precise, an observer moving with velocity v relative to the lattice would need the time $\mathfrak{L}^3 = L/\sqrt{1-v^2}$ in order to gather the information collected by detectors which measure the total charge at a given instant of time $x^0 - vx^3 = \text{const.}$

⁴except for the orthogonality condition $e^{(0)} = e_{(0)}$ we have excluded

Type of BC	Periodic BCs	Fixed BCs
Conserved quantity	$\forall_{0 \leq \mu \leq 3} P_\mu(e^{\text{BC}})$	$\forall_{e: e^2 < 0} P_0(e)$

Table 3.3: Conserved quantities in the presence of boundary conditions

Thence we shall always assume that $e^{\text{quant}} \equiv e^{\text{BC}}$ unless the contrary is explicitly stated.

Thes conclusions, however, are a particularity of periodic BCs. For fixed boundary conditions, in contrast, the importance of the co-variant time-direction and the contra-variant orientation of the quantisation surface is somewhat reversed. Only the generator $P_{(0)}$ of translations in $e_{(0)}$ -direction is then a conserved quantity whereas spatial momenta cannot be conserved because fixed BCs break translation symmetry. In this case, the contra-variant co-ordinate axis or the quantisation surface have no preferred orientation. This is illustrated in **Tab. 3.3**.

5 Kinematical Equivalence of Relativistic Frames

In this subsection, we introduce the expression KINEMATICAL EQUIVALENCE of co-ordinate systems. In order to avoid lengthy expressions, we shall suppress the distracting transverse co-ordinates x^1 and x^2 whenever they are not important.

Definition 3 *We call the co-ordinate systems x^μ and $\bar{x}^\mu$ KINEMATICALLY EQUIVALENT (abridged $x^\mu \triangleq \bar{x}^\mu$) if their respective instants of time $x^0 = 0$ and $\bar{x}^0 = 0$ coincide.*

This implies, in particular, that two events x_A and x_B which are simultaneous in one frame $x_A^0 = x_B^0$ are simultaneous also with respect to the kinematically

equivalent frame $\bar{x}_A^0 = \bar{x}_B^0$. Let us write the co-ordinate transformation $T : x^\mu \rightarrow \bar{x}^\mu = T^\mu_\nu x^\nu$ in a way

$$T = \begin{pmatrix} F & -vF \\ v'F' & -F' \end{pmatrix} \quad (3.30)$$

similar to a co-ordinate boost (transverse co-ordinates are suppressed, $FF' \neq 0$). Setting $v = v'$, $F = F' = \frac{1}{\sqrt{1-v^2}}$ renders T a combined boost and parity transformation. Setting $F = F' = 1$, $v = -1$ and $v' = 1$ yields the FF co-ordinates $\bar{x}^0 = x^+ = x^0 + x^3$, $\bar{x}^3 = x^- = x^0 - x^3$. The rationale for writing T in this form is that we are interested in the relation between the FF and the IF. The CONTRA-GREDIENT of T

$$T^\dagger \stackrel{def}{=} (T^{-1})^\top = \frac{1}{(1-vv')FF'} \begin{pmatrix} F' & F'v' \\ -Fv & -F \end{pmatrix} \quad (3.31)$$

is the co-ordinate transformation $T^\dagger : x_\mu \rightarrow \bar{x}_\mu = T^\dagger{}^\nu_\mu x_\nu$ which transforms co-variant co-ordinates. Kinematical equivalence between x^μ and $\bar{x}^\mu$ implies that the canonical quantisation procedure ⁵ yields equivalent results in the sense of the following theorem

Theorem 2 *The momenta $P_\mu(e^{(0)})$ obtained through quantisation with respect to the frame x^μ and the momenta $\bar{P}_\mu(\bar{e}^{(0)})$ obtained through quantisation with respect to the kinematically equivalent frame $\bar{x}^\mu$*

$$\bar{P}_\mu(\bar{e}^{(0)}) = T^\dagger{}^\nu_\mu P_\nu(e^{(0)}) \quad (3.32)$$

are related in an interaction-independent way by a mere co-ordinate transformation

Expressed differently, let $P(\bar{e}^{(0)}, \bar{e}_{(0)})$ and $P(e^{(0)}, e_{(0)})$ be the 4-momentum obtained in the kinematically equivalent frames $\bar{e}_{(\mu)}$ and $e_{(\mu)}$ respectively.

⁵Or, if constraints are present, the Dirac-Bergmann algorithm

Then both momenta define the same object, i.e.

$$P(\bar{e}^{(0)}, \bar{e}_{(0)}) = P(e^{(0)}, e_{(0)}) = P(e^{(0)}) \quad (3.33)$$

and one can skip the label $e_{(0)}$. The only difference between the frames $\bar{e}_{(\mu)}$ and $e_{(\mu)}$ is that they project out different components $P_\mu = e_{(\mu)} \cdot P$ from the same object.

The fact that the transformation between $P(e^{(0)})$ and $\bar{P}(\bar{e}^{(0)}) = \bar{P}(e^{(0)})$ is kinematical follows already from the fact that transformations mapping the initialisation hyper-surface onto itself do not depend on interactions. This is, in fact, the very reason for why in [65] kinematical transformations are *defined* as transformations leaving the quantisation surface invariant. It is instructive, though, to show eq. (3.32) by construction. The most general non-singular linear transformation leaving the equation $x^0 = 0$ invariant

$$T = \begin{pmatrix} F & 0 \\ v'F' & -F' \end{pmatrix} \quad (3.34)$$

is obtained by setting $v = 0$. This transformation can be decomposed into a product of T with $F = 1$ and a mere rescaling of the temporal co-ordinate $x^0 \rightarrow Fx^0$. Since two conserved Hamiltonians quantised with respect to the time co-ordinates x^0 and Fx^0 respectively are equivalent —albeit rescaled—, it suffices to study $T(\varepsilon)$ with $F \equiv 1$. The canonical momenta

$$\pi(0, \mathbf{x}) = \partial^0 \varphi(0, \mathbf{x}) = \bar{\partial}^0 \varphi(0, \mathbf{x}) = \bar{\pi}(0, \mathbf{x}) \quad (3.35)$$

appearing in Θ then remain the same under the $T(\varepsilon)$ transformation —by construction. This fact is not restricted to the scalar model. Canonical momenta $\pi(x)$ are always contra-variant zero-components $\pi(x) = \pi^0(x)$ of the 4-vector $\pi^\mu(x) = \partial \mathcal{L} / \partial(\partial_0 \varphi)$ even if φ is not a scalar field. The quantisation surface $e^{(0)}$ remains the same and therefore it follows from (3.14)

$$\bar{P}_\mu(\bar{e}^{(0)}) = \bar{P}_\mu(e^{\text{quant}}) = \bar{P}_\mu(e^{(0)}) = T^{+ \nu}_\mu P_\nu(e^{(0)}) \quad (3.36)$$

that we may construct $\bar{P}_\mu(\bar{e}^{(0)})$ as a linear combination of operators $P_\mu(e^{(0)})$ obtained through a quantisation in kinematically equivalent co-ordinates. The 4-vectors $P(e^{(0)})$ and $P(\bar{e}^{(0)})$ are the same objects; only their co-ordinates $\bar{P}_\mu$ or P_μ differ. Observables such as mass spectra do not depend on which co-ordinates one quantises in —if these co-ordinates are kinematically equivalent. If, however, $e^{(0)} \neq \bar{e}^{(0)}$, then quantisation with respect to inequivalent co-ordinate systems yields different 4-momenta. Worse still. If $P_\mu(e^{(0)})$ is a conserved quantity, $P_\mu(\bar{e}^{(0)})$ cannot be conserved due to Theorem (1). We conclude that the arbitrary⁶ general time direction $\bar{e}_{(0)}$ and the generator $\bar{P}_{(0)}$ of translations in this direction are both bereft of physical information. In general (i.e. non-orthogonal) co-ordinates, $P^{(0)}(\bar{e}^{(0)})$ is the PHYSICAL HAMILTONIAN which advances the proper-time $\bar{x}_0$ of the observer. This Hamiltonian is identical in all kinematically equivalent frames —except for rescaling.

6 Defining Boost Invariance

Now we are able to dispel the claim that, contrary to the IF, the mass spectrum of the FF is automatically boost-invariant (under boosts in the three-direction to be more precise). This claim is usually based on the observation that boosts in three directions do not change the FF quantisation surface, i.e. they do not change a light-like hyper-surface $e^{(+)}$ and consequently, they are kinematical. But we have to ask first what boost-invariance means. We give the following *necessary* criterion

Definition 4 (Boost invariance) *For a theory characterised by the 4-momentum operators $\hat{P}(e^{quant})$ to be boost-invariant in three direction, the expectation*

⁶except that the time axis must not lie in the quantisation surface.

value

$$M^2(P) \stackrel{def}{=} \left\langle P, \iota | \hat{M}^2 | P, \iota \right\rangle \quad (3.37)$$

of the mass-squared operator $\hat{M}^2 := \hat{P} \cdot \hat{P}$ must not depend on the kinematical component $P_3(e^{quant})$ (or on $P_-(e^{quant})$). Here, $|P, \iota\rangle$ denotes an eigenstate of $\hat{P}$, ι stands for conserved quantum numbers other than the momentum.

In the IF, this simply means that computed physical masses must not depend on the three-momentum P_3 for $|\mathbf{P}| \ll \Lambda$. In the FF, this means that physical masses must not depend on the minus-momentum P_- . In praxis, however, QFTs need to be regularised, e.g. by introducing a lattice. A lattice, characterised by its boundary vector $\mathfrak{L}$ or by its cut-off 4-vector $\mathbf{a}$, breaks the Poincaré invariance of the Lagrangian density and therefore, the computed physical masses can at best be *approximately* independent of the momenta. For renormalisable theories, the degree to which $M(\mathbf{P})$ is independent of $\mathbf{P}$ should increase towards total independence in the continuum limit. It is evident that neither the FF nor the IF can be exactly boost-invariant *on a lattice* according to the criterion given above. In Chapter $\langle 5 \rangle$, we shall even see that, ironically, the IF is boost-invariant to a much higher accuracy than is the FF in a practical computation. Why, then, is this phenomenon nowhere described? In the FF literature, no distinction is made, between boost-invariance under CO-ORDINATE boosts and invariance under the PHYSICAL BOOSTS discussed above. In the FF, a CO-ORDINATE BOOST in three direction transforms a frame with $\bar{e}^{(0)} = e^{(+)}$ into a kinematically equivalent frame, i.e. it leaves the quantisation surface $e^{(+)}$ intact. In the IF, a CO-ORDINATE BOOST renders the form of the 4-momenta as well as the expression for the commutation relations more cumbersome. But: physics is not changed by choosing co-ordinates which are not adapted to the physical

problem in the same way that physics is not changed if a spherically symmetric system is expressed in elliptical co-ordinates. The only penalty for using the "wrong" co-ordinates is a practical one: cumbersome expressions, as we have discussed above. In particular the commutation relations(3.12) look ugly when expressed in a frame whose instant of time does not coincide with the quantisation surface. The reason for why CO-ORDINATE BOOSTS are kinematical in the FF is the same reason for why these boosts are kinematical in the IF, too: *Co-ordinate boosts are always kinematical*. They do not involve interactions— by definition.

The invariance of the FF under *kinematical* boosts has some unpleasant side-effects. The cut-off $\Lambda = \pi/a^-$ and the lattice size $\mathfrak{L}^-$ are irrelevant quantities in the FF. In two space-time dimensions there is no continuum limit or infinite volume limit as it does not matter if $\mathfrak{L}^- \rightarrow \infty$ or if $\mathfrak{L}^- \rightarrow 0$: the mass spectrum is always the same; only the number of lattice sites $N_L \stackrel{def}{=} \frac{2\mathfrak{L}^-}{a^-}$ is relevant. The deeper reason behind this is that a space-like boundary vector $\mathfrak{L}$ defines an invariant renormalisation scale $L \stackrel{def}{=} \sqrt{-\mathfrak{L} \cdot \mathfrak{L}}$ whereas a light-like boundary vector $\mathfrak{L} \cdot \mathfrak{L} = 0$ does not: the FF is scale invariant in two space-time dimensions. We shall make use of this peculiar property on several occasions in this thesis.

7 The Front Form and ε Co-ordinates

7.1 A Sketch of the Problem

Often, the FF is defined as the limit $\varepsilon \rightarrow 0$ in so-called ε -co-ordinates [69–71], a co-ordinate system othogonal with respect to the Euclidean metric $\delta^{\mu\nu}$ but non-orthogonal with respect to the Minkowski metric $g^{\mu\nu}$. A quantisation

in these co-ordinates⁷ $\bar{x}^\mu(\varepsilon) \stackrel{def}{=} T^\mu_\nu x^\nu$ with

$$T(\varepsilon) = \begin{pmatrix} 1 + \varepsilon & 1 - \varepsilon \\ 1 & -1 \end{pmatrix} \quad (3.38)$$

is canonical (as is the IF quantisation) as long as $\varepsilon \neq 0$. No constraints as in the FF appear. Only for $\varepsilon = 0$ do the FF constraints arise; then the procedure of canonical quantisation has to be replaced by the more sophisticated Dirac-Bergmann quantisation algorithm [73, 74]. At first sight, it seems that these co-ordinates provide a smooth transition between both forms since $\bar{x}^0(0) = x^+$ and $\bar{x}^3(0) = x^-$. The ε co-ordinates seem to achieve what boosted co-ordinates cannot: a smooth transition between usual co-ordinates and co-ordinates on the light-front. We argue, however, that any construction based on the notion CLOSENESS TO THE LIGHT-FRONT is doomed⁸; it cannot be reconciled with the special theory of relativity. We direct attention to the fact that a particular co-ordinate system does not carry physical information. The actual carrier of information is the boundary 4-vector defining the BCs. For it is this 4-vector which actively breaks Poincaré invariance. The boundary 4-vector $\mathfrak{L} = (0, \mathfrak{L}^3)$ has to be *actively* boosted in terms of ε in order for $\mathfrak{L}$ to lie inside the hyper-plane $\bar{x}(\varepsilon) = 0$ lest non-conserved charges arise. Therefore, the components $\bar{\mathfrak{L}}^\mu(\varepsilon)$ diverge for $\varepsilon \rightarrow 0$ if the lattice size $L \stackrel{def}{=} \sqrt{-\mathfrak{L} \cdot \mathfrak{L}}$ is kept constant. This means that physical information is destroyed for $\varepsilon = 0$ since lattices with different invariant size $2L$ are all mapped onto the same co-ordinates $\bar{\mathfrak{L}}^-(0) = \infty$. Consequently, AMBIGUITIES must be expected and, indeed, we shall encounter them later on in this chapter. Similarly, lattices with different LATTICE SPACING a are all mapped onto the same lattice spacing $\bar{a}^- = \infty$. One might want to avoid these infinities⁹ by an

⁷ There are several conventions in order to define *varepsilon* co-ordinates. Here we use the definition in [70, 71] replacing —without loss of generality— ε/L by ε .

⁸Cf. Chapter (2)

⁹ Solely the infinities — not the ambiguities — can be avoided!

explicit reduction in the *physical* size of the lattice $L \rightarrow L(\varepsilon)$ such that L goes to zero in the limit $\varepsilon \rightarrow 0$. But even then is the transition $\varepsilon \rightarrow 0$ not smooth. For we are now prepared to show that if the physical size of the system becomes infinitesimally small in the limit $\varepsilon \rightarrow 0$, then the resulting Hamiltonian is kinematically equivalent to the IMF rather than to the FF. We shall see that the commutative diagram

$$\begin{array}{ccc}
\mathcal{L} & \xrightarrow[\varepsilon=0]{\text{quantise}} & \hat{P}(\varepsilon=0) = \hat{P}(e^{(+)}) \\
\text{quantise} \downarrow \varepsilon \neq 0 & & \text{subtract left-movers} \uparrow \text{from self-energy } \Delta\omega \\
\hat{P}(\varepsilon) \triangleq \hat{P}(e^{(0)}) & \xrightarrow[\varepsilon \rightarrow 0]{L \rightarrow 0} & \hat{P}(\varepsilon \rightarrow 0) \triangleq \hat{P}(e^{(0)})(\text{IMF})
\end{array} \quad (3.39)$$

only closes if the contributions of left-movers to the self-energy of a right-mover are *explicitly* removed (including the self-energy terms they create). Without explicit destruction of these self-energy terms, the limit $\varepsilon \rightarrow 0$ leads to the IMF and not to the FF. The diagram

$$\begin{array}{ccc}
\mathcal{L} & \xrightarrow[\varepsilon=0]{\text{quantise}} & \hat{P}(\varepsilon=0) = \hat{P}(e^{(+)}) \\
\text{remove left-movers, constrain} \downarrow \text{zero modes} & & \varepsilon \rightarrow 0 \uparrow L \rightarrow 0 \\
\mathcal{L}_R & \xrightarrow[\varepsilon=0]{\text{quantise}} & \hat{P}_R(\varepsilon) \triangleq \hat{P}_R(e^{(0)})
\end{array} \quad (3.40)$$

obtained by *first* removing left-movers and then quantising the theory, commutes however. This means that the limit $\varepsilon \rightarrow 0$ is continuous on the classical level but not on the quantum level. We shall demonstrate this in the next section.

One comment is in order: A connection between the FF and the zero volume IF has been inferred already in the article [70]. **Herein**, we go some steps further, completely separating the small volume effects from a particular co-ordinate system and demonstrating that the $\varepsilon \rightarrow 0$ limit does *not* yield the FF but rather the IMF: the transition from $\varepsilon \rightarrow 0$ to $\varepsilon =$

0 is *discontinuous* —even if the lattice size shrinks to zero in this limit. Quantisation and the $\varepsilon \rightarrow 0$ limit do not commute¹⁰. **Additionally**, we are able to clarify the nature of the constraints arising in this picture. We demonstrate that light-like and space-like BCs are allowed to co-exist in the IMF and that the selection of the former are responsible —even in an IF quantisation— for the elimination of left-movers. **Finally**, there are two points where we disagree with the authors of [70]: Firstly, we show that the size of the lattice cannot be chosen to be arbitrarily small even if only Lorentz-contracted objects which fit into the lattice are described. Secondly, we show that the FF can, in general, *not* be regarded as an effective theory of the IF on small lattices —let alone on large lattices.

7.2 Kinematical Equivalence

The kinematical equivalence of ε quantisation and conventional IF quantisation for $\varepsilon \neq 0$ is swiftly demonstrated using the formalism developed in the last section. The ε co-ordinates defined via $T(\varepsilon)$ are kinematically equivalent to a boosted frame $\check{x}^\mu \stackrel{def}{=} B_\nu^\mu x^\nu$ defined via the co-ordinate boost

$$B(\varepsilon) = \frac{1}{\sqrt{1-v^2}} \begin{pmatrix} 1 & -v \\ v & -1 \end{pmatrix} = \frac{1}{2\sqrt{\varepsilon}} \begin{pmatrix} 1+\varepsilon & 1-\varepsilon \\ \varepsilon-1 & -1-\varepsilon \end{pmatrix} \quad (3.41)$$

with boost-velocity

$$v(\varepsilon) = -\frac{1-\varepsilon}{1+\varepsilon} \quad . \quad (3.42)$$

They are also equivalent to the boosted and rescaled frame $\tilde{x}^\mu = bB_\nu^\mu x^\nu$ where

$$b(\varepsilon) = 2\sqrt{\varepsilon} = 2\sqrt{\frac{1+v}{1-v}} \quad (3.43)$$

¹⁰The only exception to this rule are certain fermionic models such as the massive Schwinger model in the unphysical axial gauge treated in Chapter 4 where the IMF is almost equivalent to the FF.

is the scale transformation. We have sloppily referred to $B(\varepsilon)$ as a "boost" even though B is actually a combination of a boost and a parity transformation. The parity transformation in B originates from our sticking to the wide-spread convention of using x^+ as time co-ordinate and x^- as space co-ordinate. A pure boost would have sufficed if we had used x^- as the time co-ordinate.

The kinematical equivalence of $\bar{x}^\mu$ and $\tilde{x}^\mu$ follows from the fact that the hyper-surfaces $\tilde{x}^0 = 0$ and $\bar{x}^0 = T_\nu^0 x^\nu = 0$ are equivalent. The co-ordinates associated with T or bB define the same hyper-surface $x^0 = vx^3$ moving with velocity v relative to the fixed reference frame x^μ . Other co-ordinates such as $\check{x}^\mu \stackrel{def}{=} B_\nu^\mu x^\nu$ would be kinematically equivalent, too, but we have chosen the frame which does not rescale time x^0 (i.e. $\bar{x}^0 = x^0$ or $F = 1$). Note that both B and b are singular in the $\varepsilon = 0$ limit. The combined transformation bB is singular, too, in the sense that it is not invertible. Only $\bar{x}^\mu(\varepsilon)$ is non-singular for $\varepsilon = 0$. For every finite ε , however, each of these transformations is well-defined and non-singular.

The metric of the $\tilde{x}^\mu$ co-ordinates

$$\tilde{g}^{\mu\nu} = b^2(\varepsilon)g^{\mu\nu} = 4\varepsilon g^{\mu\nu} \quad (3.44)$$

is diagonal but rescaled. The 4-momenta $\bar{P}_\mu(\bar{e}^{(0)})$ of the ε quantisation

$$\bar{P}_0(\bar{e}^{(0)}) = \tilde{P}_0(\tilde{e}^{(0)}) - \tilde{P}_3(\tilde{e}^{(0)}) \quad (3.45)$$

$$\bar{P}_3(\bar{e}^{(0)}) = 2\varepsilon \tilde{P}_3(\tilde{e}^{(0)}) \quad (3.46)$$

are —according to Theorem **(2)**—trivially related to the 4-momenta $\tilde{P}_\mu(\tilde{e}^{(0)})$ from $\tilde{x}^\mu$ quantisation, i.e. quantisation in the non-orthogonal ε co-ordinates can be reduced to a much simpler quantisation in orthogonal, rescaled co-ordinates. Here, we have used the fact that the matrix

$$T' \stackrel{def}{=} T \cdot (Bb(\varepsilon))^{-1} = \begin{pmatrix} 1 & 0 \\ \frac{1}{2\varepsilon} & \frac{1}{2\varepsilon} \end{pmatrix} \quad (3.47)$$

which relates the two frames $\bar{x}^\mu = T'^\mu_\nu \tilde{x}^\nu$, has the contra-gredient

$$T^{\mu'}_\nu = \begin{pmatrix} 1 & -1 \\ 0 & 2\varepsilon \end{pmatrix} \quad . \quad (3.48)$$

7.3 Boosting the Lattice

We require the boundary vector $\mathfrak{L}$ to lie inside the quantisation hyper-plane, i.e. $\bar{\mathfrak{L}} \cdot \bar{e}^{(0)}(\varepsilon) = 0$, in order to avoid non-conservation of P^μ . Cf. Theorem(1). Let $L \stackrel{def}{=} \sqrt{-\mathfrak{L} \cdot \mathfrak{L}}$ be the invariant length of the lattice then and let the boundary 4-vector lie inside the quantisation surface $\bar{x}^0 = 0$, i.e. $\bar{\mathfrak{L}}^0 = \tilde{\mathfrak{L}}^0 = 0$. For L fixed, the periodicity $\bar{\mathfrak{L}}^3 = L/\sqrt{\varepsilon}$, $\tilde{\mathfrak{L}}^3 = 2\sqrt{\varepsilon}L$ diverges in any of the quantisation frames which we have introduced. If we want to obtain a *finite* periodicity $\mathfrak{L}^- = L_c$ in the FF limit $\varepsilon \rightarrow 1$, we have to reduce the physical length L to zero in the FF limit by setting $L(\varepsilon) = L_c\sqrt{\varepsilon}$. Only the explicit reduction of the invariant length in terms of ε is able to convert the space-like $\mathfrak{L} \cdot \mathfrak{L} < 0$ 4-vector $\mathfrak{L}$ into a time like 4-vector $\mathfrak{L}(0) \cdot \mathfrak{L}(0) = 0$. If $\tilde{P}^3$ lies on the mass-shell

$$\tilde{P}_0^2 - \tilde{P}_3^2 = M^2/b^2 = (M/2)^2/\varepsilon \quad (3.49)$$

then the dispersion relation reads

$$\tilde{P}_0 = \sqrt{\tilde{P}_3^2 - M^2/b^2} = \sqrt{\tilde{P}_3^2 - M^2\varepsilon} \quad (3.50)$$

in these co-ordinates and consequently it reads

$$\bar{P}_0 = \frac{1}{2\varepsilon}(\sqrt{\bar{P}_3^2 + M^2\varepsilon} - \bar{P}_3) \quad (3.51)$$

in ε co-ordinates. Eq.(3.51) seems to become gradually identical to the dispersion relation

$$\bar{P}_0 \approx \frac{M^2}{4\bar{P}_3} \quad (3.52)$$

in ε co-ordinates in the $\varepsilon \rightarrow 0$ limit where the square-root may be developed in powers of ε . This argument is deceptive since P^3 can only assume multiple(n) values of

$$\Delta \bar{k}_3 = \frac{\pi}{\bar{\mathfrak{L}}^3} = \frac{\pi}{L} \sqrt{\varepsilon} \quad (3.53)$$

and therefore

$$\bar{P}_0 = \frac{\varepsilon}{2} (\sqrt{n^2 (\pi/L)^2 + M^2}) \quad (3.54)$$

can only be brought into the FF form if L is small– irrespective of ε . The same holds for the $\tilde{x}^\mu$ co-ordinates with momentum lattice spacing and

$$\Delta \tilde{k}_3 = \frac{\pi}{\tilde{\mathfrak{L}}^3} = \frac{\pi}{2L\sqrt{\varepsilon}}, \quad (3.55)$$

of course.

7.4 Conclusion

We conclude that, in spite of contrary claims in the literature [75,76], ε quantisation is completely equivalent to IF quantisation in boosted co-ordinates as long as $\varepsilon \neq 0$. Both quantisations define the same object $P(\tilde{e}^{(0)}) = P(\bar{e}^{(0)})$. The Hamiltonian of ε -quantisation can be seen as the projection $\bar{e}^{(0)} \cdot P$ in $\bar{e}^{(0)}(\varepsilon)$ direction; the Hamiltonian in $\tilde{x}^\mu$ co-ordinates is the projection $\tilde{e}^{(0)} \cdot P$ in $\tilde{e}^{(0)}$ -direction. By no means do ε co-ordinates justify the unfortunate notion of "CLOSENESS TO THE LIGHT-CONE", a notion completely alien to the theory of relativity. In fact, the special theory of relativity is built upon the very absence of such a notion: every observer measures the same speed of light, i.e. the relative velocity between any instant of time and any light-front is the speed of light. The RELATIVE VELOCITY¹¹

$$\Delta v_{kl} \stackrel{def}{=} \frac{k \cdot l}{\sqrt{(k \cdot l)^2 - l \cdot l \, k \cdot k}} \quad (3.56)$$

¹¹or functions of it such as the rapidity

is the proper, Lorentz-invariant, measure of closeness between two hyper-surfaces with normal vectors k and l . If one hyper-surface, say l , is light-like $l \cdot l = 0$ then the relative velocity $\Delta v_{kl} = 1$ is *always* the speed of light, for any space-like hyper-surface k . Cf. $\langle 2 \rangle$.

Only *relative* velocities have physical significance, absolute velocities have not. The ABSOLUTE VELOCITY $v^3 = k^3/k^0$ of k of the lattice depends on the co-ordinate frame: consequently, it is irrelevant. Now, the parameter ε is but a complicated way of expressing the *absolute* velocity v (cf. Eq. (3.42)) of the quantisation surface $\bar{e}^{(0)}(\varepsilon)$ and therefore, any $\varepsilon \neq 0$ is equally close or far to the light-front. A 4-vector lies either *on* the light-front or *off* the light-front. There is no in-between.

8 The Infinite Momentum Frame

8.1 Operational definition of the IMF

As the co-ordinates $\tilde{x}^\mu(\varepsilon)$ and $\tilde{x}^\mu(\varepsilon)$ are kinematically equivalent, we should be able to reproduce all properties of ε co-ordinates in ordinary co-ordinates. This is what we shall do in this section. The limit $\varepsilon \rightarrow 0$ will lead us to the IMF if the invariant lattice size $L = \sqrt{\frac{1+v}{1-v}}L_c = \sqrt{\varepsilon}L_c$ is explicitly reduced to zero. We have to distinguish two inequivalent definitions of the IMF to start with.

Definition 5 *Let $\mathbf{P}$ and M be the momentum and the mass of a particle defined on a momentum lattice with momentum lattice spacing Δk^3 in three-direction. We say that we are in the WEAK INFINITE MOMENTUM FRAME if $|\mathbf{P}|/M \rightarrow \infty$. We say that we are in the STRONG INFINITE MOMENTUM FRAME if $|\mathbf{P}|/M \rightarrow \infty$ and $\Delta k^3/M \rightarrow \infty$.*

The following theorems hold

Theorem 3 *The strong infinite momentum frame is equivalent to choosing an arbitrarily small lattice size $L < 1/M$. Therefore it is unphysical.*

Proof: $\Delta k^3 = \frac{\pi}{L} \rightarrow \infty$ is equivalent to $L \rightarrow 0$. It is well known that the correlation length $a\xi \stackrel{\text{def}}{=} \frac{1}{M}$ has to be larger than the lattice size $2L$ in order for the theory to be physical [4] (except for the non-relativistic limit or QFTs such as the Schwinger model which not require a scaling window).

Theorem 4 *If the EFFECTIVE LATTICE SIZE defined as $N \stackrel{\text{def}}{=} \frac{P^3}{\Delta k^3}$ is finite, then the weak IMF is equivalent to the strong IMF*

Proof: $P^3/M = N\Delta k^3/M \rightarrow \infty$ is equivalent to $\Delta k^3/M \rightarrow \infty$.

In other words: Only if $N = \infty$, has the IMF a chance of being physical (it may or may not, depending on the theory and on the way the limit $N \rightarrow \infty$ is defined).

Theorem 5 *An infinite effective lattice corresponds to a theory which is continuous in the variable $x_B \stackrel{\text{def}}{=} k^3/P^3$ if k^3 is replaced by x_BP^3 .*

Proof: $\Delta x_B = \frac{\Delta k^3}{P^3} = 1/N = 0$.

Theorem 6 *In the IMF, the continuum limit $\Lambda/M \rightarrow \infty$ is meaningless.*

Proof: P^3 has to be smaller than Λ . Thus $P^3/M \rightarrow \infty$ implies $\Lambda/M \rightarrow \infty$. The weak IMF on a finite lattice is therefore ill-defined in general due to the infinities which arise without cut-off. The strong infinite momentum frame is well defined if $N < \infty$ or else if the ultra-violet divergences can be removed by vacuum-subtraction alone. The reader is referred to Chapters <4> and <5> for examples.

8.2 Constraints and Boundary Conditions in the IMF

We have seen that in all reference frames with space-like instances of time are equivalent and that only an explicit reduction of the lattice size $2L$ brings the dispersion relation of ε co-ordinates into FF shape. Therefore, it is natural to ask the question: does the *simultaneous* limit

$$\lim_{\varepsilon, L \rightarrow 0} \hat{P}_\mu(\varepsilon) \quad (3.57)$$

reproduce the FF operators $P(e^{(+)})$? Again, this question has to be answered negatively; again, the specific form of the co-ordinate system does not matter. We are going to show now, that it is possible to construct the FF as a *classical* effective theory of the *strong* IMF without ever moving the instant of time x^0 towards the light-cone. The *quantised FF*, however, is not equivalent to the *quantised* IMF. In what follows we shall be using an orthogonal co-ordinate system x^μ in which the boundary 4-vector $\mathfrak{L}$ takes on the form $\mathfrak{L}_{(3)} = (0, 0, 0, \mathfrak{L}^3)$ with $\mathfrak{L}^3 = L$. We are imposing boundary-conditions in 3-direction and in the transverse direction but we shall not explicitly mention the latter ones. A free scalar field $\varphi(x)$ subjected to space-like periodic boundary conditions in 3-direction

$$\varphi(x + \mathfrak{L}_{(3)}) - \varphi(x - \mathfrak{L}_{(3)}) = \varphi(x + \mathfrak{L}) - \varphi(x - \mathfrak{L}) = 0 \quad (3.58)$$

may be expanded

$$\varphi(x) = \sum_{\mathbf{k}} \frac{1}{\sqrt{(2\pi^3)^3 2\omega(\mathbf{k})}} (a_{\mathbf{k}} e^{-i\omega(\mathbf{k})x^0 + i\mathbf{k}\cdot\mathbf{x}} + a_{\mathbf{k}}^\dagger e^{+i\omega(\mathbf{k})x^0 - i\mathbf{k}\cdot\mathbf{x}}) \quad (3.59)$$

in terms of creation $a_{\mathbf{k}}^\dagger$ and annihilation $a_{\mathbf{k}}$ operators which obey the usual

$$[a_{\mathbf{k}}^\dagger, a_{\mathbf{k}'}] = \delta_{\mathbf{k}, \mathbf{k}'} \quad (3.60)$$

commutation relations. The three-component k^3 takes on the discrete values

$$k^3 = \Delta k^3 n = \frac{\pi}{\mathfrak{L}^3} n \quad (3.61)$$

where n denotes an integer. The free fields φ and $\partial_0 \varphi = \hat{\pi}$ at $x^0 = 0$ are —by construction— a realisation of the commutation algebra **(3.17)**, **(3.18)**.

Inserting the fields $\varphi(0, \mathbf{x})$ into the Hamiltonian

$$H[\varphi] = P_0(e^{(0)}) = \int d^3x \left(\frac{1}{2}(\partial_0 \varphi)^2 + \frac{1}{2}(\nabla \varphi)^2 + \frac{m_0^2}{2}\varphi^2 + \frac{g_0}{4!}\varphi^4 \right), \quad (3.62)$$

yields the expression

$$\begin{aligned} H = & \sum_{\mathbf{k}} [\omega(\mathbf{k}) + \Delta\omega(\mathbf{k})] a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \Delta k^3 \sum_{\mathbf{k}\mathbf{l}\mathbf{m}} \frac{(\Delta k_\perp)^2 g_0}{4(2\pi)^3 4!} \times \\ & \left[4 \cdot \frac{a_{\mathbf{k}}^\dagger a_{\mathbf{l}} a_{\mathbf{m}} a_{\mathbf{k}+\mathbf{l}+\mathbf{m}}}{\sqrt{\omega(\mathbf{k})\omega(\mathbf{l})\omega(\mathbf{m})\omega(\mathbf{k}+\mathbf{l}+\mathbf{m})}} + 6 \cdot \frac{a_{\mathbf{k}}^\dagger a_{\mathbf{l}}^\dagger a_{\mathbf{m}} a_{\mathbf{k}+\mathbf{l}-\mathbf{m}}}{\sqrt{\omega(\mathbf{k})\omega(\mathbf{l})\omega(\mathbf{m})\omega(\mathbf{k}+\mathbf{l}-\mathbf{m})}} + \right. \\ & \left. 4 \cdot \frac{a_{\mathbf{k}+\mathbf{l}+\mathbf{m}}^\dagger a_{\mathbf{l}}^\dagger a_{\mathbf{m}}^\dagger a_{\mathbf{k}}}{\sqrt{\omega(\mathbf{k}+\mathbf{l}+\mathbf{m})\omega(\mathbf{l})\omega(\mathbf{m})\omega(\mathbf{k})}} \right] + \mathcal{R} \end{aligned} \quad (3.63)$$

where

$$\Delta\omega(\mathbf{k}) \stackrel{def}{=} 12\Delta k^3 g_0 \frac{(\Delta k_\perp)^2}{4(2\pi)^3 \omega(\mathbf{k}) 4!} \sum_{\mathbf{l}} \frac{1}{\omega(\mathbf{l})}. \quad (3.64)$$

is the SELF-ENERGY of the virtual particle $a^\dagger(\mathbf{k})|0\rangle$ and $\mathcal{R}$ represents PAIRING terms of the form $\text{Re } a_{\mathbf{k}} a_{-\mathbf{k}}$ and $\text{Re } a_{\mathbf{k}} a_{\mathbf{l}} a_{\mathbf{m}} a_{\mathbf{k}+\mathbf{l}-\mathbf{m}}$. Due to momentum-conservation, $\mathcal{R}$ contains at least one left-moving annihilation operator $a(\mathbf{k})$ if a right-moving operator appears in a product of two or four pairing operators. The momenta components k^i run over the domain $-\Lambda^{(i)} \leq k^i \leq \Lambda^{(i)}$ in multiples of $\Delta k^i = \frac{\pi}{\mathfrak{L}^i}$. These terms are strongly suppressed on small lattices because pairs of left-movers and right-movers are suppressed in this case, as we shall show in this section.

Taking the lattice size L much smaller than the particle masses m_0 and the transverse momenta $\mathbf{k}^\perp$ (i.e. $\Lambda^\perp \ll \Lambda^3 = \Lambda$)¹² renders the momentum lattice spacing $\Delta k^3 = \frac{\pi}{L}$ divergent. This, in turn, allows us to approximate the kinetic energy

$$\omega(k) = \sqrt{m_0^2 + \mathbf{k}^2} \approx |k^3| + \frac{m_\perp^2}{|k^3|} + \mathcal{O}(\mathfrak{L}^3) \quad (3.65)$$

(with the TRANSVERSE MASS $m_\perp^2 \stackrel{def}{=} m_0^2 + k_\perp^2$) and its inverse

$$1/\omega(k) \approx 1/|k^3| + \mathcal{O}(\mathfrak{L}^3) \quad (3.66)$$

in a way that mimics the FF. The deeper reason for why the IMF and the FF have a similar appearance is that any FF lattice necessarily has zero invariant size which implies that all momenta are infinite $n\Delta k \rightarrow \infty$. Claims that the FF —contrary to the IMF— is frame independent are unfounded.

8.3 How To Construct an Effective Hamiltonian

The divergence of the scale $\Delta k^3 = \pi L^{-1}$ allows us to replace the Hamiltonian H by an EFFECTIVE Hamiltonian H_{eff} which describes bound-states with masses much smaller than this scale. To this end we divide the Hamiltonian into the kinetic energy $T = \sum_{\mathbf{k}} \omega(\mathbf{k})$ and the interaction term $H_I \stackrel{def}{=} H - T$.

We are only interested in bound states with momenta $\mathbf{P} \neq 0$ in three-direction. These momenta $\mathbf{P}$ are multiples of Δk^3 . Thus they diverge. For this reason, we may calculate the mass squared operator

$$M^2 = (H + P^3)(H - P^3) \approx 2P^3(H - P^3) = 2P^3(T + H_I - P^3) \quad (3.67)$$

¹²Note that this implies a strongly *anisotropic* lattice since the lattice-spacing in three-direction has to be much smaller than the lattice-spacing in the perpendicular directions

in a way that resembles the FF ($P^3 \approx P^0$ for $v^3 \approx 1$). The kinetic energy minus P^3

$$T - P^3 = \sum_{\mathbf{k}; k^3=0} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} \sqrt{m_0^2 + k_\perp^2} + \sum_{\mathbf{k}; k^3 \neq 0} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} \left[\frac{m_0^2 + k_\perp^2}{2|k^3|} + |k^3| - k^3 \right] \quad (3.68)$$

diverges linearly with $\Delta k^3 \propto L^{-1}$ if at least one particle is present which moves in a direction opposite to $\mathbf{P}$. This holds because the energy $2|k^3|$ of a left mover is order $\mathcal{O}(L^{-1})$ whereas a right-mover's energy is order $\mathcal{O}(L)$. Hence the contribution of LEFT-MOVERS $k^3 < 0$ to the mass squared is order $\mathcal{O}(L^{-2})$, the contribution of RIGHT-MOVERS order unity $\mathcal{O}(L^0)$ and the contribution of particles with $k^3 = 0$ (ZERO MODES) is order $\mathcal{O}(L^{-1})$. Consequently, only right-movers contribute to the *low*-energy spectrum: the kinetic masses of left-movers and zero modes diverge. The effective Hamiltonian H_{eff} may therefore be constructed by excluding modes with divergent energy. There are two ways of excluding divergent modes:

(1) WEAK EXCLUSION or EXCLUSION ON THE QUANTUM LEVEL, i.e. quantising all modes and then removing the divergent modes from the normal-ordered Hamiltonian **(3.63)**.

(2) STRONG EXCLUSION or EXCLUSION ON THE CLASSICAL LEVEL, i.e. exclusion of the divergent fields from the Lagrangian density $\mathcal{L}$ *before* quantisation. The resulting Lagrangian density $\mathcal{L}_R$ and Hamiltonian H_R become non-local quantities except for $m_0 = 0$.

We are going to show now, (1) that only the first approach yields the correct effective Hamiltonian and (2) that the second approach is completely equivalent to the FF approach with periodic, light-like BCs.

8.4 The Classically Effective Hamiltonian

Let us describe the second approach first. The (free) field $\varphi(x) = \varphi_r(x) + \varphi_l(x) + \varphi_0(\mathbf{x}^\perp)$ may be written as a sum of a right-moving field

$$\varphi_r(x) \stackrel{def}{=} \sum_{\mathbf{k}; k^3 > 0} \frac{1}{\sqrt{(2\pi^3)^3 2k^3}} (a_{\mathbf{k}} e^{-ik^3 x^- - i\frac{m^2}{2k^3} x^0 + i\mathbf{k}^\perp \cdot \mathbf{x}^\perp} + a_{\mathbf{k}}^\dagger e^{+ik^3 x^- + i\frac{m^2}{2k^3} x^0 - i\mathbf{k}^\perp \cdot \mathbf{x}^\perp}) \quad (3.69)$$

a left-moving field

$$\varphi_l(x) \stackrel{def}{=} \sum_{\mathbf{k}; k^3 < 0} \frac{1}{\sqrt{(2\pi^3)^3 2|k^3|}} (a_{\mathbf{k}} e^{-ik^3 x^+ + i\frac{m^2}{2k^3} x^0 + i\mathbf{k}^\perp \cdot \mathbf{x}^\perp} + a_{\mathbf{k}}^\dagger e^{+ik^3 x^+ - i\frac{m^2}{2k^3} x^0 - i\mathbf{k}^\perp \cdot \mathbf{x}^\perp}) \quad (3.70)$$

and the zero mode $\varphi_0(\mathbf{k}^\perp)$. Formally we may define the linear PROJECTION OPERATORS Π_r , Π_l and Π_0 which project onto right-movers φ_r , left-movers φ_l or zero modes φ_0 respectively. E.g. $\varphi_r(x) = \Pi_r \varphi(x)$. Now we define the CLASSICALLY EFFECTIVE HAMILTONIAN

$$H_R \stackrel{def}{=} H[\Pi_r \hat{\pi}, (\Pi_r + \Pi_0)\varphi] \quad (3.71)$$

as a functional of right-moving fields $\varphi(x)$ and field-momenta $\hat{\pi}(x)$ only. The reader will have noticed that we have excluded the canonical momentum $\hat{\pi}_0 = \partial_0 \varphi$ of the zero modes but not the zero mode φ_0 itself. The rationale for this is the fact that $\varphi_0 = 0$ would lead to constraints on right-movers via the equations of motion. Imposing the condition $\varphi_0 = 0$ on the classical level is too strong if $g_0 \neq 0$ since it is not compatible with the equations of motion

$$[-\partial_0^2 + \partial_3^2] \varphi = [-\nabla_\perp^2 + m_0^2] \varphi + \frac{g_0}{3!} \varphi^3 \quad (3.72)$$

as they stand. We can see this by applying $\int \frac{dx^3}{2L}$ on both sides. This unveils the constraint

$$(\nabla_\perp^2 - m_0^2) \varphi_0 = \frac{g_0}{3!} \int dx^3 \varphi^3 \quad (3.73)$$

where we have used that (a) the integral over the partial derivative

$$\int_{-L}^L dx^3 \partial_3 \partial_3 \varphi(x^3) = \partial_3 \varphi(L) - \partial_3 \varphi(-L) = 0 \quad (3.74)$$

vanishes with periodic BCs and (b) the temporal derivative $\partial_0 \varphi_0$ vanishes by assumption. This consistency constraint has the same form as the constraint which arises in the Dirac-Bergmann quantised FF, cf. Eq. (3.21). If we had set $\varphi_0 = 0$ then the constraint (3.73) would have been a constraint on φ_r . Retaining $\varphi_0 \neq 0$, however, makes this a constraint on the zero mode. This constraint can in principle be solved for the zero mode. The kinetic operator $T - P^3$ reads

$$T - P^3 = \int d^3x \frac{1}{2} (\partial_+ \varphi)^2 + \frac{1}{2} \int d^3x \varphi \left[-\nabla_\perp^2 + m_0^2 \right] \varphi \quad (3.75)$$

in terms of the fermionic fields. Projecting out the left-moving field φ_l and the zero-mode velocity $\partial_0 \varphi_0$ and using the constraint (3.73), we obtain

$$\begin{aligned} T - P^3 &= \int \frac{d^3x}{2} \left[(\partial_+ \varphi_r)^2 + (\nabla_\perp \varphi_r)^2 + m_0^2 \varphi_r^2 \right] + \int \frac{d^2x}{2} \varphi_0 \left[\partial_0^2 - \nabla_\perp^2 + m_0^2 \right] \varphi_0 \\ &= \int \frac{d^3x}{2} \left[(\partial_+ \varphi_r)^2 + (\nabla_\perp \varphi_r)^2 + m_0^2 \varphi_r^2 \right] + \int \frac{d^3x}{2} \frac{g_0}{3!} \varphi^3 \quad . \end{aligned} \quad (3.76)$$

This demonstrates that the zero modes no longer contribute to the kinetic mass except for an indirect contribution via interactions. The right-movers obey the commutation relations

$$[\varphi_r(0, \mathbf{x}), \varphi_r(0, \mathbf{y})] = \frac{1}{2i\partial_x^3} \delta(\mathbf{x} - \mathbf{y}) = \frac{1}{2i\partial_3^y} \delta(\mathbf{x} - \mathbf{y}) \quad (3.77)$$

which means that $2\partial_3 \varphi_r|_{x^0=0} = 2\partial_- \varphi_r|_{x^0=0}$ plays the rôle of the canonical momentum of φ_r . Now we neglect the order $\mathcal{O}(L^{-1})$ contribution $\partial_+ \varphi_r$ and

end up with the CLASSICALLY EFFECTIVE HAMILTONIAN

$$H_R = P^3 + \int d^4x \delta(x^0) \left[\frac{1}{2}(\nabla_\perp \varphi)^2 + \frac{1}{2}m_0^2 \varphi^2 + V(\varphi) \right] \quad (3.78)$$

with

$$[\varphi(x^0 = 0, \mathbf{x}), \varphi(y^0 = 0, \mathbf{y})] = \frac{1}{2i\partial_3} \delta(\mathbf{x} - \mathbf{y}) \quad (3.79)$$

which has now exactly the same form as the FF Hamiltonian $P^-(e^{(+)}) = 2P_+(e^{(+)})$ (3.19) except that x^+ and x^0 are interchanged. Alternatively, the reader will easily convince her/himself that the Dirac-Bergmann quantisation of the Lagrangian density

$$\mathcal{L}_R \stackrel{def}{=} -2\partial_3 \varphi \partial_0 \varphi - (\partial_3 \varphi)^2 - \left[\frac{1}{2}(\nabla_\perp \varphi)^2 + \frac{1}{2}m_0^2 \varphi^2 + V(\varphi) \right] \quad (3.80)$$

leads to the Hamiltonian H_R with the commutation relations (3.79) and the zero-mode constraint (3.73). Please note that this Lagrangian density—which results from the elimination of high-energy modes on the classical level—is not equivalent to the physical Lagrangian density $\mathcal{L}$ we started with. This completes our construction.

If we had imposed ANTI-PERIODIC BCs on the field φ the discussion would have become much easier because anti-periodic BCs eliminate the zero mode. The principal result, however, would have been the same: *the elimination of high-energy modes on the classical level reproduces the FF.*

8.5 The Quantum Effective Hamiltonian

We define the QUANTUM EFFECTIVE HAMILTONIAN

$$H_{\text{eff}} \stackrel{def}{=} (\Pi_r + \Pi_0) H[\hat{\pi}, \varphi] \quad (3.81)$$

by removing left-movers through a weak constraint, i.e. instead of the strong constraint $\varphi_r = 0$ we merely impose the condition $\varphi_r | \text{eff} \rangle \equiv 0$ in order to

restrict the total Fock-space to states $|\text{eff}\rangle$ which do not contain left-movers. This Hamiltonian is guaranteed to be the low energy effective Hamiltonian for the strong IMF since eigenstates of the IMF Hamiltonian which mix left-movers and right-movers have infinite energy in the limit $\Delta k^3 \rightarrow \infty$. The same does not hold true for the classically effective Hamiltonian H_R which is formally equivalent to the FF Hamiltonian. Even though these two constructions, H_R and H_{eff} , are equivalent on the classical level, they are no longer equivalent on the quantum level. The second construction—which is equivalent to the FF—does not have the same low-energy mass spectrum as the IMF. It can therefore not be called effective Hamiltonian. This is so because H_R fails to reproduce the correct self-energy term $\Delta\omega(\mathbf{k})$. The self-energy receives equally important contributions from normal-ordering of left-moving particles and right-moving particles.

$$\begin{aligned}\Delta\omega(\mathbf{k}) &= 12\Delta k^3 g_0 \frac{(\Delta k_\perp)^2}{4(2\pi)^3 |k^3| 4!} \left[\sum_{\mathbf{l}; l^3=0} \frac{1}{\omega(0, \mathbf{l}^\perp)} + \sum_{\mathbf{l}; l^3 \neq 0} \frac{1}{|l^3|} \right] \\ &= \Delta\omega_r(\mathbf{k}) + \Delta\omega_l(\mathbf{k}) + \Delta\omega_0(\mathbf{k}) \quad ,\end{aligned}\tag{3.82}$$

where $\omega_r(\mathbf{k})$, $\omega_l(\mathbf{k})$ and $\omega_0(\mathbf{k})$ are defined as the contribution to $\omega(\mathbf{k})$ stemming from left-movers, right-movers and the zero mode respectively.

Removing left-movers from the fields (as opposed to removing them from the Hamiltonian after normal ordering) fails to reproduce the correct self-energy term and, consequently, the FF Hamiltonian can *not* be considered as an effective Hamiltonian to the IMF. It describes a completely different theory. With these results we can now answer the question, whether quantisation in ε co-ordinates smoothly becomes identical to FF quantisation in the limit $\varepsilon \rightarrow 0$, as long as the invariant lattice size L becomes zero in this limit: The answer is no. As the results of ε quantisation and IF quantisation are completely equivalent, the only relevant quantity is the invariant lattice

size L . **The limit $L \rightarrow 0$ leads us to the IMF, not to the FF unless the FF constraints are imposed by hand.** We have seen, however, that the FF mass operator cannot be considered as an effective IMF mass operator. *Therefore, the limit $\varepsilon \rightarrow 0$, $L \rightarrow 0$ is discontinuous.* This negative result holds for bosons only. The fermionic self-energy of a right-moving particle in QED(1 + 1) receives contribution from right-moving particles only. We shall see this in Chapter [5](#) (the FF Hamiltonian for Fermions is derived in [\[77, 78\]](#)). The self-energy contributions of left-movers in the bosonic case, in contrast, are far from being negligible. In fact, the mass-spectrum of the scalar φ^4 theory is dominated by the self-energy contribution because physical states are almost purely one-particle states¹³. (See Chapter [6](#)).

Another reason for why one should remove left-movers from the Hamiltonian and not from the fields is that two fields $\varphi_r(\mathbf{x})$ at different positions do not commute due to [\(3.77\)](#) which means that they are causally connected. This is part of a violation of microcausality which shall be treated in the next section: **the classical effective theory strongly violates microcausality.**

The explicit removal of left-movers from the fields $\varphi(x)$ is intimately connected with the implementation of light-like BCs. If only right-moving modes are allowed, a field which is periodic in a space like direction with periodicity L is periodic also in the light-like direction¹⁴ with boundary 4-vector $\mathcal{L}' = (L/2, 0, 0, -L/2)$. If both left-moving modes and right-moving modes are present in $\varphi(x)$, then this field is only periodic in spatial direction.

In conclusion: *the FF is an unphysical approximation of yet another unphysical approximation: the IMF.* The FF is neither an effective theory of

¹³In fact, the self-energy $\Delta\omega$ is responsible for the mean-field part of the critical exponents whereas the tiny logarithmic corrections stem from many-body effects.

¹⁴This statement holds for $|x^0| < L$ with arbitrary accuracy in the limit $L \rightarrow 0$. The *physical* reason for the restriction $|x^0| < L$ will be discussed in the next subsections.

the IF nor is it an effective theory of the IMF, except for the non-relativistic limit where all relativistic forms coincide.

8.6 Some Comments on Zero Modes in the FF

Light-like periodic BCs are incompatible with relativistic equations of motion as they stand

$$4\partial_+\partial_-\varphi(x) = (m_0^2 - \nabla_\perp^2)\varphi(x) + \frac{\partial}{\partial\varphi}V(\varphi(x)) \quad ; \quad (3.83)$$

they have to be supplemented by a constraint on the zero-modes $\tilde{\varphi}(x^+) = \int \frac{dx^-}{2\mathfrak{L}^-}\varphi(x^+, x^-)$ which follows from integrating these equations

$$0 = (m_0^2 - \nabla_\perp^2)\tilde{\varphi}(x^+) + \int_{-\mathfrak{L}^-}^{+\mathfrak{L}^-} \frac{dx^-}{2\mathfrak{L}^-} \frac{\partial}{\partial\varphi}V(\varphi(x)) \quad (3.84)$$

with respect to x^- . This constraint corresponds to the IF constraint **(3.73)** which arises if one tries to construct an effective Hamiltonian on the level of fields. The constraint **(3.84)** does not arise from the FF as such nor is it connected with the quantisation surface e^{quant} ; being another by-product of light-like BCs it is independent of the quantisation surface. If the field $\varphi(x)$ is subjected to anti-periodic BCs, the constraint **(3.84)** does not arise. There is no justification whatsoever for using a constraint which results from the unphysical character of periodic BCs (see Chapter **(6)**) in order to extract physics such as spontaneous symmetry breaking.

Our FF construction on a space-like quantisation plane sheds some light on why this constraint appears to describe spontaneous symmetry breaking. In the framework of our construction, this constraint arises out of the requirement that the kinetic energy of zero-modes vanish. This, in turn, requires that the velocity $\partial_0\varphi_0$ of the zero-mode vanish as well. These are, perchance, the ingredients of Landau's celebrated *mean field* treatment of the scalar

φ^4 theory. The Landau mean field theory uses a field which is constant in terms of x^0 and x^3 (the same condition as in (3.73)). *Coincidentally*, this is exactly the property of the zero mode subjected to light-like BCs. This finding explains a paradox which would otherwise be quite puzzling: a QFT in an infinitesimal volume¹⁵ would seem to describe, at least approximately so, spontaneous symmetry breaking which *solely* occurs in the infinite volume limit. The real danger of the FF is not that the FF is consistently wrong in all circumstances. Worse than that. Sometimes, as in this case, the FF comes close to reality for the wrong reasons or because the FF is extremely ambiguous (due to the omission of half of the degrees of freedom). A mere change of BCs, for instance, may drastically change the outcome even in situations where this should not happen: Anti-periodic BCs do not induce the constraint (3.84) after all.

9 The Breakdown of Causality in the Front Form

9.1 The Causality Region

There are problems even worse than those encountered before: *Light-like BCs are incompatible with microcausality*. For the sake of a comparison with [79] we do this part in $1 + 1$ dimensions. We may restrict ourselves, as [79], to a free field theory since the violation of microcausality arises already for free fields. It should be evident that the major result will persist in $3 + 1$ dimensions. The breakdown of microcausality is intimately connected to the fact that light-like boundary conditions, contrary to space-like boundary conditions, are *not* invariant under parity or time-reversal. A boundary vector

¹⁵A vanishing physical volume is always implied by light-like BCs




Figure 3.4: The shrinking of the causality region $\mathcal{C}$ in a fixed frame




Figure 3.5: The shrinking of the causality region $\mathcal{C}_\varepsilon$ in ε co-ordinates

$\mathfrak{L}$ with $\mathfrak{L}^+ = 0$ and $\mathfrak{L}^- \neq 0$ is transformed into $\mathfrak{L}'$ with $\mathfrak{L}'^- = 0$ and $\mathfrak{L}'^+ \neq 0$. This symmetry can not be restored (except for the non-relativistic limit) even if $\mathfrak{L}^- \rightarrow \infty$.

Relativistic equations of motion have always causal and acausal solutions. Hence it does not suffice to find any solutions to the relativistic equations of motion at all. **A Hamiltonian has to generate causal solutions only.** A necessary condition for relativistic field theories to be causal ist that any commutator of local operators must

$$[\mathcal{O}_1(x), \mathcal{O}_2(y)] \quad (3.85)$$

vanish for space-like $(x - y)^2 < 0$ distances $x - y$. For our purposes it will suffice to take $\mathcal{O}_1, \mathcal{O}_2 = \varphi$. Non-interacting scalar operators $\varphi(x)$ quantised in the IF fulfill this requirement since the Klein-Gordon propagator

$$i\Delta(x - y) = [\varphi(x), \varphi(y)] = \int \frac{dp^3}{2\pi\omega(\mathbf{p})} \sin(-px) \quad (3.86)$$

vanishes for space-like distances. This fact is not changed if we impose *space-like* periodic BCs $\varphi(t, -L) = \varphi(t, +L)$ as long as we restrict ourselves to the finite region

$$\mathcal{C} \stackrel{def}{=} \{x | -\mathfrak{L}^3 < x^3 < \mathfrak{L}^3; -\mathfrak{L}^3 < x^0 < \mathfrak{L}^3\} \quad (3.87)$$

which we shall refer to as the CAUSALITY REGION illustrated in **Fig. 3.4**. For times $|x^0| < \mathfrak{L}^3$, the periodic propagator coincides exactly with the propagator in an infinite volume since the effects of the boundary conditions cannot propagate faster than the speed of light. This can easily be seen by inspection of the periodic propagator

$$\Delta(x - y; \mathfrak{L}^3) = \sum_{p^3} \frac{1}{\omega(\mathbf{p})(2\mathfrak{L}^3)} \sin(-px) = \sum_n \Delta(x^0 - y^0, x^3 - y^3 + 2\mathfrak{L}^3 n) \quad (3.88)$$

Contrary to this, the FF propagator with light-like periodicity $\mathfrak{L}$

$$\Delta_{\text{FF}}(x, \mathfrak{L}^-) = \sum_{p_-} \frac{1}{2p_- \mathfrak{L}^-} \sin(-px) \quad (3.89)$$

does *not* converge against $\Delta(x)$ in the region $-\mathfrak{L}^- < x^- < \mathfrak{L}^-$, $-\mathfrak{L}^- < x^+ < \mathfrak{L}^-$. How can this be? Firstly, the projection

$$\sum_n \Delta(x^+ - y^+, x^- - y^- + 2\mathfrak{L}^- n) \quad (3.90)$$

of Δ onto its periodic part is divergent unless we *explicitly* subtract the $p_- = 0$ mode responsible for this divergence

$$- \frac{1}{2p_- \mathfrak{L}^+} \sin\left(\frac{m^2}{4p_-}\right) \quad (3.91)$$

from the propagator¹⁶. Secondly, let us assume that one can approximate an integral of the type $\int_{-\infty}^{\infty} dk f(k)$ by the Riemann sum $\sum_n \Delta k f(n\Delta k)$ such

¹⁶It is precisely the FFConstraint (3.84) which avoids such divergences since this constraint reads $\varphi(x) = 0$ for $g_0 = m_0 = 0$. Cf. the discussion of zero-modes on page 89.

that this series converges against the integral. Then there is no guarantee that the same integral

$$\int_{g^{-1}(-\infty)}^{g^{-1}(+\infty)} g'(l) dl f(g(l)) \quad (3.92)$$

can be approximated by a (equidistant) sum if the co-ordinate transformation $g : k = g(l)$ contains a singularity. Now, the transformation $g : k^+ = g(k^3) = \omega(k^3) + k^3$ which transforms k^3 into k^+ is indeed singular since $dk^3 = \frac{dk^+}{k^+} \omega(k^3)$ diverges at $k^+ = 0$. Furthermore, the integrand $\sin \left[\frac{m^2 x^+}{2k^+} \right]$ oscillates violently in terms of k^+ , close to $k^+ = 0$ which is incompatible with Riemann integrability. An explicit calculation confirms this: Since the FF propagator transforms as

$$\Delta_{\text{FF}}\left(\frac{x^+}{\kappa}, x^- \kappa; \mathfrak{L}^- \kappa\right) = \Delta_{\text{FF}}(x^+, x^-; \mathfrak{L}^-) \quad (3.93)$$

under CO-ORDINATE BOOSTS it does not make sense to say $\mathfrak{L}^-$ is large or small. We can make it arbitrarily large or small by means of a co-ordinate boost. Therefore we can always assume that it is arbitrarily small such that $\Delta k^- \gg m$. Thus in the neighbourhood of the quantisation surface (i.e. x^+ sufficiently small)

$$\Delta_{\text{FF}}(x^+, x^-; \mathfrak{L}^-, m) \approx \Delta_{\text{FF}}(x^+, x^-; \mathfrak{L}^-, 0) = -\frac{1}{4} \text{sgn}(x^-) + \frac{x^-}{4\mathfrak{L}^-} \quad (3.94)$$

This means a full-scale violation of microcausality since Δ_r does not disappear in the space-like region $x^+ < 0, x^- > 0$; it disappears in the time-like region instead. The fact that Δ_r does not vanish for arbitrarily small x^+ in the space-like region close to light-front $x^+ = 0$ cannot be changed by boosts, and therefore microcausality is violated for large $\mathfrak{L}^-/\kappa$ also. This may be reexpressed in a somewhat sloppy way: Since there is no preferred frame in the FF, we can make every particle behave as if it were massless to

any accuracy by choosing a frame with almost infinite momentum spacing Δk_- where particle-masses can be neglected. Now, massless left-movers cannot be quantised in the FF-1 without a full-scale violation of microcausality. Consequently, the FF violates causality irrespective of particle masses.

9.2 The Instant Form Perspective

Our FF construction on a space-like surface (in section $\langle 8.2 \rangle$) is able to illustrate these arguments. Inside the causality region, the massive propagator $\Delta(x, m) \approx \Delta(x, 0)$ coincides with the massless propagator up to order $\mathcal{O}(L^2)$ corrections. We divide the IF field $\varphi(x^0, x^3)$ into the right-moving part φ_r and the left-moving part $\varphi_l(x^0, x^3)$ which contains all operators which create or annihilate left-movers and $\varphi_0(x^0)$ which contains the IF zero-modes. Since φ_r , φ_l and φ_0 commute mutually, it follows that

$$[\varphi(x), \varphi(y)] = [\varphi_r(x), \varphi_r(y)] + [\varphi_l(x), \varphi_l(y)] + [\varphi_0(x), \varphi_0(y)] \quad (3.95)$$

and, therefore, the causal propagator

$$\begin{aligned} i\Delta(x - y; \mathfrak{L}^3, m) &= i\Delta_l(x - y; \mathfrak{L}^3) + i\Delta_0(x^0 - y^0; \mathfrak{L}^3) + i\Delta_r(x - y; \mathfrak{L}^3) \\ &\approx -\frac{1}{4} [\text{sgn}(x^+) + \text{sgn}(x^-)] + \mathcal{O}(L^2) \end{aligned} \quad (3.96)$$

is the sum of three acausal parts where

$$\Delta_l(x^0, x^3) = \Delta_r(x^0, -x^3) \approx \Delta_{\text{FF}}(x^+, x^-) \quad (3.97)$$

is just the parity-reversed massless propagator Δ_r and

$$\Delta_0(x^0) = \frac{1}{2m\mathfrak{L}^3} \sin(-mx^0) \approx -\frac{x^0}{2\mathfrak{L}^3} = -\frac{1}{4\mathfrak{L}^3}(x^+ + x^-) \quad . \quad (3.98)$$

The last part only holds for $|t| < \mathfrak{L}^3$. This nicely illustrates how a balanced, parity-invariant combination of left-movers and right-movers ensures microcausality even though microcausality is strongly violated by both Δ_l and Δ_r .

The mathematical reason for why $\Delta(x)$ vanishes in the space-like region is that there is a parity reversed partner $-k^3$ for every k^3 : both contributions cancel. Such a cancellation is rendered impossible in the FF-1 since neither light-like BCs nor light-like quantisation surfaces can be parity-invariant.

Elimination of left-movers on the classical level (cf. Sec. [\(8\)](#)) yields the same unphysical propagator as FF quantisation. The restriction of micro-causality to the causality region $\mathcal{C}$ has salient consequences of the STRONG IMF as well. In the limit $L \rightarrow 0$ the causality region of the IMF collapses into a point, as illustrated in [Fig. 3.4](#). This means, that the strong IMF is only capable of describing space-time events that take place in an infinitely small space-time region. In particular, this means that only instantaneous events may be described in this frame. In an arbitrarily small region, a massive propagator looks like a massless propagator. The propagation of free particles is almost massless. The causality region of the strong IMF defined in ε co-ordinates, however, does not collapse into a space-time point. It collapses into a space-time line instead as illustrated in [Fig. 3.5](#). Inside the causality region, these results, obtained for free fields, are likely to hold for interacting fields, too, as long as $g/\Delta k^3$ is order $\mathcal{O}(L)$ or, equivalently, if g/m is order $\mathcal{O}(1)$.

These results show why the IMF works so well, perturbatively, in order to formulate Feynman's parton model. In the naïve parton model, the time it takes for the electron in order to scatter off a quark is arbitrarily short and the entire process may take place in an arbitrarily short volume. This legitimises the use of the strong IMF even though the strong IMF is only capable of describing events which take place arbitrarily fast and inside an arbitrarily small volume. If vacuum effects are explicitly excluded, the IMF may even be used in order to derive the leading order DGLAP equations:

these equations do not contain loops, i.e. they are classical equations except for the usage of the running coupling constant. In the light of our results on the IMF, it is our conviction that a successful *non-perturbative* computation of QCD distribution functions in the strong IMF is not possible. We would also draw attention to the fact that the propagator Δ_r propagates free fields φ faster than the speed of light; a fact which is, of course, intimately related to the breakdown of microcausality. So is the unphysical possibility of time-travel which we have discussed in the last section.

9.3 Light-Like Boundary Conditions in the Instant Form

The periodic free field $\varphi_r(x)$ (or $\varphi_l(x)$) Eq. (3.69) is compatible with light-like periodic BCs $\mathfrak{L}'$ up to order $\mathcal{O}(L)$

$$\varphi(x + \mathfrak{L}') - \varphi(x - \mathfrak{L}') \approx 0 \quad (3.99)$$

inside the causality region. Here, $\mathfrak{L}' = (L/2, -L/2)$. The complete field $\varphi(x)$, however, does not fulfill this requirement. Excluding φ_l and φ_0 is, therefore, equivalent to the introduction of light-like BCs if a free theory is concerned. Inside the causality region, this connection can be expected to remain valid even for an interacting theory, as long as the coupling constant g is small when compared to the scale Δk^3 .

10 The Limit of Infinite Light-Like Volume

We have shown that the FF propagator $\Delta_{\text{FF}}(x)$ is completely unphysical if the light-like length $\mathfrak{L}^-$ is finite. So are time-ordered propagators or Feynman propagators. Yet FF perturbation theory, in the limit $\mathfrak{L}^- \rightarrow \infty$, seems to be equivalent to covariant perturbation theory at least in the case of some simple Feynman diagrams [51, 80]. What happens in the limit $\mathfrak{L}^- \rightarrow \infty$?

In this limit,

$$\lim_{\mathfrak{L}^- \rightarrow \infty} \Delta_{\text{FF}}(x, \mathfrak{L}^-) = \int \frac{dk_-}{2\pi k_-} \sin(-k \cdot x) \quad (3.100)$$

one is faced with the problem that the quantity

$$1/k_- = \mathcal{P}[1/k_-] + \delta(k_-) \alpha(x^+) \quad (3.101)$$

is ambiguous¹⁷. There is an infinite number of possible prescriptions — parametrised by the arbitrary function $\alpha(x^+)$ — in order to properly define what $1/k_-$ means. Setting $\alpha(x^+) = 0$, for instance, means choosing the PRINCIPAL VALUE PRESCRIPTION $\mathcal{P}[1/k_-]$. This means that the expression Δ_{FF} may be anything: The propagator depends crucially on the choice of PRESCRIPTION. In other words: the FF theory has an infinite number of free parameters. Such a theory is not sensible. The freedom of choosing physically inequivalent prescriptions for $\mathfrak{L}^- = \infty$ signals a defect of the FF reflecting the fact that the limit

$$\lim_{\mathfrak{L}^- \rightarrow \infty} \Delta_{\text{FF}}(x; \mathfrak{L}) \quad (3.102)$$

does not exist except for the FF CAUSALITY REGION¹⁸ $x^+ = 0$ where all prescriptions yield the same propagator. Of course, this ambiguity is intimately related to the fact that two light-like planes are needed to define a relativistic initial value problem.

One might argue that the prescription is not ambiguous but determined as the prescription which yields the correct physical result. Therefore, it seems that one need not worry about the prescription as long as the correct propagator is obtained. There is a flaw in this argument (a *petitio principii*),

¹⁷ In fact, this expression does not even exhaust the set of all possible prescriptions. It suffices, however, to make clear that there is an infinite number of possible prescriptions.

¹⁸ Cf. **Fig. 3.5**.

however. **There is reason to worry.** By choosing the correct propagator, we feed the theory with the data we would like to obtain. *The correct value of α does not follow from the FF quantisation procedure.* IF quantisation, in contrast, unambiguously defines the propagator without additional *ad hoc* assumptions. The fact that a prescription may be found which reproduces the correct IF propagator¹⁹ does not mean that the FF and the IF are equivalent for $\mathfrak{L}^- = \infty$. If they were then the FF would also be "equivalent" —in the same sense— to an arbitrary number of completely unphysical theories.

The choice of prescription is often referred to as *regularisation* in the literature. We do not agree with this misleading parlance. In the IF, the propagator $\Delta(x)$ is defined *without* regularisation in a *free* QFT contrary to other quantities such as the energy H which has to be regularised even if a free QFT is concerned. Furthermore, in the IF, a regularisation does *not* introduce ambiguities into quantities which need to be regularised as long as the correlation length is sufficiently large²⁰ and as long as one is dealing with a renormalisable theory. Similarly, in the FF the problem is *not* that the expression $1/k_-$ would require regularisation —it does not. The problem is that Δ_{FF} is *ambiguous* yet *finite*. Calling the choice of α a regularisation is a misnomer.

It is interesting to establish a connection between the α -ambiguity and ε co-ordinates. We have seen, that the causality region of ε co-ordinates collapses into the line $x^+ = 0$. Cf. **Fig. 3.5**. A boost $B(v = 1)$ even collapses the universe onto the subspace $x^+ = 0$. The net result is the same, even though a boost $B(v)$ with $v = 1$ is singular and $T(\varepsilon = 0)$ is not: Quantisation with respect to the transformed co-ordinates unambiguously describes

¹⁹This may no longer be true in the presence of interactions

²⁰ More precisely, ambiguities are present but related to irrelevant operators. The influence of these operators is strongly suppressed in a system close to the critical point.

the collapsed subspace $x^+ = 0$ only. Outside this region, neither of both approaches is able to provide (unambiguous) predictions.

11 Situations Where the FF May Be a Good Approximation

11.1 The Non-relativistic Limit

In the non-relativistic limit $c \rightarrow \infty$, all space-like and light-like hyper-planes

$$x^0 - \frac{v}{c^2}x^3 = 0 \quad (3.103)$$

collaps onto the unique, non-relativistic quantisation surface $x^0 = 0$. The FF and the IF coincide in this limit. The question arises, if the FF remains accurate close to the relativistic limit. The *zero-hypothesis* is that the FF-corrections to the non-relativistic limit are not accurate.

11.2 Perturbation Theory With External Input

Only free propagators go into perturbation theory. The ambiguity of free propagators in the FF allows to specify the parameter of ambiguity α such that the free time-ordered propagators coincide with the corresponding IF propagators. Therefore, it is possible to correctly describe some elementary diagrams of perturbation theory but *only if* external input is used: One has to choose amongst an infinity of possible prescriptions in order to regularise $1/k_-$ singularities, i.e. one has to fix an infinity of free parameters α . FF quantisation itself does not specify the prescription α .

11.3 Theories Without Vertices Which Connect at Least Four Bosons

We shall see in Chapter [5](#) that fermionic self-energies need not *necessarily* be inaccurately described by the FF because they do not contain left-movers. The FF may or may not come close to the correct solution to a QFT which does not contain vertices which connect at least four bosons (except for the ultra-relativistic limit). QCD, however, contains a four-gluon vertex.

11.4 Accidental Cases

There is a case where the FF *accidentally* reproduces the exact mass spectrum of a QFT. The massless SCHWINGER MODEL, treated in [4](#), is equivalent to the CHIRAL SCHWINGER MODEL in the unphysical limit $L \rightarrow 0$. The chiral Schwinger model does not contain left-movers. We shall see in [4](#) how these features lead to the fact that the mass spectrum of the massless Schwinger model is exactly described in the IF, the IMF and the FF.

11.5 Phenomenology

The FF may be used as a bookkeeping device in phenomenology. The zero-hypothesis is that every phenomenological problem which can be easily treated in the FF, may equally well be treated in the IF if the volume is taken to be sufficiently small. We do not know of any example where the FF is useful but the IF is not.

12 Other Relativistic Forms

Every relativistic form represented in [Tab. 3.1](#) is a valid quantisation surface except for the FF. May these forms be useful in order to quantise QFTs?

There is reason to doubt this. Firstly, the classification given in [65] does not take discrete symmetry into account. Yet time reversal and parity symmetry are as important as the continuous part of the Lorentz group. There is only one form whose stability group comprises time-reversal: the instant form. The quantisation surface $x^0 = 0$ is the only quantisation surface from **Tab. 3.1** which is invariant under time reversal. Important theorems such as the PCT theorem naturally apply in the IF and only in the IF. In every other form, time reversal invariance can be realised dynamically at best. The FF is worst in this respect since even parity invariance is destroyed.

Another aspect, which is neglected in the literature is the aspect of implicit time dependence. For instance, even though the boosts J_{03} and $J_{\perp,-}$ are in the FF stability group, they do not commute with the FF time evolution operator

$$U_+(x^+) \stackrel{def}{=} \exp(-i\hat{P}_+x^+) \quad . \quad (3.104)$$

This means that these operators have an *explicit* time dependence: $J_{03} = J_{03}(x^+)$. Such a symmetry is not very helpful. In particular, the diagonalisation of the Hamiltonian is not facilitated. We therefore introduce the notion of the STABILITY GROUP PROPER which consists of operators which do not explicitly depend on time and the EPHEMERAL STABILITY GROUP which consists of operators which do not commute with the time evolution operator. Geometrically, this difference of usefulness may be visualised as follows. In the IF, a valid quantisation surface $x^0 = 0$ is transported into another valid quantisation surface $x^0 = T^0$ by means of the operator $U_0(T^0)$ of temporal evolution. Any quantisation surface obtained by means of a temporal evolution is again symmetrical with respect to the full IF stability group. The IF is the only form which boasts this feature. The light-front, for instance, is transported onto a quantisation surface which is no longer symmetrical

Relativistic form	Quantisation-surface	Algebra of the stability group proper
Front form	$x^+ = e^{(+)} \cdot x = 0$	$\mathbf{P}_\perp, \mathbf{J}_{12}; P_-$
Instant form	$x^0 = e^{(0)} \cdot x = 0$	$\mathbf{P}, \mathbf{J}_{mn}$
Point form	$x \cdot x = \kappa^2, x^0 > 0$	$\mathbf{J}_{mn}$
Line form	$(x^0)^2 - (\mathbf{x}^\perp)^2 = \kappa^2,$ $x^0 > 0$	$P_3, \mathbf{J}_{12}$
Extended front form	$x^+ x^- = \kappa^2, x^0 > 0$	$\mathbf{P}_\perp, \mathbf{J}_{12}$

Table 3.4: Relativistic forms under the aspect of usefulness. The parameter κ is a real number.

under the original symmetry group. It is easy to realise that the elements of the original symmetry group which cease to be elements of a time-evolved quantisation surface do not commute with the time evolution operator of a given relativistic form. Hence all forms, except for the IF, single out an arbitrary instant of time. Another implication is this: in the IF, symmetries are stable. The "time" evolution operator does not transform an irreducible representation of the stability group into a different irreducible representation. This does not hold true in any other form. Temporal evolution of irreducible representations of a time-dependent stability group are mixed by the evolution operator. These irreducible representations should, of course, become irreducible representations of the stability group of the *evolved* quantisation surface. The relation between the original representation and the evolved representation is, however, dynamical, i.e. it depends on the interaction. A realistic comparison of different relativistic forms based on the stability group proper is given in table **Tab. 3.4**.

Chapter 4

The Massless Schwinger Model

Introduction

The SCHWINGER MODEL [81, 82] is quantum electrodynamics in two space-time dimensions with massless fermions. This model can be solved *analytically* by BOSONISATION [82]: the Schwinger model with coupling constant g is equivalent to a model of free bosons with mass $M_B = \frac{g}{\sqrt{\pi}}$. The original solution to the Schwinger model was given by J. Schwinger [82] using the action formalism; in this chapter we shall be using a solution in the Hamiltonian formalism given by N.S. Manton [83]. We are interested in the numerical treatment of the MASSIVE SCHWINGER MODEL, (i.e. QED(1+1)) with massive fermions ($m = 0$) because we shall *numerically* approximate the solutions to the MASSIVE SCHWINGER MODEL in Chapter 5. This numerical solution will be based on various assumptions and approximations that require justification: In particular, we shall assume that (1) left-movers may be neglected in small volumes, i.e. the number of virtual fermions moving contrary to the motion of physical boson states is negligible as long as the lattice size $2L$ is sufficiently small, (2) axial gauge—which is unphysical in a large volume—preserves the masses of the physical bosons as long as long

as these move sufficiently fast, (3) the vacuum has a simple, almost trivial structure on a small lattice, (4) left-movers are negligible in fast-moving states on a small lattice.

While the good description of this model which we shall obtain in $\langle 5 \rangle$, can already be regarded as an *a posteriori* justification of the approximations made, it is preferable, nonetheless, to seek a more thorough understanding of why these approximations preserve the physical character of QED(1 + 1). Thence, the exact solution to the Schwinger model in the *Hamiltonian* framework [83] constitutes an ideal laboratory to study the impact of various approximations in more detail.

Furthermore, we would like to investigate to what extent physics is affected if

(a) a naïve cut-off regularisation is employed instead of a proper gauge-invariant regularisation or

(b) to illustrate that in the infinite volume limit, the usual distinction of valence- and sea partons has to be supplemented by VACUUM partons which are related to the left-movers. These vacuum-partons do not contribute to the structure functions in the rest-frame.

(c) to calculate the proper distribution functions showing that they *never* coincide with the raw distribution functions on a large lattice, not even in the infinite momentum frame

(d) to demonstrate that non-triviality of the vacuum in general and of the chiral condensate in particular is inseparably linked to the presence of left-movers.

We argue that topological effects contribute primarily to the rest-frame sector $\mathbf{P} = 0$ where the De Broglie wave-length $\lambda = \frac{2\pi}{|\mathbf{P}|} = \infty$ is larger than any finite lattice size $2L$. Some introductory comments are in order.

1. We tried to keep our notation as close as possible to both the notation used for DIS as well as to Manton's notation. However, both notations do not coincide and hence some compromise had to be made. For example, we shall refer to particles with positive chirality as right-handed which corresponds to the usual definition of this word whereas Manton refers to negative chirality particles as right-handed. Left-handed fermions in [83] seem to move right and right-handed fermions seem to move left, contrary to the usual convention. The reason for the different terminologies is a subtle one: Manton implicitly abridges $x = x^1$ and $p = p_x = p_1 = -p^1$ as the reader may easily verify (We are using the 3-direction instead of the 1-direction and therefore Manton's p corresponds to $-p^3$ in our notation). This may be a good idea from the point of view of Chapter ⟨3⟩ as $\hat{P}_3$ rather than $\hat{P}^3$ is a generator of translations. Such a convention implies, however, that the state $a_{1,r}^\dagger |0\rangle$ has momentum $k^3 = -k_3 = -r$ rather than $+r$ ($r \in \mathbb{N}$) since the argument of Manton's operators are assumed to be *co*-variant rather than *contra*-variant operators. A state with $k_3 = r > 0$ moves LEFT rather than RIGHT, i.e. it moves against the x^3 axis.
2. It cannot be the scope of this thesis to repeat the article Ref. [83] in every detail; herein, we shall take most results for granted. The reader will **NOT** be able to understand this section unless he has studied Ref. [83].
3. A finite periodic box (circle) with length $2L = 2\pi$ is used as in Ref. [83]. It will be shown that without this volume-regularisation, the mean number of fermions in physical states—even in the vacuum—would be infinite.

4. The raw distribution functions we calculate can not be related to the structure functions since F_1 does not exist in two dimensions and the remaining longitudinal structure-function F_L vanishes due to the Callan-Gross relation which is not spoiled by perturbative corrections because quarks cannot radiate gluons with non-zero momentum.
5. In one space dimension, the symbols $\mathbf{k}$ and k^3 may be used interchangeably.

1 The Exact Solution to the Schwinger Model: A Brief Review

1.1 Terminology and Notation

We start with a brief review of the solution to the Schwinger model given by N. S. Manton [83].

The Lagrangian of the Schwinger model in standard notation is

$$\mathcal{L} = \bar{\psi}(i \not{\partial} - g \not{A})\psi - \frac{1}{4}F^{\mu\nu}F_{\mu\nu} \quad (4.1)$$

with $\psi = (\psi_R, \psi_L)$ the fermionic field and $F^{\mu\nu}$ the field strength tensor. We require the fermionic fields to be $2L$ -periodic, i.e. the Schwinger model is defined on a circle with circumference $2L$.

In order to be compatible with the usual three-dimensional notation (where the three-direction is the longitudinal direction) we use the co-ordinates $x = (x^0, x^3)$ rather than $x = (x^0, x^1)$. The metric g is the Minkowsky metric with signature $(+1, -1)$. Absorbing the coupling g into the connection (vector potential) A^μ

$$A \stackrel{def}{=} gA \quad (4.2)$$

and expressing the field strength $F^{\mu\nu}$ in terms of $\mathbf{A}$ yields Manton's form

$$\mathcal{L} = \bar{\psi}(i \not{\partial} - \not{\mathbf{A}})\psi + \frac{1}{2g^2}(\partial_0 \mathbf{A}_3 - \partial_3 \mathbf{A}_0)^2 \quad (4.3)$$

of the Lagrangian except that he uses the co-ordinates (x^0, x^1) instead of (x^0, x^3) . Please note that the field $\mathbf{A}$ which includes the coupling is written in *sans serif* style to facilitate the distinction. Since we are working in two dimensions, the generators γ^μ of the spinor group $\text{SPIN}(2) \equiv \text{SU}(2)$ are represented by the familiar σ matrices. Here, the chiral representation

$$\gamma^0 = +\sigma^1, \quad \gamma^3 = -i\sigma^2, \quad (4.4)$$

$$\gamma^5 = \sigma^3 = \gamma^0 \gamma^3 \quad (4.5)$$

from Ref. [83] will be used. The Lagrangian is invariant under the GAUGE TRANSFORMATION $\iota(x)$

$$\psi(x) \xrightarrow{\iota(x)} e^{-i\iota(x)} \psi(x) \quad (4.6)$$

$$\mathbf{A}_\mu(x) \xrightarrow{\iota(x)} \mathbf{A}_\mu(x) + \partial_\mu \iota(x) \quad . \quad (4.7)$$

The gauge parameter $\iota(x)$ is a function subjected to the constraint

$$\frac{1}{2\pi} [\iota(x^0, x^3 + L) - \iota(x^0, x^3 - L)] = n_W \in \mathbb{Z} \quad (4.8)$$

but otherwise arbitrary. This constraint has to be imposed in order to protect the periodicity of the fermionic field $\psi(x)$. Cf. Eq. (4.6). The number $n_W \in \mathbb{Z}$ is called WINDING NUMBER. Gauge transformations with non-vanishing winding number —i.e. non-periodic gauge transformations $\iota(x)$ — are called TOPOLOGICALLY NON-TRIVIAL GAUGE TRANSFORMATIONS or —shorter— LARGE GAUGE TRANSFORMATIONS. The constraint (4.8) prevents the ZERO-MODE OF THE GAUGE FIELD $\mathbf{A}_3(x)$

$$\mathbf{A}_3 \stackrel{\text{def}}{=} \int_{-L}^L \frac{dx^3}{2L} \mathbf{A}_3(x) \quad (4.9)$$

from being gauged away. Following Manton's minor abuse of notation, the symbol $\mathbf{A}_3$ now stands simultaneously for the gauge field *and* its zero mode. The zero-mode $\mathbf{A}_3$ is reduced to the domain¹ $0 \leq \mathbf{A}_3 < \pi/L$

$$\mathbf{A}_3 \xrightarrow{\iota(x)} \mathbf{A}_3 + \int_{-L}^L \frac{dx^3}{2L} \partial_3 \iota = \mathbf{A}_3 + n_W \pi/L \quad (4.10)$$

by a large gauge transformation. This reflects the fact that the GLOBAL WILSON LOOP

$$\exp \left[i \int_{-L}^L dx^3 \mathbf{A}_3(x) \right] = \exp \left[i \int_{-L}^L dx^3 \mathbf{A}_3(x) + 2\pi i \mathbb{Z} \right] \quad (4.11)$$

is the only gauge degree of freedom on a circle. The requirement that the global Wilson loop be invariant under gauge transformations would have been an alternative way of deducing the constraint (4.8). Axial gauge $\mathbf{A}_3 = 0$ would remove the zero-mode. Therefore, axial gauge is incompatible with periodic BCs. COULOMB GAUGE, in contrast,

$$\partial_3 \mathbf{A}_3(x) = 0 \quad (4.12)$$

is weaker than axial gauge. While leaving large gauge transformations intact, Coulomb gauge completely eliminates every *local* gauge transformation generated by Gauss law², i.e. it eliminates the gauge transformations with trivial winding number $n_W = 0$. Thus GAUSS' LAW

$$-\partial_3^2 \mathbf{A}_0 = g^2 \psi^\dagger \psi \quad (4.13)$$

¹The so-called FUNDAMENTAL MODULAR DOMAIN.

²Gauss' law is a FIRST CLASS CONSTRAINT in the terminology of the Dirac-Bergmann quantisation procedure [73, 74] and as such it may be considered the generator of *local* gauge transformations.

can be solved for A_0 such that the (un-regularised) HAMILTONIAN

$$\begin{aligned} H_{\text{raw}} &= \int_{-L}^L dx^3 \psi^\dagger (i\partial^3 - A^3) \gamma^0 \gamma^3 \psi + \frac{g^2}{2} \int_{-L}^L dx^3 (\psi^\dagger \psi) \frac{1}{-\partial_3^2} (\psi^\dagger \psi) + \frac{L}{g^2} E^2 \\ &= \int_{-L}^L dx^3 \psi^\dagger i\partial^3 \gamma^5 \psi + A_3 \mathcal{Q}_A + \frac{g^2}{2} \int_{-L}^L dx^3 (\psi^\dagger \psi) \frac{1}{-\partial_3^2} (\psi^\dagger \psi) + \frac{L}{g^2} E^2 \end{aligned} \quad (4.14)$$

contains only one dynamical boson field degree of freedom: the gauge field zero mode A_3 . The electric field $E = F^{03} = gF^{03}$

plays the rôle of the canonical momentum

$$[E, A_3] = -i \frac{g^2}{2L} \quad (4.15)$$

conjugate to the zero mode A_3 of the gauge field.

1.2 Operators in Momentum Space

Particles are most conveniently described in momentum space. There are several equivalent ways of expanding the fermionic field $\psi(x)$ in terms of creation and annihilation operators. Usually, one would choose an expansion in terms of particles and anti-particles with positive kinetic energies. It turns out, however, that the description of the Schwinger mode is drastically simplified in the PARTICLE-HOLE REPRESENTATION [83–86] wherein anti-particles are realised as holes in the Dirac sea. In this representation, the axial anomaly has an intuitive interpretation as the effect of a fluctuating Fermi level. We therefore expand the fermionic field in terms of particle annihilation operators

$$\psi(0, x^3) = \sum_{k^3} \frac{1}{\sqrt{2\pi}} (w_R a_{Rk^3} e^{ik^3 \cdot x^3} + w_L a_{Lk^3} e^{+ik^3 \cdot x^3}) \quad ; \quad (4.16)$$

with CHIRAL SPINORS

$$w_R = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad w_L = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (4.17)$$

in a way *similar*³ to Ref. [83]. The momenta k^3 assume multiple values of Δk , i.e. $k^3/\Delta k \in \mathbb{Z}$. The creation operator $a_{R,\mathbf{k}}^\dagger$ creates a *right*-handed particle with momentum $\mathbf{k}$ and $a_{L,\mathbf{k}}^\dagger$ creates a *left*-handed particle. The creation operators obey the usual anti-commutation relations

$$\{a_{\alpha,k^3}, a_{\beta,k^{3'}}^\dagger\} = \delta_{\alpha,\beta} \delta_{k^3,k^{3'}} \quad (4.18)$$

where $\alpha, \beta = R, L$.

1.3 The Hamiltonian in Manton's basis

Now we are able to express the Hamiltonian H in terms of the annihilation and creation operators defined above

$$\begin{aligned} H_{\text{raw}} = & \sum_{p^3} a_{R,p^3}^\dagger a_{R,p^3} (p^3 + A_3) + \sum_{p^3} a_{L,p^3}^\dagger a_{L,p^3} (-p^3 - A_3) \\ & + \frac{g^2}{4L} \sum_{p^3 \neq 0} \tilde{j}_{p^3}^0 \frac{1}{(p^3)^2} \tilde{j}_{-p^3}^0 + \frac{L}{g^2} E^2 \quad . \end{aligned} \quad (4.19)$$

This Hamiltonian is *formally* equivalent to the expression given in [83]. Note, however, the subtle yet important difference that we are using contra-variant momenta p^3 in lieu of co-variant momenta. The symbol $\tilde{j}^\mu$ stands for the spatial Fourier transform

$$\tilde{j}^0(P^3) \stackrel{\text{def}}{=} \int dx^3 e^{iP^3 x^3} j^0(x) = \rho_R(P^3) + \rho_L(P^3) \quad (4.20)$$

of the VECTOR CURRENT $j^\mu \stackrel{\text{def}}{=} \bar{\psi} \gamma^\mu \psi$ where

$$\rho_\alpha(\mathbf{P}) \stackrel{\text{def}}{=} \int dx^3 e^{ik^3 x^3} \psi_\alpha^\dagger \psi_\alpha = \sum a_{\alpha,\mathbf{P}+\mathbf{k}}^\dagger a_{\alpha,\mathbf{k}} \quad (4.21)$$

³ Please note that $a_{R,n}^\dagger$ creates a *right*-handed particle with momentum $k^3 = n$ whereas in Ref. [83], the similar operator $a_{1,n}^\dagger$, creates a *left*-handed particle with momentum $k^3 = -k_3 = -n$.

with $\alpha = R, L$. Classically, both the VECTOR CURRENT

$$j^\mu(x) \stackrel{def}{=} \bar{\psi}(x) \gamma^\mu \psi(x) \quad (4.22)$$

and the AXIAL CURRENT

$$j_A^\mu(x) \stackrel{def}{=} \bar{\psi}(x) \gamma^\mu \gamma^5 \psi(x) \quad (4.23)$$

are conserved quantities because the Lagrangian is invariant under the vectorial $U(1)$ transformation

$$\psi(x) \longmapsto e^{-i\mathbb{1}\theta_V} \psi(x) \quad (4.24)$$

and the axial $U(1)$ transformation

$$\psi(x) \longmapsto e^{-i\gamma^5 \theta_A} \psi(x) \quad . \quad (4.25)$$

In the quantised theory, however, the Hamiltonian H , the VECTOR CHARGE $\mathcal{Q}$ and the AXIAL CHARGE $\mathcal{Q}_A$

$$\mathcal{Q} \stackrel{def}{=} \int_{-L}^L dx^3 j^0 = \mathcal{Q}_R + \mathcal{Q}_L \quad , \quad \mathcal{Q}_A \stackrel{def}{=} \int_{-L}^L dx^3 j_A^0 = \mathcal{Q}_R - \mathcal{Q}_L \quad (4.26)$$

with $\mathcal{Q}_\alpha = \rho_\alpha(0)$ have to be RENORMALISED by subtracting their respective vacuum expectation values. The vacuum energy is the only part of the Hamiltonian that contains ultra-violet divergences. The renormalisation constants are unity. This property is related to the fact that the Schwinger-model is SUPER-RENORMALISABLE (i.e. the coupling constant g has the dimension of a mass).

Manton performs the renormalisation in the limit $g \rightarrow 0$ at first. The divergent vacuum energy of H_{raw} is contained in the first part of H_{raw}

$$T_{\text{raw}} = \sum_{p_3} a_{R,p^3}^\dagger a_{R,p^3} (p^3 + \mathbf{A}_3) + \sum_{p_3} a_{L,p^3}^\dagger a_{L,p^3} (-p^3 - \mathbf{A}_3) \quad . \quad (4.27)$$

Eigenvectors of T_{raw} can be constructed as follows: Let us define the A-VACUUM $|\Omega_a : \mathbf{A}_3\rangle$ as an eigenstate $|\Omega_a : \mathbf{A}_3\rangle$ of the vector potential operator

$$\hat{\mathbf{A}}_3 |\Omega_a : \mathbf{A}_3\rangle = \mathbf{A}_3 |\Omega_a : \mathbf{A}_3\rangle \quad (4.28)$$

which is annihilated by the operators a_{α,k^3}

$$\forall_{\alpha;k^3} a_{\alpha,k^3} |\Omega_a : \mathbf{A}_3\rangle = 0 \quad . \quad (4.29)$$

This state is an eigenstate to T_{raw} with eigenvalue zero. For $\mathbf{A}_3 = 0$ we may construct the state with lowest (naïve) energy (i.e. the vacuum of T_{raw}) by filling the a-vacuum with left-moving right-handed particles and with right-moving right-handed particles as illustrated in **Fig. 4.1**. Choosing $\mathbf{A}_3 \neq 0$, the so-constructed state is no longer the state with lowest energy. Filling up the a -vacuum around a different DISCRETE FERMİ LEVEL $\mathcal{M}$ with right-handed fermions $a_{R,k^3}; k^3 \leq \mathcal{M}$ and left-handed fermions $a_{L,k^3}; k^3 > \mathcal{M}$ we obtain the AXIAL VACUUM

$$|\Omega : \mathbf{A}_3, \mathcal{M}\rangle \stackrel{def}{=} \prod_{k^3 > \mathcal{M}} a_{R,\mathcal{M}+1-\mathbf{k}}^\dagger a_{L,\mathbf{k}}^\dagger |\Omega_a : \mathbf{A}_3\rangle \quad (4.30)$$

with NAÏVE AXIAL CHARGE : $\mathcal{Q}_A := 2\mathcal{M}$. This construction is illustrated in **Fig. 4.1**. Please note that the vacuum symbols cannot be found in Manton's article. We have introduced them for later convenience.

Now, there is a serious problem with this argument, of course. The energies of the states $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ are, in fact, infinite. So is their charge $\mathcal{Q}$. Little does it make sense, consequently, to speak of "the state with lowest energy". The axial charge, finally, is finite but ill-defined. The subtraction of infinite quantities has to be rendered well-defined by means of a controlled regularisation procedure. The difference of an infinite number of left-movers and an infinite number of right-movers is then defined via the CONTINUUM

LIMIT, i.e. the limit in which the regularisation is removed. It turns out that different regularisation procedures yield different results in the continuum limit. A cut-off regularised Hamiltonian, for instance, does not become gauge-invariant in the continuum limit whereas the vectorial charge and the axial charge are both conserved for every cut-off parameter Λ . Manton employs the gauge-invariant HEAT-KERNEL REGULARISATION in order to regularise the vacuum energy of $\mathcal{Q}$, $\mathcal{Q}_A$ and H_{raw} . Basically, this regularisation can be seen as the gauge-invariant version of a SMOOTH CUT-OFF Λ_h which consists in damping the number of right-handed fermions $a_{R,\mathbf{k}}^\dagger a_{R\mathbf{k}}$ and




Figure 4.1: **Symbolic representation of the vacuum $|\Omega : A_3, \mathcal{M}\rangle$ with $\mathcal{M} = 0$ and $A_3 = 0$.** The circles represent right-handed fermions filling the Fermi sea from below. The rounded squares represent left-handed fermions filling the Fermi sea from above.

left-handed fermions $a_{L,\mathbf{k}}^\dagger a_{L,\mathbf{k}}$ with the factors $\exp(\frac{k^3 + \mathbf{A}_3}{\Lambda_h})$ and $\exp(-\frac{k^3 + \mathbf{A}_3}{\Lambda_h})$ respectively⁴. This procedure renders infinite sums such as $\sum_{k^3 \leq 0} 1$ finite. Manton finds that the divergent vacuum expectation values

$$\langle \Omega : \mathbf{A}_3, \mathcal{M} | \mathcal{Q}_R | \Omega : \mathbf{A}_3, \mathcal{M} \rangle = c_\infty + (\mathcal{M} + \mathbf{A}_3 + \frac{1}{2}) \quad (4.31)$$

$$\langle \Omega : \mathbf{A}_3, \mathcal{M} | \mathcal{Q}_L | \Omega : \mathbf{A}_3, \mathcal{M} \rangle = c_\infty - (\mathcal{M} + \mathbf{A}_3 + \frac{1}{2}) \quad (4.32)$$

$$\langle \Omega : \mathbf{A}_3, \mathcal{M} | T_{\text{raw}} | \Omega : \mathbf{A}_3, \mathcal{M} \rangle = c_\infty^H + (\mathcal{M} + \mathbf{A}_3 + \frac{1}{2})^2 \quad (4.33)$$

may be decomposed in a finite, field-dependent part and field-independent constants $c_\infty \approx \Lambda_h + \mathcal{O}(1/\Lambda_h)$, $c_\infty^H \approx -2\Lambda_h^2 + \mathcal{O}(1/\Lambda_h)$ which diverge in the continuum limit $\Lambda_h \rightarrow \infty$. Subtracting these constants from the regularised operators $\mathcal{Q}_R^{\text{reg}} \stackrel{\text{def}}{=} \mathcal{Q}_R - c_\infty$, $\mathcal{Q}_L^{\text{reg}} \stackrel{\text{def}}{=} \mathcal{Q}_L - c_\infty$ and $H_{\text{reg}} \stackrel{\text{def}}{=} H_{\text{raw}} - c_\infty^H$ yields the renormalised operators. It is crucial that gauge-invariant regularisation produce expressions which depend on the gauge-field. The finite gauge-dependent part of the *vacuum energy* acts now as an additional interaction term of the Hamiltonian. The finite gauge-dependent part of the *axial charge* is not conserved and will be responsible for the anomaly. The regularised charge $\mathcal{Q}_{\text{reg}}$ is zero by construction: a non-vanishing vectorial charge would be incompatible with Gauss' law (4.13). The regularised axial charge may be written as $\mathcal{Q}_A^{\text{reg}} = 2\hat{\mathcal{M}} + 2\mathbf{A} + 1$, where $\hat{\mathcal{M}}$ is the discrete Fermi level. As a result, the quantity $\mathcal{Q}_A^{\text{reg}}/2$ may now be regarded as a continuous Fermi label. This means that we end up with the —somewhat paradoxical— result that the difference between the (infinite) number of right-movers and the infinite number of left-movers is not an integer. This result becomes less paradoxical, of course, if one realises that heat-kernel regularisation consists in giving fermions a fractional, $\mathbf{A}_3$ -dependent particle number. Hence

⁴ A conventional/gauge-variant smooth cut-off would employ the factors $\exp(\pm \frac{k^3}{\Lambda_h})$ instead.

we should not be overly surprised that the axial charge remains fractional and A_3 -dependent in the continuum limit. The same comment applies to alternative gauge-invariant regularisation techniques.

The heat-kernel regularised charges are invariant under LARGE GAUGE TRANSFORMATIONS U_G

$$U_G |\Omega : A_3, \mathcal{M}\rangle = |\Omega : A_3 - 1, \mathcal{M} + 1\rangle \quad (4.34)$$

which simultaneously change $\mathcal{M}$ and A_3 . Gauge-invariant states are irreducible representations of the group of large gauge transformations, generated by U_G with group characters $e^{-i\theta}$. They are called θ -STATES.

Naïve renormalisation based on cut-off regularisation partially destroys gauge invariance. Conversely, if the renormalisation procedure chosen is gauge-invariant, then the *renormalised* axial charge ceases to be conserved. This phenomenon is called AXIAL ANOMALY. The technical details involved in order to describe the axial anomaly vary remarkably, depending on the way quantisation is performed: (1) In PERTURBATION THEORY, the anomaly arises due to ambiguities in divergent loop integrals of a certain class of Feynman diagrams, the easiest of which is the so-called TRIANGLE DIAGRAM. For details, the reader is referred to Ref. [87,88]. (2) In PATH-INTEGRAL QUANTISATION [89], the renormalisation ambiguity can be traced back to the path integral measure which has to be regularised. All of these manifestations have in common that the anomaly stems from the necessity of choosing amongst different regularisation procedures one of which is gauge invariant. If one uses an infinite volume or fixed BCs, then the anomaly manifests itself in yet another way. In this case axial gauge may be chosen and A_3 disappears. The θ angle is then due to the fact that adding an *unperiodic* term θx^3 to a solution A_0 to Gauss' law (4.13) transforms it into another solution [90,91]. This term may then be interpreted as a constant background field [90,91].

We note in passing that one may regularise the axial charge such that it is conserved and the vector charge is not. In the Hamiltonian approach, however, the non-conservation of (vector) charge is incompatible with Gauss' law.

In order to diagonalise the heat-kernel regularised Hamiltonian, Manton expresses the Hamiltonian in terms of the bosonic fields Π and Φ

$$\Pi(\mathbf{p}) = +\frac{\rho_R(\mathbf{p}) - \rho_L(\mathbf{p})}{\sqrt{2}} = +\frac{1}{\sqrt{2}}\tilde{j}_{\mathbf{p}}^3 \quad (4.35)$$

$$\Phi(\mathbf{p}) = -\frac{\rho_R(\mathbf{p}) + \rho_L(\mathbf{p})}{\sqrt{2}ip^3} = -\frac{\tilde{j}_{p^3}^0}{\sqrt{2}ip^3} \quad (4.36)$$

$$\Pi(0) = \sqrt{2}(\tilde{A} + \frac{1}{2}) \quad (4.37)$$

$$\Phi(0) = \frac{1}{g^2\sqrt{2}}E \quad (4.38)$$

$$[\Pi(-p^3), \Phi(p^{3'})] |\text{phy}\rangle = [\Pi_{p^3}^\dagger, \Phi(p^{3'})] |\text{phy}\rangle = -i\delta(p, p') |\text{phy}\rangle \quad . \quad (4.39)$$

The bosonic commutation relations **(4.39)** hold on a PHYSICAL SUBSPACE of the total Fock space only. Physical states $|\text{phy}\rangle$ are states with finite regularised energy, i.e. $\langle \text{phy} | H_{\text{reg}} | \text{phy} \rangle < \infty$. With these operators, the Hamiltonian assumes a form

$$H_{\text{reg}} = \frac{1}{2} \sum_{p^3} (\Pi^\dagger(p^3)\Pi(p^3) + \Omega_{p^3}^2 \Phi^\dagger(p^3)\Phi(p^3)) \quad (4.40)$$

which is manifestly equivalent to a free theory of bosons with mass $M_B \stackrel{\text{def}}{=} g/\sqrt{\pi}$ and energy is

$$\Omega(p^3) \stackrel{\text{def}}{=} \sqrt{(p^3)^2 + \frac{g^2}{\pi}} \quad . \quad (4.41)$$

These bosons are called SCHWINGER BOSONS. Let us finally look, however briefly, at the nature of the axial anomaly exhibited by the Schwinger model.

The axial symmetry of the Lagrangian is not realised in the mass spectrum. This might easily lead us to the conclusion that the heat kernel regularisation breaks the axial symmetry in the sense that $[H_{\text{reg}}, : \mathcal{Q}_A :] = 0$ no longer holds. This is wrong, $[H_{\text{reg}}, : \mathcal{Q}_A :] = 0$ *does* hold. Axial symmetry and gauge symmetry are incompatible on the quantum mechanical level since both symmetries are now merged into a larger, non-commutative group: $[U_G, : \mathcal{Q}_A :] \neq 0$. In other words, U_G and $: \mathcal{Q}_A :$ form a non-commutative algebra, i.e. they cannot be measured simultaneously. The axial symmetry is not realised in the spectrum because states with different axial charges are related via U_G : therefore, states with different axial charges must have the same energy.

AXIAL GAUGE breaks the symmetry generated by U_G and, consequently, destroys the degeneracy of the mass spectrum: spurious states appear.

2 Translation into the Particle-Antiparticle Picture

In the framework of the parton model, fermion fields are expressed

$$\psi(0, x^3) = \sum_{k^3} \frac{1}{\sqrt{2\pi}} (\mathbf{u}(k^3) b_{k^3} e^{+ik^3 \cdot x^3} + \mathbf{v}(k^3) d_{k^3}^\dagger e^{-ik^3 \cdot x^3}) \quad (4.42)$$

in terms of particle creation operators and anti-particle creation operators rather than in terms of particles and holes. The usual spinors (for massless fermions)

$$\mathbf{u}(k^3) = \begin{cases} +w_R & \text{if } k^3 > 0 \\ +w_L & \text{if } k^3 \leq 0 \end{cases} \quad (4.43)$$

$$\mathbf{v}(k^3) = \begin{cases} +w_R & \text{if } k^3 > 0 \\ -w_L & \text{if } k^3 \leq 0 \end{cases} \quad (4.44)$$

are used. We have absorbed the term $1/\sqrt{\omega(k^3)}$ usually appearing in the expansion into the spinors $\mathbf{u}$ and $\mathbf{v}$. The choice for $k^3 = 0$ is arbitrary, reflecting the fact that there are two degenerate vacua of the free, massless fermionic theory. Naturally, the operators defined above fulfill the standard anti-commutation relations

$$\{b_{k^3}, b_{k^3'}^\dagger\} = \delta_{k^3, k^3'} \quad \{d_{k^3}, d_{k^3'}^\dagger\} = \delta_{k^3, k^3'} \quad . \quad (4.45)$$

In what follows we shall set $\Delta k = 1$ (i.e. $L = \pi$). In order to transform the exact solution to the Schwinger model into the partonic picture, we relate the different creation and annihilation operators. This can be done by a simple comparison of these two expansions

$$\begin{pmatrix} a_{R, k^3} \\ a_{L, k^3} \end{pmatrix} = \begin{cases} \begin{pmatrix} b_{k^3} \\ +d_{-k^3}^\dagger \end{pmatrix} & \text{if } k^3 > 0 \\ \begin{pmatrix} -d_{-k^3}^\dagger \\ b_{k^3} \end{pmatrix} & \text{if } k^3 \leq 0 \end{cases} \quad (4.46)$$

We call the vacuum $|\Omega : \mathbf{A}_3, 0\rangle$ which is annihilated by $b_{\mathbf{k}}$ and $d_{\mathbf{k}}$

$$\forall_{k^3} b_{k^3} |\Omega : \mathbf{A}_3, 0\rangle = 0 \quad \forall_{k^3} d_{k^3} |\Omega : \mathbf{A}_3, 0\rangle = 0 \quad (4.47)$$

the bd-VACUUM. This vacuum is identical to the vacuum $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ constructed in the last subsection if we set $\mathcal{M} = 0$. In particular, $|0\rangle \stackrel{def}{=} |\Omega : \mathbf{A}_3 = 0, 0\rangle$ is the PERTURBATIVE VACUUM of the theory in axial gauge $\mathbf{A}_3 = 0$.

A change of the discrete Fermi level $\mathcal{M}$ has to be seen as a creation or annihilation of particles if the bd-basis is used rather than the a-basis. For example, $|\Omega : \mathbf{A}_3, -1\rangle$ is obtained from $|\Omega : \mathbf{A}_3, 0\rangle$ by lowering the Fermi-level with the operator $a_{R,0} a_{L,0}^\dagger$ and therefore

$$|\Omega : \mathbf{A}_3, -1\rangle = a_{L,0}^\dagger a_{R,0} |\Omega : \mathbf{A}_3, 0\rangle = b_0^\dagger d_0^\dagger |\Omega : \mathbf{A}_3, 0\rangle \quad . \quad (4.48)$$

Similarly, the vacuum for $\mathcal{M} = -2$ is obtained as

$$|\Omega : \mathbf{A}_3, -2\rangle = a_{L,0}^\dagger a_{R,0} |\Omega : \mathbf{A}_3, -1\rangle = b_{-1}^\dagger d_{+1}^\dagger b_0^\dagger d_0^\dagger |\Omega : \mathbf{A}_3, 0\rangle \quad (4.49)$$

and so on.

Now we have to regularise the Hamiltonian. Heat-kernel regularisation works only on an infinite lattice. Numerical methods, however, can handle a finite number of (effective) degrees of freedom only. Therefore, we shall not repeat this procedure in spite of its elegance. Instead, we suggest a *heuristic* procedure which will turn out to be equivalent to heat-kernel regularisation. We start with a naïve cut-off regularisation and regularise the divergent vacuum energy by normal-ordering. As this destroys the invariance of the Hamiltonian under LARGE GAUGE TRANSFORMATIONS (i.e. the topologically non-trivial gauge transformations that remain in spite of Coulomb gauge [83]) we have to restore gauge invariance *by hand* in order to obtain physically meaningful results. We shall show that gauge-invariance can be repaired simply by adding a term to the Lagrangian which disappears in the infinite volume limit. It will turn out that the addition of this term exactly reproduces the same Hamiltonian as heat-kernel regularisation if the cut-off is removed. The un-regularised Hamiltonian in the bd basis reads

$$H_{\text{raw}} = T + \mathbf{A}_3 \mathcal{Q}_A - \Lambda(\Lambda + 1) + \frac{L}{g^2} \mathbf{E}^2 + \frac{g^2}{4L} \sum_{p^3 \neq 0} \tilde{j}_{p^3}^0 \frac{1}{(p^3)^2} \tilde{j}_{-p^3}^0 \quad (4.50)$$

where T is the normal-ordered KINETIC ENERGY

$$T = \sum_{p^3} (b_{p^3}^\dagger b_{p^3} + d_{p^3}^\dagger d_{p^3}) |p^3| \quad (4.51)$$

and $\Lambda(\Lambda + 1)$ is the constant which arises when the kinetic energy is normal ordered. This constant has to be regularised since it contains the short-distance singularity $\Lambda(\Lambda + 1)$. Normal ordering the Hamiltonian with respect

to the bd operators corresponds to removing this term. Subtracting the kinetic energy of the state $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$

$$\hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) \quad (4.52)$$

from the kinetic energy T , we define the MODIFIED KINETIC ENERGY

$$\mathfrak{T} \stackrel{def}{=} \sum_{p^3} (b_{p^3}^\dagger b_{p^3} + d_{p^3}^\dagger d_{p^3}) |p^3| - \hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) \quad (4.53)$$

which is insensitive to the discrete Fermi-level $\mathcal{M}$. While the states

$$b_{k^3}^\dagger d_{p^3}^\dagger |\Omega : \mathbf{A}_3, 0\rangle \text{ and } b_{k^3}^\dagger d_{p^3}^\dagger |\Omega : \mathbf{A}_3, \mathcal{M}\rangle \quad (4.54)$$

have different kinetic energies T , for instance, (i.e. $|k^3| + |p^3|$ resp. $|k^3| + |p^3| + \mathcal{M}(\mathcal{M} + 1)$) their modified energies $|k^3| + |p^3|$ coincide. The same holds for more complicated states. We have introduced $\mathfrak{T}$ because it is invariant under topologically non-trivial gauge transformations U_G which change the axial charge whereas the conventional kinetic energy is *not*. Both operators have the same eigenstates but their eigenvalues coincide solely in the zero axial charge $\mathcal{Q}_A = 0$ sector. The distinction between T and $\mathfrak{T}$ (not explicitly introduced in Ref. [83]) is very important. The bosonized kinetic energy in Ref. [83] corresponds in fact to the *modified* kinetic energy $\mathfrak{T}$ rather than the usual kinetic energy T . Now we insert the new quantity $\mathfrak{T}$ into the normal-ordered (with respect to bd) Hamiltonian

$$H_{\text{N.O.}} = \mathfrak{T} + \mathcal{M}(\mathcal{M} + 1) + 2\mathcal{M}\mathbf{A}_3 + \mathbf{A}_3 + \frac{L}{g^2} \mathbf{E}^2 + \frac{g^2}{4L} \sum_{p^3 \neq 0} \tilde{j}_{p^3}^0 \frac{1}{(p^3)^2} \tilde{j}_{-p^3}^0 \quad (4.55)$$

because we want to isolate the terms which are not invariant under LARGE GAUGE TRANSFORMATIONS U_G

$$\hat{\mathcal{M}} \rightarrow U_G \hat{\mathcal{M}} U_G^{-1} = \hat{\mathcal{M}} + 1 \quad (4.56)$$

$$\hat{\mathbf{A}}_3 \rightarrow U_G \hat{\mathbf{A}}_3 U_G^{-1} \hat{\mathbf{A}}_3 - 1. \quad (4.57)$$

After completion of the square

$$H_{\text{N.O.}} = \mathfrak{T} + (\hat{\mathcal{M}} + \mathbf{A}_3 + \frac{1}{2})^2 - \mathbf{A}_3^2 - \frac{1}{4} + \frac{L}{g^2} \mathbf{E}^2 + \frac{g^2}{4L} \sum_{p^3 \neq 0} \tilde{j}_{p^3}^0 \frac{1}{(p^3)^2} \tilde{j}_{-p^3}^0 \quad (4.58)$$

we see that the quadratic term $-\mathbf{A}_3^2$ is the only term which varies under large gauge transformations. Subtracting $\mathbf{A}_3^2$ from the Lagrangian or, equivalently, adding it to the Hamiltonian

$$H_{\text{reg}} = H_{\text{N.O.}} + \mathbf{A}_3^2 + \frac{1}{4} \quad , \quad (4.59)$$

the Hamiltonian becomes identical to the one obtained by HEAT-KERNEL REGULARISATION. **This means that the quantisation of the *gauge-variant* Lagrangian $\mathcal{L}' \stackrel{\text{def}}{=} \mathcal{L} - \frac{L}{\pi} \mathbf{A}_3^2$ with a naïve cut-off regularisation, is equivalent to quantisation of the *gauge-invariant* Lagrangian $\mathcal{L}$ with a heat-kernel regularised Hamiltonian (in the limit $\Lambda \rightarrow \infty$).** The addition of one single field variable $\mathbf{A}_3^2$ might seem small a change in the infinite volume limit where an infinite number of fields is present: the term $\frac{L}{\pi} (\mathbf{A}_3)^2$ (units restored) even vanishes in the infinite volume limit $L \rightarrow \infty$ because $\mathbf{A}_3$ can always be gauged to be smaller than $\triangle k = \frac{\pi}{L}$. Yet this step has a profound influence on the spectrum and the condensate (in the rest frame and only in the rest frame). It is remarkable that Hamiltonians which have the same classical infinite volume limits should have different quantum mechanical infinite volume limits. While the heuristic procedure proposed here is not necessary in the Schwinger model where the only divergence in the Hamiltonian is the vacuum energy, a similar procedure may turn out to be helpful for theories where the Hamiltonian itself diverges (e.g. QCD(3+1)). The Hamiltonian H_{reg} is invariant under *two*⁵ symmetries: AXIAL TRANSFORMATIONS generated by $:\mathcal{Q}_A:$ and large GAUGE-TRANSFORMATIONS U_G .

⁵ Note that the anomaly does *not* mean that the axial symmetry is broken: the point is

This distinguishes H_{reg} from a large class of Hamiltonians which differ by quantity that vanishes in the limit $l \rightarrow \infty$.

3 Solutions to the Schwinger Model and the Influence of Approximations

In order to study the influence of several approximations it is convenient to introduce some Hamiltonians constructed using these approximations: The HAMILTONIAN IN AXIAL GAUGE

$$H_{\text{axial}} = H_{\text{N.O.}} - \frac{L}{g^2} \mathbf{E}^2 - (2\hat{\mathcal{M}}_A + 1)\mathbf{A}_3 \quad (4.60)$$

would have been obtained through the choice of the *unphysical* (for large volumes, see [83]) axial gauge plus normal-ordering. This Hamiltonian is not invariant under large gauge transformations. If we subtract the term $\mathcal{M}(\mathcal{M} + 1)$, however, the resulting ARTIFICIAL HAMILTONIAN

$$H_{\text{arti}} = H_{\text{axial}} - \hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) = H_{\text{reg}} - \frac{L}{g^2} \mathbf{E}^2 - (\tilde{\mathbf{A}} + \frac{1}{2})^2 \quad (4.61)$$

$$= \mathfrak{T} + \frac{g^2}{4L} \sum_{p^3 \neq 0} \tilde{j}_{p^3}^0 \frac{1}{(p^3)^2} \tilde{j}_{-p^3}^0 \quad (4.62)$$

with $\tilde{\mathbf{A}} \stackrel{\text{def}}{=} \mathcal{M} + \mathbf{A}_3$ is again invariant under large gauge transformations as it differs from the heat-kernel Hamiltonian by a gauge-invariant quantity.

Now, we would like to find a complete solution to all of these Hamiltonians and interpret them in terms of quark- and gluon- degrees of freedom (in the literature, the fermions of the Schwinger model are often suggestively referred to as QUARKS, the gauge boson field as GLUONS. We follow this convention).

that the generator of the axial symmetry does not commute with the generator of the large gauge symmetry. As physical states have to be gauge invariant, the conserved quantity $: \mathcal{Q}_A :$ cannot be observed whereas the non-conserved quantity $\mathcal{Q}_A^{\text{reg}}$ can be: the axial symmetry is invisible.

In addition to Ref. [83] it is useful to introduce the operators

$$\begin{aligned}\mathcal{A}_{k^3}^\dagger &\stackrel{def}{=} \frac{1}{\sqrt{2}}(\sqrt{\Omega(k^3)}\Phi^\dagger(-k^3) - \frac{1}{\sqrt{\Omega(k^3)}}\Pi(k^3)i) \\ &= -i \frac{\rho_R(k^3)(\frac{\Omega(k^3)}{k^3} + 1) + \rho_L(k^3)(\frac{\Omega}{k^3} - 1)}{2\sqrt{\Omega}}\end{aligned}\quad (4.63)$$

$$\begin{aligned}\mathcal{A}_0 &= \frac{\sqrt{\Omega}}{\sqrt{2}}(\Phi(0) + \frac{1}{\Omega}\Pi^\dagger(0)i) \\ &= \frac{\sqrt{M_B}}{\sqrt{2}}(\frac{i}{\sqrt{2}}\frac{d}{d\tilde{\mathbf{A}}} + \frac{1}{M_B}\sqrt{2}(\tilde{\mathbf{A}} + \frac{1}{2}))i \\ &= -\frac{1}{\sqrt{M_B}}(\mathbf{E}/M_B - (\tilde{\mathbf{A}} + \frac{1}{2}i))\end{aligned}\quad (4.64)$$

which create and absorb Schwinger bosons. The artificial Hamiltonian

$$H_{\text{arti}} = \sum_{p^3 \neq 0} \mathcal{A}_{p^3}^\dagger \mathcal{A}_{p^3} \Omega(p^3) \quad (4.65)$$

written in this form is then equivalent to the heat-kernel regularised Hamiltonian except that H_{arti} lacks the pure gauge sector

$$H_{\text{gauge}} \stackrel{def}{=} H_{\text{reg}} - H_{\text{arti}} = \frac{L}{g^2}\mathbf{E}^2 + (\tilde{\mathbf{A}} + \frac{1}{2})^2 = \mathcal{A}_0^\dagger \mathcal{A}_0 \Omega(0) \quad (4.66)$$

describing the rest frame. Consequently, the axial Hamiltonian

$$H_{\text{axial}} = \sum_{p^3 \neq 0} \mathcal{A}_{p^3}^\dagger \mathcal{A}_{p^3} \Omega(p^3) + \hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) = H_{\text{arti}} + \hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) \quad (4.67)$$

is not Lorentz co-variant except for states with $\mathcal{M} = 0, -1$: all other states are SPURIOUS STATES whose masses depend strongly on $\mathbf{P}$ because their four-momenta

$$P_{\mathcal{M}} \stackrel{def}{=} \binom{\Omega(\mathbf{P})}{\mathbf{P}} + \binom{\mathcal{M}(\mathcal{M} + 1)}{0} \quad (4.68)$$

do not lie on the mass-shell, i.e. $P_{\mathcal{M}}^2 \neq M_B^2$. In summary, axial gauge destroys both Lorentz covariance and the axial anomaly.

It is interesting though that these illnesses of axial gauge are swept under the carpet in the (unphysical) limit $L \rightarrow 0$. The unphysical term $\mathcal{M}(\mathcal{M} + 1)\Delta k$ diverges if physical units are restored (i.e. $L \neq \pi$) because Δk diverges. Consequently, spurious states with $\mathcal{M}(\mathcal{M} + 1) \neq 0$ are removed from the low-energy mass spectrum. If only the mass-spectrum is concerned, the peculiar limit $L \rightarrow 0$ mimicks the anomaly. This phenomenon is a *fake-realisation* of the anomaly since two wrongs —i.e. zero lattice size and axial gauge— make a right and this is not acceptable.

4 The Schwinger Model on an Infinitesimal Lattice

In this section, we point out that the mass spectrum of the Schwinger model may be reproduced in the CONSTITUENT QUARK MODEL *without phenomenological input* if (1) the lattice size L is sufficiently small and (2) no θ states are constructed from axial states. We demonstrate that the vacuum of the Schwinger model is trivial and that the Schwinger bosons are pure two-fermion states if the limit of an infinitesimally small lattice size is performed. This holds except for the rest-frame where the Schwinger boson may be interpreted as a topological glue-ball. Remarkably, the limit $L \rightarrow 0$ does not affect the mass-spectrum. In the ensuing section, we shall then demonstrate that the large volume solutions to the Schwinger model may be obtained by a simple, unitary transformation. Herein, the ZERO VOLUME BASIS

$$\mathbb{A}_{P^3} \stackrel{def}{=} \begin{cases} \frac{i}{\sqrt{|P^3|}} \rho_R(-P^3) & \text{for } P^3 > 0 \\ -\frac{i}{\sqrt{|P^3|}} \rho_L(-P^3) & \text{for } P^3 < 0 \end{cases} \quad (4.69)$$

will be used. This basis appears in the limit

$$M_B L \rightarrow 0 \Leftrightarrow g/\Delta k \rightarrow 0 \quad (4.70)$$

which may be interpreted either as the zero-volume limit $2L \rightarrow 0$ (if the physical mass M_B is constant) or as the limit of vanishing Schwinger boson mass M_B (if the volume is kept constant).

$$\lim_{p^3/M_B \rightarrow \infty} \mathcal{A}_{p^3}^\dagger = -i \frac{\rho_R(p^3)}{2\sqrt{p^3}} = \mathbb{A}_{p^3} \quad (4.71)$$

where $\Omega(\mathbf{p}) \rightarrow |\mathbf{p}|$. The zero volume basis has the interesting property that these operators annihilate the perturbative vacuum for $\mathbf{P} \neq 0$. The same holds for every axial vacuum $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ —a fact that is immediately clear in Manton’s a -basis: the operators ρ_R cannot move fermions onto sites with negative momenta as they are already occupied. The same way, ρ_L is unable to shift fermions deeper into the occupied Fermi sea. This proves

Theorem 7 (Triviality of the Vacuum) *The axial vacua $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ and $|\Omega : 0, \mathcal{M}\rangle$ are eigenstates of the Hamiltonians H_{axial} and H_{arti} if $g \ll \Delta k$. The axial vacua $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ are vacua (i.e. lowest eigenstates) of H_{arti} . H_{axial} , in contrast, has only two vacua $|\Omega : 0, 0\rangle$ and $|\Omega : 0, -1\rangle$.*

The reason for why the axial Hamiltonian H_{axial} has only two vacua $|\Omega : 0, 0\rangle$ and $|\Omega : 0, -1\rangle$ is that the energies $\mathcal{M}(\mathcal{M} + 1)$ of the other states $|\Omega : 0, \mathcal{M}\rangle$ are larger than zero. Theorem (7) implies that the PERTURBATIVE VACUUM $|0\rangle \stackrel{def}{=} |\Omega : 0, 0\rangle$ is a vacuum of both Hamiltonians. In other words, both Hamiltonians have one TRIVIAL vacuum and the other vacua are almost trivial. We may construct θ vacua as projectively gauge-invariant linear combinations of the trivial vacua

$$|\Omega : \tilde{\mathbf{A}}; \theta\rangle = \sum_{\mathcal{M}} |\Omega : \tilde{\mathbf{A}} - \mathcal{M}, \mathcal{M}\rangle \frac{1}{\sqrt{2\pi}} e^{-i\theta\mathcal{M}} \quad . \quad (4.72)$$

These states remain eigenstates of H_{arti} but not of H_{axial} .

The states $|\Omega : \mathbf{A}_3, \mathcal{M}\rangle$ are not vacua of H_{reg} because they are not eigenvectors of the sector $M_B \mathcal{A}_0^\dagger \mathcal{A}_0$ which consists of gauge fields only. Using the eigenfunctions $\mathbf{H}_n(\mathbf{A}/M_B)$ of an harmonic oscillator in the spatial representation, however, we may build harmonic oscillator states

$$|\Omega_n : \mathcal{M}\rangle \stackrel{\text{def}}{=} \int d\mathbf{A} \mathbf{H}_n\left(\frac{\mathbf{A}}{M_B}\right) |\Omega : \mathbf{A}_3, \mathcal{M}\rangle \quad (4.73)$$

that are eigenstates of H_{reg} . These, in turn, may be super-imposed to θ vacua

$$\begin{aligned} |\Omega_n; \theta\rangle &= [\mathbb{A}_0^\dagger]^n |\Omega_0; \theta\rangle = \int d\tilde{\mathbf{A}} \mathbf{H}_n\left(\frac{\tilde{\mathbf{A}}}{M_B}\right) |\Omega : \tilde{\mathbf{A}}, \theta\rangle \\ &= \int d\tilde{\mathbf{A}} \mathbf{H}_n\left(\frac{\tilde{\mathbf{A}}}{M_B}\right) \sum_{\mathcal{M}} |\Omega : \tilde{\mathbf{A}} - \mathcal{M}, \mathcal{M}\rangle \frac{1}{\sqrt{2\pi}} e^{-i\theta\mathcal{M}} \end{aligned} \quad (4.74)$$

of H_{reg} . The index n in $|\Omega_n; \theta\rangle$ or $|\Omega_n, \mathcal{M}\rangle$ indicates the number of Schwinger bosons with momentum $\mathbf{P} = 0$. Consequently, the state $|\Omega_0; \theta\rangle$ is the physical vacuum. The states labeled with $\mathcal{M}$ are eigenstates of H_{reg} and the naïvely axial charge : $\mathcal{Q}_A := 2\mathcal{M}$. States labeled with θ are eigenstates of H_{reg} and the heat-kernel regularised axial charge $\mathcal{Q}_A^{\text{reg}} = 2\mathcal{M} + 2\mathbf{A}_3 + 1$. Only the latter states are physical because the former states are not gauge-invariant⁶. All eigenstates of the Hamiltonians under consideration can now simply be obtained by applying the creation operators $\mathbb{A}_{\mathbf{P}}^\dagger$ onto the respective vacua. A corollary of theorem (7) reads as follows

Theorem 8 (Constituent Picture) *The eigenstate $\mathbb{A}_{\mathbf{P}}^\dagger |\Omega_n : 0\rangle$ of H_{reg} contains either right-movers and zero modes only ($P^3 > 0$), left-movers and zeromodes only ($P^3 < 0$) or pure gauge configurations only ($P^3 = 0$).*

⁶One might question that θ states are more physical than axial states. This question is interesting but hypothetical since only for massless fermions does this question arise. We do not observe massless particles which bear colour.

In other words: the state of a Schwinger boson is a TOPOLOGICAL GLUEBALL in the rest-frame and a *pure two-fermion state* in all other frames as long as $M_B \ll \Delta k$ and as long as this state is an eigenstate to $\hat{\mathcal{M}}$ rather than to $\mathcal{Q}_A^{\text{reg}}$. Proof: the state

$$\begin{aligned} \mathbb{A}_{\mathbf{P}}^\dagger |\Omega_n : 0\rangle &= \frac{1}{\sqrt{P^3}} \sum_{1 \leq k^3 \leq P^3} a_{R,k^3}^\dagger a_{R,k^3-P^3} |\Omega_n : 0\rangle \\ &= \frac{1}{\sqrt{P^3}} \sum_{1 \leq k^3 \leq P^3} b_{k^3}^\dagger d_{P^3-k^3}^\dagger |\Omega_n : 0\rangle \end{aligned} \quad (4.75)$$

contains only right-movers for $P^3 > 0$ as is easily seen in Manton's a-basis. This is illustrated in **Fig. 4.2** for the case $P^3 = 3$. Analogously for $P^3 < 0$. The vacuum $|\Omega_0 : 0\rangle$ does not contain fermions. Neither does a boson at rest $\mathbb{A}_0^\dagger |\Omega_0 : 0\rangle = |\Omega_1 : 0\rangle$. The vacuum state and bosons at rest are purely gluonic except that the presence of fermions is responsible —via the anomaly— for the potential energy $(\tilde{A} + 1/2)^2$ of the boson. The potential energy $(\tilde{A} + 1/2)^2$ may thus be considered as an effective interaction due to the presence of fermions. These features bear some resemblance with zero-mode dominance of QCD glueballs described by Van Baal et al. [28, 29].

The $\mathcal{M} = 0$ vacuum and the bosonic states are even simpler for the axial or the artificial Hamiltonian: Their $\mathcal{M} = 0$ vacuum is the perturbative vacuum $|0\rangle$ and their Schwinger boson state

$$\mathbb{A}_{\mathbf{P}}^\dagger |0\rangle = \frac{1}{\sqrt{P^3}} \sum_{1 \leq k^3 \leq P^3} b_{k^3}^\dagger d_{P^3-k^3}^\dagger |0\rangle \quad (4.76)$$

is built on this vacuum without gluonic structure.

5 Theta Vacua and Cut-Off Regularisation

Since the vacua $|\Omega : A_3, \mathcal{M}\rangle$ of H_{reg} are degenerate, every linear combination of these vacua is again a vacuum. For $\Lambda = \infty$, a special case of this statement

perturbative vacua $ \Omega \dots\rangle, L \rightarrow 0$	Symbol
perturbative axial vacuum	$ \Omega : \mathbf{A}_3, \mathcal{M}\rangle$
perturbative θ -vacuum	$ \Omega : \tilde{\mathbf{A}}; \theta\rangle$
n boson vacua, $L \rightarrow 0$	Symbol
perturbative axial vacuum	$ \Omega_n : \mathcal{M}\rangle$
perturbative θ -vacuum	$ \Omega_n; \theta\rangle$
artificial vacua $ \Psi \dots\rangle, L \rightarrow \infty$	Symbol
artificial axial vacuum	$ \Psi : \mathbf{A}_3, \mathcal{M}\rangle$
artificial θ -vacuum	$ \Psi : \tilde{\mathbf{A}}; \theta\rangle$
n boson vacua, $L \rightarrow \infty$	Symbol
full axial vacuum	$ \Psi_n : \mathcal{M}\rangle$
full θ -vacuum	$ \Psi_n; \theta\rangle$

Table 4.1: Vacua: a systematic collection

are the θ vacua defined in (4.74). States like

$$|\Omega : \tilde{\mathbf{A}}; \theta, N_\theta\rangle = \sum_{\mathcal{M}=-N_\theta}^{N_\theta} |\Omega : \tilde{\mathbf{A}} - \mathcal{M}, \mathcal{M}\rangle \frac{1}{\sqrt{2\pi}} e^{-i\theta\mathcal{M}} \quad (4.77)$$

are an other special case. These states are no irreducible representations of the gauge group; for they are not eigenstates of the gauge-transformation operator U_G unless $N_\theta = \infty$. They may, however, be considered as an approximation of θ vacua in the sense that the expectation value of U_G with respect to these states is approximately $e^{i\theta}$. We introduce a CYCLIC GROUP $\mathfrak{C}_{2N_\theta+1}$ with generator $U_G^{(N_\theta)}$ defined as

$$U_G^{(N_\theta)} |\Omega : \mathbf{A}_3, \mathcal{M}\rangle \stackrel{def}{=} \begin{cases} |\Omega : \mathbf{m}(\mathbf{A}_3 - 1), \mathbf{m}(\mathcal{M} + 1)\rangle & \text{if } |\mathcal{M}| \leq N_\theta \\ |\Omega : \mathbf{A}_3, \mathcal{M}\rangle, & \text{else} \end{cases} \quad (4.78)$$

where $\mathbf{m}(a) \stackrel{def}{=} [(a + N_\theta) \bmod (2N_\theta + 1)] - N_\theta$ denotes the MODULO DIVISION which imposes the periodicity $2N_\theta + 1$ upon the argument a . The states

$|\Omega : \tilde{\mathbf{A}}; \theta, N_\theta\rangle$ which we shall call CYCLIC VACUA form irreducible representations of this group if

$$\theta = \theta_n \stackrel{def}{=} \frac{\pi n}{N_\theta + 1} \quad (4.79)$$

with $n \in \mathbb{Z}$. Therefore, $\mathfrak{C}_{2N_\theta+1}$ may be considered as a finite approximation to the infinite gauge group. We shall use the irreducible representations of $U_G^{(N_\theta)}$ in $\langle \mathbf{5} \rangle$ for the special case $N_\theta = 0$. **i.e. we shall consider right-movers and zero-modes only.** According to (4.79), this implies that we shall be able to describe two θ angles: $\theta = 0$ and $\theta = \pi$. If $N_\theta \ll \Lambda$ and if H




Figure 4.2: **Symbolic representation of a Schwinger boson.** The circles represent right-handed fermions filling the Fermi sea from below. The rounded squares represent left-handed fermions filling the Fermi sea from above.

is restricted to a subspace of n -boson states with $n|\mathbf{P}| + N_\theta < \Lambda$, then $U_G^{(N_\theta)}$ is a genuine symmetry of the restricted Hamiltonian. This allows the infinite gauge group to be reduced in a way suitable for numerical computations.

6 Conclusions for Numerical Computations

When diagonalising the *massless* Schwinger model, we have the choice between gauge-invariant θ eigenstates and gauge-variant axial eigenstates. The eigenstates of the *massive* Schwinger model, however, are necessarily θ states. This renders numerical computations (in a partonic framework) very difficult since θ states are states with an infinite parton content. Replacing Coulomb gauge by axial gauge seems to solve this numerical problem. Alas, axial gauge is unphysical and introduces spurious eigenstates. Herin we propose a simple numerical technique which allows us to identify and repair the damage inflicted on the mass spectrum by axial gauge. In Chapter [5](#), we shall demonstrate that and how this technique works.

We have shown that the spectrum of the artificial Hamiltonian and the heat-kernel regularised Hamiltonian are identical for all momenta $\mathbf{P}$ except for $\mathbf{P} = 0$. This fact has important consequences for numerical calculations in the unphysical axial gauge. Axial gauge destroys the invariance of the Hamiltonian under large gauge transformations (gauge invariant regularisation does not make sense in this gauge). Since the degeneracy of ground states of the Hamiltonian is a result of the invariance of the Hamiltonian under large gauge transformations, axial gauge destroys the degeneracy of the axial vacua except for $\mathcal{M} = 0, -1$. States with $\mathcal{M} \neq 0, -1$ manifest themselves as spurious states polluting the mass spectrum. The spectrum of

states with $\mathcal{M} \neq 0, -1$, however, does not contain spurious states. Hence it is correctly described by H_{axial} (except for the rest frame spectrum ($\mathbf{P} = 0$) the only frame affected by axial gauge).

The discussion above suggests that the problem of spurious states can be solved—even for $\mathcal{M} \neq 0, -1$ —by the following *prescription* (1) Diagonalise the axial Hamiltonian (2) subtract the energy $\mathcal{M}(\mathcal{M} + 1)$ of the spurious state in the $: \mathcal{Q}_A := 2\mathcal{M}$ sector from the energies E'_n of all other states in this sector $E_n = E'_n - \mathcal{M}(\mathcal{M} + 1)$. This procedure is equivalent to a computation of the mass spectrum of the artificial Hamiltonian and, therefore, restores the degeneracy of the mass spectrum (except for the rest-frame $\mathbf{P} = 0$). Furthermore, it is easy to show that the degeneracy of $2N_\theta$ states is restored as well if the energy of the vacua (4.77) is subtracted from the energies of states built on these vacua. In Chapter 5, finally, we shall demonstrate that this technique works even for the *massive* Schwinger model (if the fermion masses are small) where only the θ vacua are degenerate whereas the boson-masses acquire a θ dependence: it will be shown that the θ dependence of the vector boson and the scalar boson is correctly reproduced by the above-described technique.

Needless to say that the anomaly is irreparably destroyed by the axial gauge if we choose the rest-frame.

7 The Schwinger Model on a Large Lattice

We have seen, that a CONSTITUENT PICTURE arises for $L \approx 0$. For large volumes $2L$, however, the perturbative vacua (labeled with Ω) are no longer eigenstates of the Hamiltonian. But it is not difficult to construct vacua belonging to large volumes in terms of the already constructed zero-volume

vacua. The operators $\mathcal{A}_{\mathbf{P}}$ ($P^3 > 0$) can be expressed

$$\mathcal{A}_{\mathbf{P}} = \frac{\mathbb{A}_{\mathbf{P}}(|\mathbf{P}| + \Omega(\mathbf{P})) + \mathbb{A}_{-\mathbf{P}}^\dagger(|\mathbf{P}| - \Omega(\mathbf{P}))}{2\sqrt{\Omega(\mathbf{P})}|\mathbf{P}|} \quad (4.80)$$

$$\mathcal{A}_{-\mathbf{P}}^\dagger = \frac{\mathbb{A}_{\mathbf{P}}(\Omega(\mathbf{P}) - |\mathbf{P}|) + \mathbb{A}_{-\mathbf{P}}^\dagger(|\mathbf{P}| + \Omega(\mathbf{P}))}{2\sqrt{\Omega(\mathbf{P})}|\mathbf{P}|} \quad (4.81)$$

in terms of the $L \rightarrow 0$ basis and therefore, a finite-volume vacuum $|\Psi \dots\rangle$ (all these vacuum states are represented in **Tab. 4.1**, e.g. $|\Psi \dots\rangle = |\Psi_n : \theta\rangle$) corresponding to a large volume Hamiltonian H_{reg} is obtained as the solution to the equations

$$\mathcal{A}_{\mathbf{P}} |\Psi \dots\rangle \equiv 0 \quad (4.82)$$

Since this equation allows to construct $|\Psi_0; \theta\rangle$ in terms of $|\Omega_0; \theta\rangle$ and the $\mathbb{A}$ operators and since we know how the $\mathbb{A}$ operators are built from fermions, we are able to construct $|\Psi_0; \theta\rangle$ in terms of fermions. In fact we are able to construct a large-volume state (labeled with Ψ) from any zero-volume state (labeled with Ω) by means of this procedure. The large-volume state

$$|\Psi \dots\rangle = U_B(L) |\Omega \dots\rangle \quad (4.83)$$

may formally be obtained by applying the unitary operator $U_B(L)$ on a small-volume state $M_B L \rightarrow 0$. $U_B(L)$ can be explicitly constructed in terms of $\mathbb{A}_{\mathbf{P}}$ operators. But for our purposes, we do not need the explicit expression; what we are interested in are fermion distribution functions. For an explicit construction, the reader is referred to Ref. [92, 93]. The reason for this simple relation between zero-volume and large-volume states is the fact that the spectrum of the Schwinger model is the spectrum of a free theory. Note that for $L = \infty$ neither the $\mathcal{A}$ basis nor the $U_B(L)$ operator are well defined in terms of fermionic degrees of freedom. This is an illustration of Haag's theorem [46].

8 The Vacuum Distribution Function: Axial Vacua

Let $|\Psi \dots\rangle$ be one of the *axial* vacua listed in **Tab. 4.1**. It is now elementary to calculate the distribution of $\mathbb{A}$ bosons inside the vacuum $|\Psi \dots\rangle$

$$h(\mathbf{k}) \stackrel{def}{=} \langle \Psi \dots | \mathbb{A}_{\mathbf{k}}^\dagger \mathbb{A}_{\mathbf{k}} | \Psi \dots \rangle = \frac{(|\mathbf{k}| - \Omega(\mathbf{k}))^2}{4|\mathbf{k}|\Omega(\mathbf{k})} \quad (4.84)$$

or inside the large-volume Schwinger boson $\mathcal{A}_{\mathbf{P}}^\dagger |\Psi \dots\rangle$

$$h(\mathbf{k}; \mathbf{P}) \stackrel{def}{=} \langle \Psi \dots | \mathcal{A}_{\mathbf{P}} \mathbb{A}_{\mathbf{k}}^\dagger \mathbb{A}_{\mathbf{k}} \mathcal{A}_{\mathbf{P}}^\dagger | \Psi \dots \rangle = \delta(\mathbf{k} - \mathbf{P}) + h(\mathbf{k}) \quad . \quad (4.85)$$




Figure 4.3: **The meson cloud.** The probability of finding a $\mathbb{A}$ boson inside the vacuum for lattices with small but finite size.

The PARTON DISTRIBUTION of quarks and anti-quarks inside the zero-volume state $\mathbb{A}_{P^3} |\Omega_n : \mathcal{M}\rangle$, in turn, can be calculated

$$\begin{aligned}
f(\mathbf{k}, P^3; \Omega : \mathcal{M}) \Delta k^3 &\stackrel{def}{=} \langle \Omega_n : \mathcal{M} | \left[b_{\mathbf{k}}^\dagger b_{\mathbf{k}} - d_{-\mathbf{k}}^\dagger d_{-\mathbf{k}} \right] | \Omega_n : \mathcal{M} \rangle \\
&= \langle \Omega_n : \mathcal{M} | \left[a_{R,\mathbf{k}}^\dagger a_{R,\mathbf{k}} + a_{L,\mathbf{k}}^\dagger a_{L,\mathbf{k}} - 1 \right] | \Omega_n : \mathcal{M} \rangle \\
&= \theta(\mathcal{M} + 1 \leq k \leq \mathcal{M} + P^3) / P^3 \\
&\quad - \theta(\mathcal{M} \leq k \leq \mathcal{M} + P^3 - 1) / P^3
\end{aligned} \tag{4.86}$$

using Mantons solution (graphically represented in **Fig. 4.2**). We also need the RAW DISTRIBUTION FUNCTIONS for $\mathcal{M} = 0$

$$f(\mathbf{k}, P^3; \Omega : 0) \Delta k^3 \stackrel{def}{=} \langle \Omega_n : 0 | b_{\mathbf{k}}^\dagger b_{\mathbf{k}} | \Omega_n : 0 \rangle = \theta(1 \leq k \leq P^3) / P^3 \tag{4.87}$$

and the corresponding raw distribution of an anti-quark

$$\bar{f}(\mathbf{k}, P^3; \Omega : 0) \Delta k^3 \stackrel{def}{=} \langle \Omega_n : 0 | d_{\mathbf{k}}^\dagger d_{\mathbf{k}} | \Omega_n : 0 \rangle = \theta(0 \leq k \leq P^3 - 1) / P^3 \tag{4.88}$$

which follow from Eq (4.76). Here, $\theta(x)$ denotes the Heaviside step function. The quark distribution inside the *large volume* vacuum $|\Psi_0, 0\rangle$ is approximately the convolution

$$f(\mathbf{k}; \Psi : 0) \approx \sum_{\mathbf{P}} \Delta k^3 h(\mathbf{P}) f(\mathbf{k}; P^3 : \Omega : 0) \tag{4.89}$$

of these two functions. If $h(\mathbf{P})$ is small so is the convolution. We use this fact in order to demonstrate that the approximations utilised in Chapter **5** are accurate, i.e. we can easily show now that the vacuum is almost trivial for $g = \Delta k$ a value of g that we shall use in Chapter **5**. The graph of h is represented in **Fig. 4.3** for $g = \Delta k$. We conclude that smallness of h justifies the assumption of Chapter **5** that the vacuum is almost trivial. Moreover, we conclude from Eq (4.84) that the total number of particles in the vacuum

$\mathcal{N}_{\text{vac}} \stackrel{\text{def}}{=} \sum_{\mathbf{k}} h(\mathbf{k})$ diverges in the limit $L \rightarrow \infty$ (i.e. $M/\Delta k \rightarrow \infty$): another illustration of Haag's theorem. It is useless to start with $L = \infty$ and — strictly speaking— even wrong.

9 The Vacuum Distribution Function: Gauge Invariant Vacua

So far we have been calculating distribution functions in axial states. Physical states, however, are gauge-invariant. In a *gauge-invariant* θ state, the expectation value of the particle number operator $a_{\alpha,\mathbf{k}}^\dagger a_{\alpha,\mathbf{k}}$ is no longer well-defined because the number of partons is no gauge-invariant quantity. In order for distribution functions of θ vacua to be well defined, they have to be defined relative to the fermion level, i.e. via the gauge invariant particle number operators $a_{\alpha,\mathbf{k}+\mathcal{M}}^\dagger a_{\alpha,\mathbf{k}+\mathcal{M}}$ or $a_{\alpha,\mathbf{k}-\mathbf{A}_3}^\dagger a_{\alpha,\mathbf{k}-\mathbf{A}_3} = a_{\alpha,\mathbf{k}+\mathcal{M}-\tilde{\mathbf{A}}}^\dagger a_{\alpha,\mathbf{k}+\mathcal{M}-\tilde{\mathbf{A}}}$. In the first case, the distribution functions of θ states are identical to the distribution functions in states with $\mathcal{M} = 0$: the calculation of gauge-invariant distribution functions in a θ state can thus be replaced by the simpler calculation of a gauge-dependent distribution function in a axial state. In the second case, the distribution functions of θ states are smeared through the oscillation of $\tilde{\mathbf{A}}$ with an oscillation amplitude of roughly $\Delta\tilde{\mathbf{A}} \approx M_B$.

In fact, structure functions are related to distribution functions of the second type (cf. the remark on p.24) if axial gauge cannot be chosen: Therefore, gauge-invariant distribution functions of theta states are obtained by "smearing out" gauge-variant distribution functions of axial states. These considerations illustrate nicely why the inclusion of a gauge string U_A into the correlation function $\tilde{\mathbf{q}}_\mu(y|PS)$ is necessary.

10 Epilogue: The Front Form and the Schwinger Model

If the Schwinger model is quantised in the FF, only right-movers are present. We have already demonstrated that this quantisation is not equivalent to QED but rather to chiral QED in zero volume. In this theory, $: \mathcal{Q}_A =: \mathcal{Q} := 0$. Therefore, θ vacua are impossible to implement in this approach⁷, as they constitute superpositions of states with different axial charge $\mathcal{Q}_A$. McCartor and Robertson [94] treated the Schwinger model by quantising the fermionic fields on two light-like quantisation surfaces. They quantised right-movers on the quantisation surface $x^+ = 0$ and left-movers on the quantisation surface $x^- = 0$. This allowed them to include θ vacua. They found, however, that the condensate becomes zero in the limit of infinite light-like periodicity. We are now able to explain this puzzle. Consider the quantisation surface used by McCartor and Robertson depicted in **Fig. 4.4**. Also depicted is an equivalent quantisation surface which we shall prefer for reasons that will become clear soon. The proper way of quantising on a 'bent' quantisation surface is using bent co-ordinates $\tilde{x}^\mu$ defined as

$$\tilde{x}^3 \stackrel{def}{=} x^3 \quad \text{and} \quad \tilde{x}^0 = \begin{cases} x^+ & \text{if } x^3 < 0 \\ x^- & \text{if } x^3 \geq 0 \end{cases} \quad (4.90)$$

in terms of which the bent quantisation surface reads $\tilde{x}^0 = 0$.

Now we have to impose boundary conditions upon the fields φ . If we require periodicity in $\tilde{x}^3$ then this periodicity is *space*-like because the four-vector $2\mathfrak{L}$ which joins A and B is space-like. As such, $\mathfrak{L}$ defines a finite, non-zero

⁷I.e. they cannot be implemented using *periodic BCs*




Figure 4.4: Admissible BCs for the FF

renormalisation scale. Quantisation in a space with space-like periodicity, however, is most conveniently performed in the IF formalism; quantisation in bent co-ordinates is much more cumbersome to put it mildly.

One might as well—following McCartor and Robertson—introduce TWO *light*-like BCs on the right-moving and the left-moving component respectively joining the point O with the points A and B respectively. This defines two *light*-like boundary four-vectors. Consequently, one is working in an infinitesimally small box. Such a theory is not equivalent to the IF as such. Such a theory is equivalent to the STRONG IMF instead, i.e. to the IF on an infinitesimally small lattice $L = 0$.

In the Schwinger model, the kinetic energy does not necessarily render left-movers infinitely heavy in the IMF. The reason for this is that the kinetic energy T is replaced by the modified kinetic energy $\mathfrak{T}$ which is zero for every axial vacuum even though axial vacua may contain an arbitrary number of left-movers. Consequently, the FF Schwinger model is not an effective the-

ory for the IMF Schwinger model because an effective theory for the IMF Schwinger model must not exclude the left-movers which are responsible for the θ vacuum. What is the effective Hamiltonian then? Even though we cannot eliminate left-movers, we are able to eliminate every operator which *mixes* left-movers and right-movers. The justification for this is the fact that axial vacua $|\Omega_0 : \mathcal{M}\rangle$ are exact solutions to the zero volume Schwinger model, as we have already shown. This means that operators which enforce mixing of left-movers and right-movers are completely suppressed in the limit $L \rightarrow 0$: From these considerations we conclude that the effective IMF Hamiltonian for the Schwinger model is the IMF Hamiltonian with left-right-mixing operators removed. Once these terms are removed, the plus-component $P^+(e^{(0)})$ of the IF quantised Hamiltonian consists merely of right-movers and the minus-component $P^-(e^{(0)})$ consists merely of left-movers. This effective Hamiltonian has now exactly the same structure as a theory constructed on two light-fronts [94]. This also explains why this theory is unable to describe the axial condensate of the Schwinger model. The condensate is due to the fluctuating particle number in the vacuum generated by the Bogolubov transformation $U_B(L)$: it can only be correctly described on a large volume $L \rightarrow \infty$. The volume of the strong IMF, however, is infinitesimally small and so is the volume of the two-light-front theory [94]. We shall elaborate these arguments in a later publication. We summarise: **The two-light-front theory is equivalent to the zero volume Schwinger model whereas the FF is equivalent to the chiral zero volume Schwinger model.**

Bornons ici cette carrière.
Les longs ouvrages me font peur.
Loin d'épuiser une matière
On n'en doit prendre que la fleur.
Il s'en va temps que je reprenne
Un peu de forces et d'haleine
Pour fournir à d'autres projets.

[La Fontaine: Livre VI]

Chapter 5

The Massive Schwinger Model

In this chapter we diagonalise the Hamiltonian of the massive Schwinger model in a frame where the VECTOR BOSON (i.e. the lightest physical particle) with four-momentum

$$P = (E, 0, 0, P^3) \tag{5.1}$$

moves at a velocity $v^3 = P^3/E$ close to the speed of light. This frame will be referred to as LARGE MOMENTUM FRAME (LMF). We compare our results with the computations of Hamer et al. [95] who used the Kogut-Susskind Hamiltonian¹. The LMF has several advantages. *Firstly*, the RADIUS R of the lightest physical particle is Lorentz-contracted to

$$R' = \sqrt{1 - v_3^2} R \tag{5.2}$$

which is smaller than the CORRELATION LENGTH $a\xi = \frac{1}{M}$ if the velocity v^3 is sufficiently high (The actual size of R is not important for our argument). $M = M_V$ is the mass of the vector boson. In the rest-frame, in contrast, the correlation length of a particle is usually smaller than its extension. Therefore, the physical requirement that *both* the correlation length

¹For the first computations with Wilson fermions see [96–99]. Since we are working in momentum space, we do not have to deal with fermion doubling.

and the physical size of the boson be smaller than the size L of the lattice

$$a\xi, R < 2L \quad (5.3)$$

may be replaced by the much weaker condition

$$a\xi < 2L \quad (5.4)$$

which allows us to use a LATTICE SIZE L much smaller than if we had chosen the rest-frame— albeit not an infinitely small one. This advantage is crucial since pair-creation leads to a finite density of VIRTUAL PARTICLES in any interacting field theory. The number of virtual particles is thus roughly proportional to the volume and, consequently, the total number of particles is drastically reduced if the volume $2L$ is small. A *second* advantage is the fact that most constituents inside a fast-moving object can be expected to move in the same direction as the fast moving object itself *as long as* the number of VIRTUAL PARTICLES ASSOCIATED WITH THE VACUUM can be neglected when compared to the number of virtual particles associated with the moving object. Under such circumstances, the neglect of LEFT-MOVERS, i.e. particles moving in the direction of the negative 3-axis, may thus be a very good approximation. Having no left-movers, in turn, constitutes a *third advantage* since it renders the number of states in the Fock space finite (except for bosonic zero-modes) as has been explained in Ref. [100, 101]. We are going to show now that this approximation results indeed in an excellent description of the Schwinger-boson mass as well as of the condensate of the massless Schwinger model.

The defining Lagrangian density of the massive Schwinger model is

$$\mathcal{L} = \bar{\psi}(i \not{D} + m)\psi - \frac{1}{4g^2} F^{\mu\nu} F_{\mu\nu} \quad (5.5)$$

where m , g and $D = \partial + i\mathbf{A}$ denote the mass of the fermions, the coupling constant and the covariant derivative, respectively. A^μ is the gauge-field connection and $\mathbf{F}$ the corresponding field-strength. We have absorbed

$$\mathbf{A}^\mu = gA^\mu \quad (5.6)$$

the coupling constant g in the photon fields A^μ . Canonical quantisation in axial gauge $A^3 = 0$ yields the Hamiltonian

$$H = \int_{-L}^L dx^3 (\bar{\psi} \gamma^3 i \partial_3 \psi + m \bar{\psi} \psi) + \frac{g^2}{2} \int_{-L}^L dx^3 (\psi^\dagger \psi) \frac{1}{-\partial_3^2} (\psi^\dagger \psi) \quad (5.7)$$

where we have introduced a finite lattice size $2L$. Periodic boundary conditions are assumed from now on. Strictly speaking, axial gauge is incompatible with periodic boundary conditions since the global Wilson loop

$$\exp(i \int_{-L}^L dx^3 \mathbf{A}^3) \quad (5.8)$$

is a physical observable. It will turn out, however, that removing this physical degree of freedom does not hamper the computation of the mass-spectrum and the vacuum condensate if the lattice size is sufficiently small. The deeper reason for this is the fact that topological effects are less important when working in tiny volumes as we have already demonstrated in Chapter 4

We expand the fermionic fields

$$\psi(t=0, \mathbf{x}) = \sum_{\mathbf{k}} \frac{1}{2\sqrt{L\omega(\mathbf{k})}} (u_{\mathbf{k}} b_{\mathbf{k}} e^{+i\mathbf{k}\cdot\mathbf{x}} + v_{\mathbf{k}} d_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{x}}) \quad (5.9)$$

in the usual way in terms of annihilation and creation operators where

$$\omega(\mathbf{k}) = \sqrt{m^2 + \mathbf{k}^2} \quad (5.10)$$

spinors appearing in this equation are defined as solutions of the free, massive Dirac equation with the normalisation:

$$\bar{u}_k u_k = 2m \quad (5.11)$$

$$\bar{v}_k v_k = -2m \quad . \quad (5.12)$$

We diagonalise the Hamiltonian in the $\mathbf{P} = 0$ sub-sector of the total Fock space and obtain the vacuum energy E'_0 . This sub-sector is spanned by a finite number of Fock-states since we have

1. truncated the total Fock space to states that do not contain any left-moving particles with $k^3 < 0$
2. at most $N = P^3/\Delta k$ particles with $k^3 > 0$ are able to share the total momentum P^3
3. there is only a finite number (i.e. two) of fermionic zero-modes due to the Pauli exclusion principle
4. we have removed the bosonic zero-mode by our (unphysical) choice of axial gauge (contrary to fermions, there may be an arbitrary numbers of bosons at $\mathbf{k} = 0$).

In fact, there are two vacua corresponding to theta angles $\theta = 0$ and $\theta = \pi$. If we had restricted the fermion momenta to the region

$$-n\Delta k \leq k^3 \leq P^3 + n\Delta k \quad (5.13)$$

we would have had $2n$ θ -vacua (or rather approximants thereof, cf. (4.77)). No restriction on the momenta at all yields an infinity of them (the number of these vacua is, of course, equal to the number of distinct values that the naïve axial charge can take on). Here, we shall primarily be concerned with the case $\theta = 0$ as our approximation is able to describe the sector $\theta = \pi$ for small fermion masses only.

Numerical diagonalisation of the Hamiltonian for the momentum $\mathbf{P}$ yields the energy spectrum E'_n (relative to the energy of the perturbative vacuum).

The physical energies are now

$$E_n(\theta) = E'_n(\theta) - E'_0(\theta) \quad (5.14)$$

where $E'_0(\theta)$ is the energy of a θ -vacuum. The physical masses, in turn, are obtained from the relativistic dispersion relation

$$M_n = \sqrt{E_n^2 - \mathbf{P}^2} \quad (5.15)$$

Let $M = M_1 = \frac{1}{a\xi}$ be the mass of the VECTOR BOSON, i.e. the boson with smallest mass. $\Lambda = \frac{\pi}{a}$ is the cut-off and $a \rightarrow 0$ is the lattice-spacing. As in the case of the massless Schwinger model, we are allowed to the continuum-limit $a \rightarrow 0$. The only renormalisation necessary in the *super-renormalisable* Schwinger model is the subtraction of the vacuum energy E'_0 .

1 Scaling Window and Region of Validity

On a finite lattice, not every pair of parameters m, g is physically meaningful. It is well-known that the computed mass spectrum is physically meaningful [4] only if the correlation length $a\xi$ is smaller than the lattice size $2L$ and larger than the lattice spacing a . Therefore, the parameters m and g have to be tuned such that the computed correlation length lies inside the scaling window

$$a < \xi a < 2L \quad . \quad (5.16)$$

We have set $a = 0$ and therefore the scaling window seems to be open

$$\xi a < 2L \quad (5.17)$$

in the ultra-violet region ². The physical region in the parameter space, however, is smaller than the scaling window (5.16) since we have used the

²(it is remarkable that — in contrast to the φ^4 model— the cut-off $\Lambda = \pi/a$ cannot be made smaller than $2P^3$ without a massive deterioration of the numerical results)

approximation that left-moving particles are not important— an assumption which can only be justified if the velocity $v^3 = P^3/E$ of the boson is sufficiently close to the velocity of light, i.e. if $Ma = \xi^{-1} \ll aP^3$. Therefore, our results can only be trusted inside the *accuracy window*

$$(1/P^3) \ll \xi a < 2L \quad . \quad (5.18)$$

This constraint is still not strong enough, however, since it only ensures the absence of left-moving virtual particles associated with the moving object. Since the vacuum in interacting relativistic field theories is not empty, there are always VIRTUAL PARTICLES ASSOCIATED WITH THE VACUUM (rather than with the moving object itself). In particular, the parity invariance of the vacuum implies that it contains an equal number of right-movers and left-movers. The mere presence of a fast-moving physical particle does not suffice to modify the properties of the vacuum, let alone render it trivial. This is why we choose the length $2L$ of the system as small as possible such that the number of particles associated with the vacuum is negligible or such that at least most particles associated with the vacuum are concentrated at $\mathbf{k} = 0$. The choice $L = \infty$ right from the onset is impossible [46] and unnecessary. In the context of lattice gauge theory one works in a volume $2L$ which is large enough to comfortably contain the physical object of diameter $2R$ and yet small enough for practical computations. Our choice of the moving frame allows us to boost the size $2R'$ of the physical object to arbitrarily small values thus enabling us to further reduce the number of vacuum particles dramatically when compared to what is possible in the rest-frame.

Naively we could even reduce the number of particles in the vacuum to zero (and all extensive quantities with it) by the extreme choice of $L \approx 0$ which seems to be legitimate in the INFINITE MOMENTUM FRAME (IMF) $v^3 \rightarrow 1$ where the Lorentz-contracted SIZE $R\sqrt{1-v_3^2} \rightarrow 0$ of any physical

object vanishes(cf. Chapter $\langle 3 \rangle$). This, however, would be in blatant contradiction with the scaling window (5.16) which requires that the size of the lattice be larger than $1/M_1$. To our knowledge, this deficiency of the IMF has nowhere been discussed in the literature. We shall present our results for a coupling of $g = \Delta k = \pi/L = 1$ corresponding to a Schwinger-boson mass of

$$M_B = \frac{g}{\sqrt{\pi}} \approx 0.56 \quad (5.19)$$

which minimises the volume and yet is compatible with the scaling window: The volume $2L = 2\pi$ is then small enough to drastically reduce the number of particles in the vacuum and yet sufficiently large for the correlation length $1/M_B \approx 1.77 < 2L = 2\pi$ to lie comfortably inside the scaling window (5.16): We have already demonstrated in chapter $\langle 3 \rangle$ that the vacuum of the exact solution to the massless Schwinger model is indeed almost trivial for $\Delta k = \pi/L \approx g$. Since the number of virtual particles in the vacuum must decrease when they become heavier, it follows that the vacuum is also trivial for $m/g > 0$ ($g \approx \Delta k$). For $m/g > 0$ the vector boson mass $M_1(m, g)/\Delta k$ increases and we could further reduce the volume in physical units $2LM_1 = 2L/(a\xi)$ (i.e. in units of $a\xi$) by choosing an even smaller value of g without leaving the window. Indeed, fixing $M_1(g)$ at the value of $M(g, m) \approx M_1(g, 0) = \Delta k/\sqrt{\pi}$ accelerates the convergence of physical quantities for larger fermion masses. At given $m/g \approx 0.25$, for instance, the convergence is approximately 2.3 times better for $g = 1/\sqrt{\pi}$ than for $g = 1$

$$\frac{M_1^{(N=384)}(g = 1.0) - M_1^{(N=6)}(g = 1.0)}{M_1^{(N=384)}(g = \frac{1}{\sqrt{\pi}}) - M_1^{(N=6)}(g = \frac{1}{\sqrt{\pi}})} \approx 2.3 \quad (5.20)$$

and the distance between $M^{(N)}(m/g = 0.25, g = 1/\sqrt{\pi})$ and the corresponding mass of chiral perturbation theory is divided by a factor of roughly

three when compared to $M^{(N)}(m/g = 0.25, g = 1)$. Inverting the function $M_1(g)$, however, is necessary for this purpose and requires several self-consistent diagonalisations for each value of m/g instead of only one. As our results are already excellent for fixed $g \approx 1$, we prefer the straightforward choice $g = 1$ and do not seek to further increase precision by fixing $M_1(g) \equiv \Delta k / \sqrt{\pi} = \text{const.}$ We observed that for masses as small as $m/g = 0.5$ already, the dependence on the scale $g/\Delta k$ is almost negligible.

2 Convergence and Covariance

Physical particle masses are Lorentz scalars. Therefore, an important test of a relativistic model is the approximate independence of the computed physical masses of the momentum. Diagonalising the mass squared operator in sectors of different momenta must *approximately* yield the same result. Ideally, this should hold for every momentum $-\Lambda \ll P^3 \ll \Lambda$. In praxis, however, the physically relevant region of $\mathbf{P}$ may be further restricted by additional approximations as, for instance, the assumption that only right-movers are important. In what follows, we shall use the requirement that physical masses be scalars in order to localise the momenta for which our approximation *does not* hold: For if the computed masses depend strongly on P^3 in some region of momentum space then this indicates that our approximation cannot be trusted in this region. This is the case for the region of small total momenta P^3 (i.e. momenta that are small compared to the boson mass M_n) because left-movers do become important if the physical particle moves slowly.

A few comments are in order here as care must be taken to make a clear distinction between the CONTINUUM LIMIT $\Lambda \rightarrow \infty$ and what we shall call the COVARIANCE LIMIT $N=P^3/\Delta k \rightarrow \infty$. In the Schwinger model, the only ultraviolet regularisation necessary is the subtraction of the vacuum energy.

Contrary to renormalisable theories, we are able to perform the continuum limit $\Lambda/M_1 \rightarrow \infty$ without rendering the Hamiltonian singular. This means that the actual number of lattice points $2\Lambda/\Delta k$ we are working with is infinite. If, however, only right-moving particles $\mathbf{k} \geq 0$ are important, this implies that particles with momenta $\mathbf{k}$ larger than the total momentum $\mathbf{P}$ cannot play a rôle, either. Hence only particles lying on the EFFECTIVE LATTICE $0 \leq k^3 \leq P^3$ have to be taken into account for the *practical* numerical calculation. Consequently, the quantity which determines the *minimal size* of the Hamiltonian matrix to be diagonalised (which determines the actual numerical effort) is the number $N + 1$

$$N \stackrel{def}{=} \frac{P^3}{\Delta k} \quad (5.21)$$

of lattice points between $\mathbf{k} = 0$ and $\mathbf{k} = \mathbf{P}$ whereas the cut-off Λ primarily influences the *form* of the Hamiltonian. For this reason we shall call the number $N + 1$ of numerically important lattice sites the EFFECTIVE LATTICE SIZE — a number which must not be confused with the real lattice size $2\Lambda/\Delta k = \infty$. One should be careful not to confuse the limit $N \rightarrow \infty$ with the CONTINUUM LIMIT $\Lambda/M_1 \rightarrow \infty$ or with the thermodynamic limit $M_1 L \rightarrow \infty$ ³.

3 Numerical Results

We have computed the mass spectrum and the parton distributions of QED(1+1) in terms of m/g in the ULTRA-RELATIVISTIC REGION $m/g \lesssim 0.3$. For the sake of comparison with the results of Hamer et al. [95] we have used the logarithmically spaced points $m_i/g = 1/8, 1/4, \dots, 16, 32$. We choose

³The last point makes clear that our results can, at best, approximate the correct results since we never perform thermodynamic limit(!)

$g = \Delta k = \pi/L = 1$ for all numerical calculations presented below for the reasons explained in the last chapter. The results for the FF that are used here were published first in [49]. The generality of our computer programme allowed us to recalculate these results in order to compare them with [49] (thus providing us with an additional control of our software) and to create the figures presented in this chapter.




Figure 5.1: **Convergence of distribution functions in the chiral limit** $m/g = 1/8$. The LMF distribution function $f(x_B, P^3)$, defined on the $N + 1$ points of the effective lattice, is represented for $P^3 = N = 6, 12, 24, \dots, 192$. In the COVARIANCE LIMIT $N \rightarrow \infty$, the variable x_B becomes continuous.

4 Distribution Functions

We define the fermion distribution function

$$\mathfrak{f}(x_B; P^3) \triangleq x_B = \mathfrak{f}_{x_B}(P^3) \stackrel{def}{=} \langle P | (b_{x_B \mathbf{P}}^\dagger b_{x_B \mathbf{P}} + d_{x_B \mathbf{P}}^\dagger d_{x_B \mathbf{P}}) | P \rangle \quad (5.22)$$

as the mean number of virtual particles or antiparticles at momentum $\mathbf{k} = x_B \mathbf{P}$ inside the boson state $|P\rangle$. **Fig. 5.1** shows rapid convergence of this function for small fermion masses ($m/g = 1/8$). This indicates that the computed distribution functions become approximately boost invariant for




Figure 5.2: **Convergence of the distribution functions in the FF**, $m/g = 1/8$. The FF distribution function $\mathfrak{f}(x_B, P^3)$, defined on the $N + 1$ points of the effective lattice, is represented for $P^3 = N = 6, 12, 24, \dots, 192$. In the COVARIANCE LIMIT $N \rightarrow \infty$, the variable x_B becomes continuous.

large momenta which is all the more remarkable as distribution functions cannot be boost invariant for *small* momenta P^3 (to see this, set $P^3 = 0$ in Def. (5.22)). The corresponding FF result **Fig. 5.2** also shows approximate boost-invariance. It is remarkable that the distribution functions computed in the large momentum frame (LMF) are boost-invariant to a much higher degree than the ones computed using the FF. This illustrates our comment of chapter 3 that a finite lattice spacing Δk destroys boost-invariance in all relativistic forms including the FF.

The (discrete) integral over the (vector boson) distribution function is 2.0




Figure 5.3: **The number of fermions in a boson.** The number of fermions in the vector boson, the scalar boson and in a two-boson bound-state respectively, plotted against the fermion mass m/g .

for every value m/g of the fermion mass as can be inferred from **Fig. 5.3**. The vector boson state is practically a pure two fermion state in our LMF approximation (the same thing happens in the FF). **Fig. 5.4** shows that the same holds true in the FF (as already remarked in [49]). Indeed, the vector boson mass remains unchanged if we truncate to the Fock space such that states with more than two bosons are excluded. The fact that the number of fermions does not fluctuate implies that the PROBABILITY $\mathcal{P}_{x_B}$ of finding




Figure 5.4: **The number of fermions in a boson (FF)**. The number of fermions in the vector boson, the scalar boson and in a two-boson bound-state respectively, plotted against the fermion mass m/g .

a fermion with momentum fraction x_B is proportional

$$\mathcal{P}_{x_B} = \frac{1}{2} \mathbf{f}_{x_B} \quad (5.23)$$

to the distribution function. This is so because the number operator becomes the projector on the two-particle state $|x_B\rangle \stackrel{def}{=} b_{x_B \mathbf{P}}^\dagger d_{x_B \mathbf{P}}^\dagger |0\rangle$

$$b_{x_B \mathbf{P}}^\dagger b_{x_B \mathbf{P}} = |x_B\rangle \langle x_B| \quad (5.24)$$

if the particle number is fixed. In **Fig. 5.5** we represent convergence of $\mathbf{f}(x_B) = \mathbf{f}(x_B, N\Delta k)$ for an almost non-relativistic value $m/g = 32$ of the




Figure 5.5: **Convergence of distribution functions for $m/g = 32$.** The distribution functions $\mathbf{f}(x_B, N)$ for a large fermion mass ($m/g = 32$) are plotted against the Bjorken scaling variable x_B .

fermion mass. On the one hand, a finer (effective) lattice is required in non-relativistic domain in order to resolve the narrow peak of the distribution function. On the other hand, however, we observe that $f(x_B)$ is zero for $x_B > 0.6$ or for $x_B < 0.4$ which means that we could have considerably reduced the numerical effort by choosing an *effective* lattice between $x_B = 0.4$ and $x_B = 0.6$ which is five times smaller than the original one. **Fig. 5.6** depicts the dependence of the distribution functions on the fermion mass m from $m/g = 1/8$ to $m/g = 32$, **Fig. 5.7** shows the corresponding results from $m/g = 0$ to $m/g = 0.3$. Comparing the LMF distribution functions **Fig. 5.7** with the




Figure 5.6: **Overview of distribution functions on the entire range of fermion masses.** The distribution functions $f(x_B, P^3)$ for different fermion masses $m/g = 1/8, 1/4, \dots, 32$ are plotted against the Bjorken variable x_B .

same functions **Fig. 5.8** obtained in the FF reveals a substantial disparity in the region where x_B or $1 - x_B$ is small. In particular, our LMF result coincides exactly with the exact solution to the Schwinger model (4.89) on a small lattice (see Chapter 4) whereas the FF result differs considerably.




Figure 5.7: **Distribution functions (LMF) in the chiral limit $m/g \rightarrow 0$.** The distribution function $f(x_B; N = 192)$ is plotted against the Bjorken variable x_B for small masses $m/g \lesssim 0.3$.

5 The Mass of the Vector Boson

Fig. 5.9 shows that vector boson mass (mass of the lightest physical particle) as a function of $m/g \lesssim 0.3$. Chiral perturbation theory predicts

$$\frac{M_1(g)}{g} \approx 0.5642 + 1.781 \frac{m}{g} + 0.1907 \left(\frac{m}{g}\right)^2 \quad (5.25)$$

for the vector meson mass [90, 91, 95, 102, 103] for small values of m/g . Agreement with second order chiral perturbation theory as well as with the points computed in Ref. [95] is observed. The excellent convergence in terms of N indicates that the physical mass becomes approximately independent




Figure 5.8: **Distribution functions (FF) in the ultra-relativistic parameter region.** The FF distribution function $f(x_B; N = 192)$ is plotted against the Bjorken variable x_B for small masses $m/g \lesssim 0.3$.

of the momentum even for very moderate values of N . The Schwinger limit $m \rightarrow 0$ is special in that the physical mass M is entirely independent of the momentum for every $N > 0$. Moreover, the mass-spectrum is correctly described even if the correlation length $1/M_1$ is much larger than the volume $2L$. These peculiarities are due to the fact that the (massless) Schwinger model is equivalent to a theory of point-like free bosons. These results have to be contrasted with the results obtained in the FF approach depicted in **Fig. 5.10** where the convergence is so slow that it is hardly possible to accurately es-




Figure 5.9: **The vector boson mass M_V . A comparison with chiral perturbation theory.** The mass of the vector boson (LMF) is plotted against the fermion mass $m/g \lesssim 0.3$ and compared to chiral perturbation theory and the results of Hamer et al. The dimensionless momentum N varies logarithmically according to $3 \cdot 2^i$.

timate the exact boson mass between $m/g = 0$ and $m/g = 0.2$ even on huge lattices. The instant form computation, in contrast, **Fig. 5.9** reproduces the linear dependence of M on the fermion mass even on effective lattices as small as $N = 6$.

We see that the physical mass computed in our IF approximation is approximately boost-invariant already at very small momenta whereas the mass computed in the FF is far from being invariant even for the largest momenta N . This is most remarkable since one of the alleged advantages of the FF




Figure 5.10: **The vector boson mass in the FF. A comparison with chiral perturbation theory.** The mass of the vector boson computed in the **FF formalism** is plotted against the fermion mass $m/g \lesssim 0.3$ and compared to chiral perturbation theory. The dimensionless momentum N varies logarithmically according to $3 \cdot 2^i$.

over the IF is the manifest boost-invariance of the former. We see that in a non-perturbative calculation on *any finite* lattice, neither the LMF *nor* the FF are boost-covariant. The IF, however, is almost boost-covariant even on small lattices whereas a gigantic effective lattice is needed in the FF in order to obtain covariance to the same degree as in the IF.

Non-perturbatively, the IF is boost-invariant to a much higher accuracy on a finite lattice than is the FF which means that it is exceedingly preferable to use the IF in order to optimise covariance.

It has been shown [50] that the FF reproduces approximately the correct




Figure 5.11: **The binding energy of the vector boson.** The binding energy of the vector boson $(M_V - 2m)/g$ is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

vector boson mass on an effective lattice which is infinitely large (i.e. if the limit $N \rightarrow \infty$ is performed). The massive Schwinger model is special in the sense that the FF, the IMF and the LMF coincide on an infinite lattice $N = \infty$. This is why the results of [50] may be regarded as the results of the FF, the IMF and the LMF for an infinite effective lattice. It is easy to show that the absence of (a) four-boson couplings in QED(1+1) and (b) massless mesons is responsible for this result. We shall show later on that this does not hold for the two-flavour massive Schwinger model. **Thence one has to realise that the successful computations in [50] are successful**




Figure 5.12: **The binding energy of the vector boson in the FF.** The binding energy of the vector boson $(M_V - 2m)/g$, computed in the FF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

only because of this quirky coincidence of three theories (FF, IMF, LMF) which do not coincide for most QFTs. A further comment is in order. Even though the results of [50] are quite successful, they are not exact nonetheless. The reason for this is that one has never performed the infinite volume limit because in the infinite volume limit, the vector boson state consists of an arbitrary number of quarks and anti-quarks. Cf. Chapter <4>. It is impossible to see this in the FF since the FF always implies an infinitely small volume and hence precludes the infinite volume limit.

Fig. 5.11 and **Fig. 5.12** depict the BINDING ENERGY $(M_V - 2m)/g$ of the vector boson with mass M_V . One observes that the manifest inequality between the IF and the FF at finite momenta N changes drastically in the non-relativistic limit $m/g \rightarrow \infty$ as well as in the moderately relativistic region $m/g \approx 1$ where the FF and the IF are practically identical. The reason for this is that matrix elements involving only right-movers $k^3 \geq 0$ are practically identical in the LMF, the IMF and the FF. Therefore, the numerical results are almost identical for larger fermion masses $m/g \lesssim 0.25$ since virtual particles with zero momenta $\mathbf{k} = 0$ are not important in this case. A glimpse at the particle distribution function $\mathfrak{f}(x_B; P^3)$ makes this point clear: In **Fig. 5.7**, the distribution function in the LMF is depicted for different values $m/g = 0, 0.05, \dots, 0.3$ of the fermion mass. For $m = 0$, our results coincide with the exact distribution function (4.87)

$$\begin{aligned} \mathfrak{f}(x_B) &= f(x_B P^3, P^3; \Omega : 0) + \bar{f}(x_B P^3, P^3; \Omega : 0) \\ &= \Theta(1 \leq x_B \leq P^3) + \Theta(0 \leq x_B \leq P^3 - 1) \end{aligned} \tag{5.26}$$

computed in Chapter <4>. **Fig. 5.8** represents the same function as computed in the FF. It is clear that the contribution of the zero-momentum sector $\mathbf{k} = 0$ cannot be neglected in the ultrarelativistic region and this is exactly the region where the FF and the LMF differ substantially. At $m = 0$

the probability $\mathcal{P}_0 = \frac{1}{2}f_0 = 0.5/N$ to find a particle at $\mathbf{k} = 0$ is as important as any other mode $0 < k^3 < P^3$ and can not be discarded. The FF distribution function for $m = 0$ differs considerably from the exact result (5.26). Only if the limit $N \rightarrow \infty$ is performed can this mode be neglected; although the probability for finding a particle at $\mathbf{p} = 0$ converges quickly $\propto 1/N$, the influence it exerts on the physical spectrum remains extremely strong even for momenta as large as $N = 384$. A comparison of **Fig. 5.9** and **Fig. 5.10** immediately makes this clear. This explains nicely the following conundrum: while Mo's and Perry's "continuum" version [50] ($N = \infty$ and therefore $\Delta x_B = \Delta k/P^3 = 0$) of the FF is able to reproduce the linear behaviour

$$M_1(m) - M_1(0) \propto m + \mathcal{O}(m^2) \quad (5.27)$$

of the vector boson mass in terms of the fermion mass. But there is always a quadratic behaviour

$$M_1(m) - M_1(0) \propto m^2 + \mathcal{O}(m^3) \quad (5.28)$$

around $m = 0$ for *any* finite N . If $N = \infty$, the probability $\mathcal{P}_0$ of finding a zero-mode fermion is zero, which allows for the possibility that the FF and the LMF are equivalent at $N = \infty$. We note in passing that this result is unlikely to hold in four-dimensional QCD since the experimentally measured quark distribution functions diverge at $x_B = 0$. It should also be stressed that the possible equivalence of the LMF and the FF in the limit $N = \infty$ does by no means imply the equivalence of the FF and the IF: We have already stressed that keeping a small volume prevents us from performing the continuum limit. Therefore, our model is an approximation of the complete IF dynamics. An excellent one— but an approximation it is.

6 The Mass of the Scalar Boson

The BINDING ENERGY $(M_S - 2m)/g$ of the scalar boson mass M_S is shown in **Fig. 5.13** with a logarithmic abscissa for the points used in [95] including the ultrarelativistic region $m \ll g$ and the non-relativistic region. The Fock-space has been truncated to the two fermion subspace. Excellent convergence in terms of the effective lattice size $N + 1$ is observed except for very small fermion masses $m \lesssim 1/4$ where the two-particle truncation is no longer valid as shown in **Fig. 5.3**. The binding energy of the scalar boson mass M_S in




Figure 5.13: **The binding energy of the scalar boson.** The binding energy of the scalar boson $(M_S - 2m)/g$ is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

the FF is presented in **Fig. 5.14**. The Fock-space has been truncated to the two fermion subspace. Excellent convergence in terms of the effective lattice size $N + 1$ is observed except for very small fermion masses $m/g \lesssim 1/4$ where the two particle truncation is no longer valid as demonstrated elsewhere.

7 The Modified Front Form

The mass of the vector boson computed using the **modified FF formalism** as compared to chiral perturbation theory and the results of Hamer et al.




Figure 5.14: **The binding energy of the scalar boson in the FF**. The binding energy of the scalar boson $(M_S - 2m)/g$, computed in the FF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

The dimensionless momentum N varies logarithmically according to $3 \cdot 2^i$. No convergence is observed in the entire ultra-relativistic region. The *only* point of convergence is the massless limit.

It is possible to expand the FF Hamiltonian in terms of massless fields rather than in terms of massive fields. Details are explained in ref. [49] where this approach was first introduced. This expansion has the advantage that zero-modes are not suppressed as in the massive expansion and the correct Schwinger-boson mass is reproduced in the limit $m = 0$. We shall




Figure 5.15: **The vector boson mass in the modified FF. A comparison with chiral perturbation theory.** The mass of the vector boson in the **modified FF** is plotted against the fermion mass $m/g \lesssim 0.3$ and compared to chiral perturbation theory and the results of Hamer et al. The dimensionless momentum N varies logarithmically according to $3 \cdot 2^i$.

refer to the massless expansion as MODIFIED FRONT FORM(MFF) since this expansion cannot be obtained by means of (Dirac-Bergmann) quantisation of the massive Schwinger model [77, 78] on a light-like quantisation plane (except for the special case $m = 0$). A comparison of the vector boson mass as obtained in the MFF **Fig. 5.15** to the corresponding LMF result **Fig. 5.9** shows that the modified FF is unable to reproduce chiral perturbation theory even though the MFF is able to reproduce the correct Schwinger-boson mass at $m = 0$. The reason for this is that the FF for $m = 0$ is equivalent to CHIRAL QED (QED with right-handed fermions only) if (a) the axial gauge




Figure 5.16: **The number of fermions in a boson (modified FF)**. The number of fermions in the vector boson, the scalar boson and in a two-boson bound-state respectively, plotted against the fermion mass m/g .

is chosen and (b) the $L \rightarrow 0$ limit is performed and (c) the physical states are restricted to the $\mathcal{M} = 0$ sector. The spectrum of this *unphysical* theory⁴ is identical to the spectrum of QED except that it contains right-moving bosons only as the reader may easily verify using the chiral Hamiltonian

$$H_{\text{chiral}} = \sum_{P^3 > 0} \mathbb{A}_{P^3}^\dagger \mathbb{A}_{P^3} \Omega(P^3) + \hat{\mathcal{M}}(\hat{\mathcal{M}} + 1) \quad (5.29)$$

obtained by removing the left-movers from $H_{\text{axial}}(L \rightarrow 0)$ (of course, this

⁴ unphysical since it does not permit large gauge transformations [83])




Figure 5.17: **The binding energy of the vector boson in the modified FF.** The binding energy of the vector boson $(M_V - 2m)/g$, computed in the modified FF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

is a strong violation of Lorentz invariance). The fact that the modified FF cannot be obtained by Dirac-Bergmann quantisation is further illustrated by a comparison of **Fig. 5.16**, **Fig. 5.17**, **Fig. 5.18** and **Fig. 5.15** with the corresponding figures **Fig. 5.3** , **Fig. 5.11** , **Fig. 5.7** and **Fig. 5.9** . In the relativistic limit as well as in the non-relativistic limit this method fails to reproduce the correct results except for $m = 0$ and an intermediate region. The breakdown of the modified FF in the non-relativistic region is due to the fact that the zero-modes are massless in the modified FF. Consequently their kinetic energy is zero which makes them "easily excited": If N is too small




Figure 5.18: **Distribution functions in the chiral limit $m/g \rightarrow 0$ (modified **FF**)**. The distribution function $f(x_B; N = 192)$ is plotted against the Bjorken variable x_B for small masses $m/g \lesssim 0.3$.

to resolve the peak at $x_B = 1/2$, the wave-function spreads towards $x_B = 0$ and $x_B = 1$, zero-modes are excited and the spectrum breaks down as can be seen in **Fig. 5.18**, **Fig. 5.17** and **Fig. 5.19**. The phenomenon that neither the FF nor the modified FF are able to reproduce chiral perturbation theory was first described (but not explained) in Ref. [104] by means of a numerical computation.




Figure 5.19: **The binding energy of the scalar boson in the modified **FF**.** The binding energy of the scalar boson $(M_S - 2m)/g$, computed in the modified FF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

8 The Infinite Momentum Frame

Are the LMF, the IMF and the FF equivalent? In general not, as we have proved in Chapter 3. The massive Schwinger model in the unphysical axial gauge exhibits a remarkable property which distinguishes it from most other field theories: the equivalence between the IMF and the FF as well as an almost exact equivalence between the IMF /FF and the LMF if the effective lattice is infinite ($N = \infty$). If N is finite, these three approximations are




Figure 5.20: **The vector boson mass in the IMF. A comparison with chiral perturbation theory.** The mass of the vector boson in the IMF is plotted against the fermion mass $m/g \lesssim 0.3$ and compared to chiral perturbation theory and the results of Hamer et al. The dimensionless momentum N varies logarithmically according to $3 \cdot 2^i$.

clearly inequivalent as we have already seen in the case of the LMF/FF.

In the previous subsections we have chosen $g \approx 1$ in order to comply with the constraint on $M_1/\Delta k$ imposed by the scaling window (5.16). In this subsection we shall investigate what happens if a very small value of $g/\Delta g$ is chosen such that $M_1(g)$ lies outside the scaling window. This corresponds to choosing the IMF as Δk is very large compared to the vector boson mass M_1 and the fermion mass m such that the approximation

$$\omega(\mathbf{k}) \approx k^3 + \frac{m^2}{2k^3} \quad (5.30)$$




Figure 5.21: **The binding energy of the vector boson in the IMF.** The binding energy of the vector boson $(M_V - 2m)/g$, computed in the IMF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

for the kinetic energy of a given virtual particle becomes exact in the $m\Delta k \rightarrow 0$ limit for every momentum $k^3 = n\Delta k > 0$. The kinetic energy at $k^3 = 0$, in contrast, remains $\omega(0) = m$. A given physical mass

$$M_1^2 = \sqrt{H^2 - P_3^2} \approx 2P^3(H - P^3) \quad (5.31)$$

is the sum of the kinetic squared mass

$$M_{1,kin}^2 = 2P^3(m(b_0^\dagger b_0 + d_0^\dagger d_0) + \sum_{0 < n < N} \frac{m^2}{2n\Delta k} (b_n^\dagger b_n + d_n^\dagger d_n)) \quad (5.32)$$

and the contribution of the interaction H_{pot} . It is immediately clear that the contribution of a zero-mode to a physical mass M is infinitely larger than




Figure 5.22: **The binding energy of the scalar boson in the IMF.** The binding energy of the scalar boson $(M_S - 2m)/g$, computed in the IMF, is plotted against $\log_2(m/g)$. The Fock-space has been truncated to the two fermion subspace.

the contribution of a right-mover since the ratio

$$\frac{m}{m^2/(2n\Delta k)} = 2n\frac{\Delta k}{m} \quad (5.33)$$

goes to infinity in the limit $\Delta k \rightarrow \infty$. Consequently, zero-modes are kinetically repressed in the IMF. As long as the kinetic contribution is not cancelled by a similar contribution from the interaction term, the diverging ratio

$$\frac{\omega(0)}{\omega(\mathbf{k}^3)} > 0 \quad (5.34)$$

has the same effect for $m > 0$ as the explicit constraint that removes zero-modes in the FF.

The contribution m/g^2 of the zero-mode to the total kinetic mass squared M^2/g^2 is finite for any finite fermion mass m/g . This contribution even diverges in the IMF where $g \rightarrow 0$, $m/g = \text{const}$ which means that the zero-mode can not be excited in the IMF limit. Therefore, only the FF contribution to the total mass squared is important. Hence the IMF and the FF are approximately equivalent for the Schwinger model (this does not hold for most other models, in particular renormalisable ones) for $m \neq 0$. For $m = 0$, however, the kinetic energy of the zero-mode cannot diverge and the zero-mode is free, even in the IMF limit. **Fig. 5.9** illustrates this. In particular, we see that the IMF is equivalent to the LMF for $m = 0$ whereas for $m \neq 0$ the IMF and the FF are almost equivalent. In **Fig. 5.20** we chose a small but finite coupling $g/\Delta k = 0.001$ in order to simulate the IMF where $a\xi \gg L$. This scale independence (exact conformal symmetry) of the LMF in the limit $m = 0$ is responsible for the observed exact covariance of the spectrum for $P^3 \geq \Delta k$. Even on a grotesquely small effective lattice with $N + 1 = 2$ only two lattice points does one obtain the exact Schwinger-boson mass.

For larger fermion masses m where dynamics becomes similar to non-relativistic dynamics, the IMF is numerically indistinguishable from the FF as can be inferred by comparing the IMF mass spectrum [Fig. 5.21](#) , [Fig. 5.22](#) with the FF mass spectrum [Fig. 5.12](#) , [Fig. 5.14](#) .

9 Why the Schwinger Model is Special

This might suggest that the FF (or the IMF) are in general equivalent to the IF as long as the continuum version $N = \infty$ of the FF or the IMF is used. It is easy to show that this assumption is wrong in general. The simplest counter-example is the two-flavour Schwinger model with small fermion mass where theories defined on an infinitesimal volume $2L$ as the IMF and the FF must fail. In [\[105, 106\]](#) it is shown that the mass M_π of the lightest boson of this theory is proportional to $m^{2/3}$ for $mL\sqrt{M_v L} \gg 1$ whereas it is proportional to m for $mL\sqrt{M_v L} \ll 1$ with $M_v = \sqrt{\frac{g^2}{2\pi}}$ the mass of the vector boson of this theory (we set $\theta = 0$). For small volumes (i.e. $L \ll 1/M_v$ and $L \ll 1/m$), however, the lightest boson is always proportional to m ; therefore, the FF and the strong IMF cannot reproduce the physical result of the infinite volume limit $M_\pi \propto m^{2/3}$. The weak IMF for $N = \infty$ (being merely the $N \rightarrow \infty$ limit of the LMF) can reproduce this result in principle but it does so with much more numerical effort necessary than that needed in the LMF on a finite lattice. Note that this counter-example cannot be blamed on the well-known difficulties of describing chiral symmetry breaking in the FF: The problem is more fundamental, for there is no spontaneous symmetry breaking in two space-time dimensions according to Coleman's theorem [\[58\]](#).

Things are likely to get worse for the FF for three-dimensional or renormalisable theories: The dominance of the FF contribution to the mass squared

operator is a *very particular* feature of the Schwinger model since, amongst others, it (a) does not contain dynamic bosons (except for the zero-mode) that would (b) fermionic self-energies

$$\begin{aligned}
\Delta\omega_b(\mathbf{k}) &\stackrel{def}{=} \langle 0 | b_{\mathbf{k}} : H : b_{\mathbf{k}}^\dagger | 0 \rangle \\
&= \sum_{l^3 \neq \mathbf{k}} \left(u_{\mathbf{k}}^\dagger u_l \frac{1}{(\mathbf{k} - l)^2} u_l^\dagger u_{\mathbf{k}} - v_l^\dagger u_{\mathbf{k}} \frac{1}{(\mathbf{k} + l)^2} u_{\mathbf{k}}^\dagger v_l \right) \\
&\approx \sum_{\mathbf{k} \neq l^3 \geq 0} \left(u_{\mathbf{k}}^\dagger u_l \frac{1}{(\mathbf{k} - l)^2} u_l^\dagger u_{\mathbf{k}} - v_l^\dagger u_{\mathbf{k}} \frac{1}{(\mathbf{k} + l)^2} u_{\mathbf{k}}^\dagger v_l \right)
\end{aligned} \tag{5.35}$$

associated with right-movers ($\Delta k^3 > 0$) vanish for fast-moving left-movers and vice versa (c) it is super-renormalisable. In general, we are not able to perform such a limit in a meaningful way, especially in renormalisable theories where $P^3 < \Lambda$ has to be *finite*. In general, the importance of zero-modes does not disappear in the limit $N \rightarrow \infty$ especially in the presence of condensates

10 The Influence of the θ Angle

The θ angle in the massless Schwinger model is a super-position of all axial vacua weighted with the factor $e^{i\mathcal{M}\theta}$. If left-movers are excluded from the Fock space, however, only two axial sectors $\mathcal{M} = 0$ and $\mathcal{M} = -1$ can be described because states with $\mathcal{M} < -1$ contain at least $\mathcal{M} + 1$ left-movers. Following Chapter $\langle 3 \rangle$, a rudimentary version of the θ vacua may be constructed,

$$|\theta = 0\rangle = \frac{1}{\sqrt{2}}(|\Omega : 0, 0\rangle + |\Omega : 0, -1\rangle) \tag{5.36}$$

and

$$\begin{aligned}
|\theta = \pi\rangle &= \frac{1}{\sqrt{2}}(|\Omega : 0, 0\rangle - |\Omega : 0, -1\rangle) \\
&= \frac{1}{\sqrt{2}}(|\Omega : 0, 0\rangle + e^{-i\pi} |\Omega : 0, -1\rangle)
\end{aligned} \tag{5.37}$$

corresponding to the theta angles $\theta = 0$ and $\theta = \pi$ which both have a non-vanishing condensate. The state $|\theta = \pi\rangle$ corresponds to the states $|\Omega : \tilde{A}; \theta, N_\theta\rangle$ introduced in Chapter **3** with $N_\theta = 0$ except that —contrary to Chapter **3**— we have chosen axial gauge for numerical computations. Axial




Figure 5.23: **The lowest-lying mass spectrum (for very small fermion masses)**. The mass of the vector meson ($\theta = 0$ and $\theta = \pi$) and the scalar meson ($\theta = 0$ and $\theta = \pi$) are plotted against m/g and compared to the respective linear approximations of chiral perturbation theory. The bifurcations of the vector mass towards $M = 0$ indicate the breakdown of our approximation at a level-crossing.

gauge destroys covariance and the degeneracy of θ states as well. Therefore, the bare vacuum energy $E'_0(\theta)$ depends on the θ angle. In axial gauge, states corresponding to $\theta \neq 0$ appear as spurious states. In our case, the energy $E'_0(-1)$ of the (approximate) vacuum with $\theta = -1$ is slightly larger than the energy $E'_0(0)$ of the $\theta = 0$ vacuum. This means that the state $|\theta = -1\rangle$ ceases to be a vacuum since a vacuum—in common parlance—is defined as the state with lowest energy. If we define the renormalised energy spectrum as $\tilde{E}_n(\theta) \stackrel{\text{def}}{=} E'_n(\theta) - E'_n(0)$ then the "vacuum" with $\theta = \pi$ appears

The Low-Lying Mass Spectrum- Raw Data




Figure 5.24: **The naïve mass spectrum (for very small fermion masses).** The mass of the vector mesons ($\theta = 0$ and $\theta = \pi$) and the scalar mesons ($\theta = 0$ and $\theta = \pi$) are plotted against m/g and compared to the respective linear approximations of chiral perturbation theory. In this figure, the energy of the $\theta = \pi$ vacuum is *not* subtracted from the energies of the $\theta = \pi$ states in order to study the level-crossings.

as a non-covariant spurious state with a finite, momentum-dependent mass.

In this section, we demonstrate that the harm inflicted upon the mass spectrum by axial gauge can be repaired —to a remarkable degree— by treating even degenerate "vacua" as vacua: this means that we define the renormalised energy $E_n(\theta) \stackrel{def}{=} E'_n(\theta) - E'_0(\theta)$ by subtracting the vacuum energy $E'_0(\theta)$ from the bare energy $E'_n(\theta)$ rather than subtracting the energy $E'_0(0)$ of the vacuum proper. This procedure renders the renormalised energies $E_0(\theta) \stackrel{def}{=} E'_0(\theta) - E'_0(\theta) = 0$ of every θ vacuum degenerate — by construction. This renormalisation prescription also allows a correct description of the mass spectrum at very small fermion mass m as shown in **Fig. 5.23**: one observes that our approximation allows us to reproduce the linear approximation (see Ref. [90, 91]) of the scalar boson mass

$$M_V \approx M_B + 2\pi \frac{C(\infty)}{M_B} m \approx M_B + 1.78 \cos(\theta) m \quad (5.38)$$

and vector boson mass

$$M_S \approx 2M_V \quad (5.39)$$

quite nicely in both θ -sectors. Here,

$$C(L) \stackrel{def}{=} \langle \Omega_0; \theta | \bar{\psi} \psi | \Omega_0; \theta \rangle = \frac{e^{\gamma_E}}{2\pi} M_B \cos(\theta) \quad (5.40)$$

is the fermionic CONDENSATE, γ_E is the EULER NUMBER.

We are faced with one practical problem, however. In order to perform the correct renormalisation of the bosonic energies, we have to find out which value of θ a physical state corresponds to. We can do this either by inspection of the wave function for *every* fermion mass or by inspection of the wave function for one fermion mass and then using the continuity of M_n in terms of the fermion mass. We chose the second possibility which is much simpler

from the numerical point of view because it can be easily automated. The first possibility is more powerful though. We could not implement it before the end of the thesis. A disadvantage of using the continuity of $M_n(m)$ is that this scheme does only work before the bare energy $E'_n(\theta)$ crosses the level $E'_{n+1}(\theta)$ of a state with a different θ angle. Then this scheme breaks down as can be seen in **Fig. 5.23** because the wrong vacuum is subtracted in this case. The spurious mass M_n dives into the vacuum and becomes imaginary. The larger the momentum N the smaller the fermion mass where this happens.

In **Fig. 5.24** we depict the vector and scalar masses computed under the assumption that there is only one vacuum: the vacuum corresponding to $\theta = 0$. In this case, the second vacuum appears as a state of its own in **Fig. 5.24**. The $\theta = \pi$ states do not only have the wrong masses without appropriate treatment of the vacuum, they also loose their covariance (i.e. they acquire N -dependence) in a drastic way. Already at very small fermion masses does the $\theta = \pi$ vector state cross the $\theta = 0$ scalar state and a level-crossing occurs. It is this level crossing which limits the fermion mass region in which we are able to describe the $\theta = \pi$ sector correctly as can be seen through a comparison of **Fig. 5.23** and **Fig. 5.24**. The subtraction of the wrong vacuum energy $E'_0(\theta)$ is responsible for the breakdown of covariance and other unphysical effects in **Fig. 5.24** and **Fig. 5.23**

Parts of this chapter have been presented on the IX th International Conference on Recent Progress in Many-Body Theories in Sidney, Australia [107]. A brief summary of these results (and additional figures) is accepted for publication [108].

Chapter 6

The ϕ^4 Theory

Our results concerning the φ^4 theory have been published in [101] and [100]. These papers are enclosed at the end of this thesis. A further paper (on both the φ^4 theory and the Schwinger model) will appear in [107]. The results described in [100, 101] can be further improved upon in order to better deal with the logarithms appearing in the four-dimensional scalar model. We shall describe how to do this in yet another publication.

Chapter 7

Discussion

Que dites-vous? ... C'est inutile? ... Je le sais!
Mais on ne se bat pas dans l'espoir du succès!

[Edmond Rostand: Cyrano de Bergerac]

1 Relativistic Forms

For more than a decade, considerable effort has been invested in order to solve QCD and other field theories using FF techniques. It was thought that the FF could accomplish what the IF could not: describing hadrons as a bound-state of few constituents. We have shown that the FF is not equivalent to FF. We have also shown that the FF is unphysical to a high degree. Some problems and insufficiencies of the FF have been known for a long time. It has been known, for instance, that the FF is unable to describe massless left-movers and that time-ordered propagators depend on an arbitrary choice of prescription. As a reaction to these previously known problems, several disparate philosophies had emerged in the literature

- A1 The **pragmatical** philosophy: some researchers argued that the known problems did not justify abandoning a promising approach: in particular, the successes of the FF in two space-time dimensions were im-

pressive, the massive Schwinger model being a prominent example. It was long claimed —without proof— that these successes could not be repeated in the conventional IF — at least not in such a simple way.

A2 **Other researchers**, however, became suspicious and conjectured that the FF and the IF were not equivalent, acknowledging intrinsic shortcomings. Fascinated by the relative simplicity of the FF, they tried to overcome these shortcomings by treating the FF as an effective theory which may be further improved upon if additional effective interactions are introduced. Some of these researchers tried to turn the badly broken Poincaré symmetry of FF quantisation into an advantage: the class of additional effective interactions —so they conjectured— is defined as the class of operators which restore these broken symmetries when added to the Hamiltonian. This programme might be called FF-bootstrap.

A3 **A third class of researchers**, finally, tried to trace every problem of the FF to the fact that most researchers neglect zero modes.

The results of this Ph.D. thesis render all of these philosophies questionable or unnecessary.

R1 **Firstly**, we have shown that the IF and the FF are not equivalent and that the FF in its pure form leads to completely unphysical predictions such as infinite speed of light, time travel and the violation of both microcausality and causality. There is one theory where the FF yields an *exact* mass-spectrum: The massless Schwinger model. We have traced this to the complete scale-invariance of the Schwinger boson mass and to the fact that the inconsistent chiral Schwinger model becomes consistent for axial boson states in an infinitesimally small box.

Already the massive Schwinger model can be solved only approximately in the FF — albeit with good accuracy.

R2 **Secondly**, we have demonstrated that the FF is not, in general, a low-energy (or low mass) effective theory since it does not do what a low-energy effective theory is supposed to do: reproducing the low-energy spectrum of the full theory. A prominent exception is the FF Hamiltonian of the Schwinger model on an *infinite* effective lattice because (a) no left-movers contribute to the self-energy and (b) the contribution of the zero mode disappears on an infinite lattice. Our findings show that symmetry considerations do not sufficiently constrain the number of possible effective interactions which could improve the FF Hamiltonian. In one spatial dimension, for example, we may replace the FF Hamiltonian H_R by the IMF Hamiltonian or by the effective IMF Hamiltonian H_{eff} . In the IMF there is no need to add additional, effective interactions in order to restore, say, parity symmetry, since the IMF Hamiltonian does not violate this symmetry. Yet the strong IMF is unphysical as we have seen. Thus, parity is not helpful in order to distinguish the unphysical IMF Hamiltonian from a physical Hamiltonian. One might argue that in three spatial dimensions, rotation symmetry is explicitly broken in the IMF, too so that rotation symmetry may serve in order to distinguish the strong IF from the true effective Hamiltonian. Yet if the symmetry argument does not work in one dimension, this FF bootstrap is questionable in three dimensions as well.

R3 **Thirdly**, the problems of the FF cannot be blamed on the zero-modes as we have shown. On the contrary, they even aggravate these problems in some circumstances. For example, a correct treatment of zero-modes

renders the violation of microcausality worse not better. Constraints on zero modes arise due to unphysical BCs or —equivalently— due to the attempt to construct an effective theory on the classical level. Constraints on zero modes are additional symptoms of unphysical assumptions rather than the solution of problems which arise due to exactly these assumptions. Our finding that the FF is the *classically* effective theory of the strong IMF explains why the zero modes of the scalar φ^4 model obey the *classical* constraint of the Landau mean field theory if other modes are discarded. This explains why the FF is unable to go beyond mean-field results in the scalar model. Thus, the problems of the FF do not go away if the zero modes are properly taken into account via the Dirac-Bergmann algorithm.

A quantisation method that is incapable of describing relativistic propagators can hardly be justified. One might argue that some FF successes in one-dimensional theories provide such a justification. Not so. We have shown for QED(1 + 1) that these successes may easily be reproduced within the traditional IF formalism. The FF is not needed. What is more, we have laid bare the mechanism which enables the FF and the IMF to be applied upon the massive Schwinger model: The LMF treatment of this model is almost *scale independent* on a small *infinite* effective lattice. The massless Schwinger model is *exactly scale-independent* even for finite lattices. We have also shown that this scale-independence — which allows for the lattice scaling window to be broken — is peculiar to the Schwinger model: already the two-flavour Schwinger model does not exhibit this feature.

The FF only works (a) if it coincides with a viable approximation in the IF formalism (e.g. the special form of fermionic contractions) or (b) if an infinity of parameters are left free which can be adjusted so as to fit into

reality (e.g. perturbation theory) or (c) in the non-relativistic limit.

More or less byzantine methods may be devised in order to salvage FF perturbation theory (or the FF description of the axial anomaly) using an infinity of parameters as input. One cannot help but noticing, however, that such efforts are but a clumsy, awkward and questionable way of repeating what can be done with relative ease in the IF or with covariant methods (beyond simple Feynman diagrams). Even some successful applications of the FF on mass spectra of two-dimensional theories become bleak when compared to analogous computations in the IF. Therefore, it is necessary to ask the question: Is there any advantage of the FF over the IF/LMF? What do we avail by using the FF if everything that the FF does can be done equally well in the IF and usually with much more ease (but not vice versa)?

2 ε Co-ordinates

We have shown that ε co-ordinates are completely equivalent to the IF quantisation for $\varepsilon \neq 0$ and equivalent to the FF quantisation for $\varepsilon = 0$. The advantages of ε co-ordinates quantisation are by no means due to ε -quantisation itself. We have traced these advantages to the **implicit** choice of small lattice sizes $2L(\varepsilon)$ in the limit $\varepsilon \rightarrow 0$. In this limit, ε quantisation becomes identical to the IMF which violates the lattice scaling window even though it is more physical than the FF and does not destroy microcausality. Only if $\bar{\mathcal{L}}^3(\varepsilon)$ is increased in the limit $\varepsilon \rightarrow 0$ does the lattice size not decrease. In this case, there is no advantage in using these co-ordinates.

3 The Massive Schwinger Model

These results seem to be rather negative since they seem to eliminate the hope of describing QFTs in the Hamiltonian approach. This would be true if the conventional IF were much too complicated in order to describe QFTs as is often claimed in the FF literature. For instance, it is often claimed that the relatively accurate description of the Schwinger model cannot be repeated in the IF, at least not with the ease in which this is done in the FF. Fortunately, this claim has no foundation: we have demonstrated, in Chapter [5](#), that the opposite is true. We have shown that the mechanism which renders the FF (or ε co-ordinates or the IMF) simple is *not* a specific quantisation surface: the relevant mechanism is rather the *implicite choice of an arbitrarily small volume*. This is why the vacuum appears to be almost trivial (i.e. trivial except for zero-modes). In the literature, the (almost) triviality of the FF vacuum is usually claimed to be a result of fact that (a) the vacuum is annihilated by P_- and (b) no constituent particle is allowed to have a negative p_- momentum. This, however, is only a *sufficient* reason for the triviality of the vacuum (if zero-modes are discarded) but by no means *necessary*. The vacuum of the IMF is (almost) trivial even though P_3 is *not* non-positive or non-negative. Unfortunately, an infinitely small volume is unphysical, as we have seen, since it is not compatible with the LGT scaling window. This is why we propose to use a volume that is sufficiently small so as to profit from a simplified vacuum yet sufficiently large so as to fulfill the scaling window of LGT.

Based on this observation, we have put forward another method, the LMF, which shares every advantage of the IMF or the FF while avoiding their respective disadvantages. We have tested this method on two models, $QED(1+1)$ (Chapter [5](#)) and the scalar φ^4 model (Chapter [6](#)) in four

space-time dimensions. Excellent agreement with other methods is found. In the scalar model, our results are compatible with the semi-analytical results obtained by Lüscher and Weisz in the domain of applicability of our method¹. For the Schwinger model our results agree well with the computations of Hamer et al. Contrary to the FF, we were able to verify predictions of chiral perturbation theory on a *finite* lattice.

Chapter 4 uses the exactly known solution of massless $QED(1+1)$ in order to justify the approximations that went into our computations. Furthermore we are able to rigorously demonstrate *that* and *why* a moving frame is advantageous when compared to the rest frame. We pointed out that topological effects, the chiral condensate and the axial anomaly are only essential in the lattice rest-frame. In the LMF, in contrast, discarding topological effects, destroying the chiral condensate or treating the anomaly inadequately does not hamper the correct description of the mass-spectrum in the LMF. We caution that the massive Schwinger model is a rather peculiar theory. Consequently, it is not very likely that all of these results hold in the case of $QCD(3+1)$ as well. Our method works best if the correlation length $a\xi = 1/M$ is sufficiently small when compared to the diameter $2R$ of the physical particle in question. For pions however, the product $2MR$ is very close to one. We may therefore expect that the assumption of a trivial vacuum might not allow us to treat pions correctly. This problem can be dealt with by allowing for a restricted number of lattice sites with negative momenta to be occupied. In addition to that, it will be useful to introduce a *spatially* anisotropic lattice for reasons that will be detailed in another publication.

¹Our application of the LMF method on the scalar model needs some improvement, though (in order to better describe logarithmic corrections). This will be the subject of another publication

4 The Massless Schwinger Model

We have found that the mass spectrum of the Schwinger model is remarkably robust against a number of severe approximations:

- The mass spectrum does not depend on the size L of the lattice. Consequently, even the strong IMF is able to reproduce the correct spectrum.
- Axial gauge does not modify the mass spectrum for $L \rightarrow 0$ except for the spectrum in the lattice rest frame.
- The diagonalisation of the Hamiltonian may be performed using a basis with definite axial charge instead of using gauge-invariant θ states.
- For $L \rightarrow 0$ the forcible exclusion of fermionic left-movers has no effect whatsoever on the mass spectrum, even if left-movers are removed on the classical level. We have traced the latter feature to the fact that *fermionic* left-movers do not contribute to the self-energy of right-movers.
- The effective size $N + 1$ of the lattice may be chosen to be ridiculously small: One obtains the *exact* Schwinger boson mass even on a effective lattice with two lattice sites.

This was elaborated in Chapter [4](#). This robustness towards severe approximations is the reason for why the FF and the IMF are able to reproduce the boson masses of the Schwinger model. If all of the approximations listed above are performed, the Schwinger model is formally equivalent to the unphysical *chiral* Schwinger model² or to the FF. See Chapter [4](#) and [5](#).

²It is remarkable that the chiral Schwinger model is even inconsistent without these approximations.

We found that the volume dependence of the massless Schwinger model differs drastically from the massive Schwinger model: the mass spectrum of massive Schwinger model is sensitive to correct implementation of the scaling window $2L > a\xi$ whereas the mass spectrum of the massless Schwinger model comes out correctly even if the scaling window is not fulfilled: Schwinger boson masses are independent of the lattice size $2L$ — even in the rest frame. *If and only if* an infinite effective lattice $N = \infty$ is used, the massive Schwinger model, too, becomes insensitive to the lattice size $2L$ and the LMF Schrödinger equation for the vector state coincides with the FF Schrödinger equation. Therefore, the results obtained in Ref. [50] for the vector state may be considered as LMF results in the limit $N = \infty$. We have traced this coincidence to the fact that the Schwinger model does not contain a four-boson interactions.

Topological effects contribute primarily to the rest-frame sector $\mathbf{P} = 0$ where the De Broglie wave-length $\lambda = \frac{2\pi}{|\mathbf{P}|} = \infty$ is larger than any finite lattice size $2L$. This implies, so we argue, that the relations between the mass-spectrum and the chiral condensate obtained in chiral perturbation theory are fundamentally different for $\mathbf{P} = 0$ and for $|\mathbf{P}| \neq 0$ respectively. Since the mass spectrum of QED(1 + 1) is almost insensitive to the lattice size $2L > 1/M$ in the LMF, masses computed in the LMF are related to the fermionic condensate $C(\infty)$ obtained at infinite volume rather than to the condensate $C(L) = \langle P | \bar{\psi}\psi | P \rangle$ at finite volume³. For this to be true, it is *de rigueur* that no approximation destroy Lorentz covariance. The spectrum at $\mathbf{P} \approx 0$, however, is related to the condensate $C(L)$ at finite volume.

³ Intuitively, one might conjecture that the chiral condensate extracted from the m -dependence of the physical mass spectrum is equal to $C(L/\sqrt{1-v^2})$ where v is the velocity of the physical particle. For $v \approx 1$ we have $C(L/\sqrt{1-v^2}) \approx C(\infty)$.

5 Structure Functions

We have demonstrated in Chapter 2 that the impulse approximation which relates structure functions to a convolution of distribution functions and parton cross sections is *frame dependent* due to the non-triviality of the vacuum. For frames similar to the Breit-frame, however, this frame dependence is negligible and the IA is a good approximation. It is well known that the rest-frame is not suited in order to interpret structure functions in terms of quarks and gluons. We have shown, additionally, that the IMF is not suited either, since close to both frames, the structure functions calculated using the IA show a strong dependence on the momentum signaling the breakdown of the IA. This frame dependence stems from the fact that the vacuum is probed in both extreme cases. The fact that this frame dependence is so strong stems from the fact that the vacuum quark distribution (which is not a physical observable) diverges for small momenta $\mathbf{k}$.

Structure functions are related to *space-like* correlation functions. They are *not* related to *light-like* correlation functions. Space-like correlation functions attain their maximal simplicity in AXIAL GAUGE and *not* in the notoriously difficult [53] AXIAL GAUGE. The Taylor series of space-like correlation functions with respect to x^3 corresponds to the OPERATOR PRODUCT EXPANSION (OPE) in the sense that the n -th Taylor coefficient is proportional to the n -th moment⁴ of the structure function F_2^{IA} . This means that the OPE is a *short-distance* expansion (at least in the IA) and *not* a problematic light-cone expansion. This is conceptually advantageous because the OPE does —strictly speaking— only make sense as a short-distance expansion.

⁴ It is well-known that the same statement holds *formally* for the Taylor series of light-like correlation-functions in terms of x^- .

Bibliography

- [1] G. Parisi, An Introduction to Scaling Violations, in *Int. Meeting on Neutrino Physics, Flaine, France*, 1976.
- [2] R. G. Roberts, *The Structure of the Proton: Deep Inelastic Scattering* (Cambridge, UK: Univ. Pr. (Cambridge monographs on mathematical physics), 1990).
- [3] H. J. Rothe, *Lattice Gauge Theories: An Introduction* (Singapore, Singapore: World Scientific, 1992).
- [4] I. Montvay and G. Münster, *Quantum Fields on a Lattice* (Cambridge, UK: Univ. Pr. (Cambridge monographs on mathematical physics), 1994).
- [5] D. Weingarten, Nucl. Phys. **B215**, 1 (1983).
- [6] W. Lee and D. Weingarten, Nucl. Phys. Proc. Suppl. **53**, 236 (1997), hep-lat/9608071.
- [7] K. G. Wilson, Phys. Rev. **D10**, 2445 (1974).
- [8] F.J.Wegner, J.Math.Phys. **10**, 2249 (1971).
- [9] G. Martinelli and C. T. Sachrajda, Nucl. Phys. **B316**, 355 (1989).

- [10] G. Martinelli and C. T. Sachrajda, Nucl. Phys. **B306**, 865 (1988).
- [11] G. Martinelli and C. T. Sachrajda, Phys. Lett. **196B**, 184 (1987).
- [12] G. Martinelli and C. T. Sachrajda, Phys. Lett. **190B**, 151 (1987).
- [13] C. Best *et al.*, Phys. Rev. **D56**, 2743 (1997), hep-lat/9703014.
- [14] M. Göckeler *et al.*, Phys. Lett. **B391**, 388 (1997), hep-lat/9609008.
- [15] M. Göckeler *et al.*, Nucl. Phys. Proc. Suppl. **53**, 81 (1997), hep-lat/9608046.
- [16] M. Göckeler *et al.*, Phys. Rev. **D54**, 5705 (1996), hep-lat/9602029.
- [17] M. Göckeler *et al.*, Nucl. Phys. Proc. Suppl. **49**, 250 (1996), hep-lat/9601007.
- [18] M. Göckeler *et al.*, (1995), hep-lat/9511025, Zeuthen 1995, Proceedings, Prospects of spin physics at HERA*339.
- [19] M. Göckeler *et al.*, Phys. Rev. **D53**, 2317 (1996), hep-lat/9508004.
- [20] K.-F. Liu and S.-J. Dong, (1994), hep-lat/9411067.
- [21] W. Wilcox and B. Anderson-Pugh, (1993), hep-lat/9312034.
- [22] M. Alford, T. R. Klassen, and G. P. Lepage, Nucl. Phys. **B496**, 377 (1997), hep-lat/9611010.
- [23] C. J. Morningstar and M. Peardon, Phys. Rev. **D56**, 4043 (1997), hep-lat/9704011.
- [24] C. Morningstar and M. Peardon, Nucl. Phys. Proc. Suppl. **53**, 917 (1997), hep-lat/9608050.

- [25] C. Morningstar, Nucl. Phys. Proc. Suppl. **53**, 914 (1997), hep-lat/9608019.
- [26] J. Kogut and L. Susskind, Phys. Rev. **D11**, 395 (1975).
- [27] M. Lüscher and G. Münster, Nucl. Phys. **B232**, 445 (1984).
- [28] J. Koller and P. van Baal, Phys. Rev. Lett. **58**, 2511 (1987).
- [29] B. van den Heuvel and P. van Baal, Nucl. Phys. Proc. Suppl. **42**, 823 (1995), hep-lat/9411046.
- [30] A. M. Chaara, H. Kröger, L. Marleau, K. J. M. Moriarty, and J. Potvin, Phys. Lett. **B336**, 567 (1994), hep-lat/9403023.
- [31] D. Bérubé, H. Kröger, R. Lafrance, and L. Marleau, Phys. Rev. **D43**, 1385 (1991).
- [32] D. Bérubé, J. F. Brière, H. Kröger, K. J. M. Moriarty, and J. Potvin, Nucl. Phys. Proc. Suppl. **17**, 744 (1990).
- [33] H. Kröger, K. J. M. Moriarty, and J. Potvin, Scattering Theory on the Lattice and with Monte-Carlo, LAVAL-PHY-89/6.
- [34] J. F. Brière and H. Kröger, Phys. Rev. Lett. **63**, 848 (1989).
- [35] C. P. van den Doel and D. Horn, Phys. Rev. **D35**, 2824 (1987).
- [36] X.-Q. Luo, S.-H. Guo, H. Kröger, and D. Schütte, (1997), hep-lat/9711042.
- [37] D. Schütte, W.-H. Zheng, and C. J. Hamer, Phys. Rev. **D55**, 2974 (1997), hep-lat/9603026.

- [38] C. J. Hamer, M. Sheppeard, W.-H. Zheng, and D. Schütte, Phys. Rev. **D54**, 2395 (1996), hep-th/9511179.
- [39] X.-Q. Luo, Q.-Z. Chen, S.-H. Guo, X.-Y. Fang, and J.-M. Liu, Nucl. Phys. Proc. Suppl. **53**, 243 (1997), hep-ph/9609389.
- [40] X.-Q. Luo and Q.-Z. Chen, Mod. Phys. Lett. **A11**, 2435 (1996), hep-ph/9604395.
- [41] S. J. Brodsky, G. McCartor, H. C. Pauli, and S. S. Pinsky, Part. World **3**, 109 (1993).
- [42] C. J. Morningstar and M. Weinstein, Phys. Rev. **D54**, 4131 (1996), hep-lat/9603016.
- [43] C. J. Morningstar and M. Weinstein, (1994), hep-lat/9409007, Published in the proceedings of the 27th International Conference on High Energy Physics, 1994:1045.
- [44] C. J. Morningstar and M. Weinstein, Phys. Rev. Lett. **73**, 1873 (1994), hep-lat/9405020.
- [45] S.-H. Guo and X.-Q. Luo, (1997), hep-lat/9706017, Experimental and Theoretical Study of Glueballs, CCAST-WL Workshop Series: Volume 76, 127-138.
- [46] R. Haag, Dan. Mat. Fys. Medd. **29**, 1, No. 12 (1955).
- [47] P. A. M. Dirac, Rev. Mod. Phys. **21**, 392 (1949).
- [48] M. Heyssler and A. C. Kalloniatis, Phys. Lett. **B354**, 453 (1995), hep-th/9505126.

- [49] T. Eller, H. C. Pauli, and S. J. Brodsky, Phys. Rev. **D35**, 1493 (1987).
- [50] Y.-Z. Mo and R. J. Perry, J. Comput. Phys. **108**, 159 (1993).
- [51] N. E. Ligterink and B. L. G. Bakker, Phys. Rev. **D52**, 5954 (1995), hep-ph/9412315.
- [52] N. C. J. Schoonderwoerd and B. L. G. Bakker, (1997), hep-ph/9702311.
- [53] G. McCartor and D. G. Robertson, Z. Phys. **C62**, 349 (1994), hep-th/9311065.
- [54] S. Mandelstam, Nucl. Phys. **B213**, 149 (1983).
- [55] G. Leibbrandt, Phys. Rev. **D29**, 1699 (1984).
- [56] M. Lüscher and P. Weisz, Nucl. Phys. **B290**, 25 (1987).
- [57] J. L. G. E. Brézin and J. Zinn-Justin, Test, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and M. S. Green, p. 125, Academic Press, London, 1976.
- [58] C. Itzykson and J. B. Zuber, *Quantum Field Theory* (New York, USA: McGraw-hill (1980) 705 P.(International Series in Pure and Applied Physics), 1980).
- [59] N. Scheu, Extraction of Structure Functions Using Discretised Light-Cone Quantisation, Diploma thesis, Ruperto Carola University, Heidelberg, 1993, (In German).
- [60] R. L. Jaffe, (1996), hep-ph/9602236, In *Lenz, F. (ed.): Lectures on QCD* 178.
- [61] K.-F. Liu and S.-J. Dong, Phys.Rev.Lett. **72**, 1790 (1994).

- [62] F. E. Close, *An Introduction to Quarks and Partons* (Academic Press, London, 1979).
- [63] G. Altarelli, Phys. Rept. **81**, 1 (1982).
- [64] S. Weinberg, Phys. Rev. **150**, 1313 (1966).
- [65] H. Leutwyler and J. Stern, Ann. Phys. **112**, 94 (1978).
- [66] R. Neville and F. Rohrlich, Nuovo Cimento **A1**, 625 (1971).
- [67] P. J. Steinhardt, Ann. Phys. **128**, 425 (1980).
- [68] T. Heinzl and E. Werner, Z. Phys. **C62**, 521 (1994), hep-th/9311108.
- [69] E. V. Prokhvatilov and V. A. Franke, Yad. Fiz. **49**, 1109 (1989).
- [70] F. Lenz, M. Thies, K. Yazaki, and S. Levit, Annals Phys. **208**, 1 (1991).
- [71] H. W. L. Naus, H. J. Pirner, T. J. Fields, and J. P. Vary, (1997), hep-th/9704135.
- [72] S. Weinberg, *Gravitation and Cosmology* (John Wiley & Sons, 1972).
- [73] K. Sundermeyer, *Constrained Dynamics with Applications to Yang-Mills Theory, General Relativity, Classical Spin, Dual String Model* (Berlin, Germany: Springer (Lecture Notes In Physics, 169), 1982).
- [74] P. A. M. Dirac, Can. J. Math. **2**, 129 (1950).
- [75] M. Burkardt, (1997), hep-ph/9709421.
- [76] M. Burkardt, Adv. Nucl. Phys. **23**, 1 (1996), hep-ph/9505259.
- [77] D. Mustaki, Phys. Rev. **D42**, 1184 (1990).

- [78] A. C. Kalloniatis and D. G. Robertson, Phys. Rev. **D50**, 5262 (1994), hep-th/9405176.
- [79] T. Heinzl and E. Werner, Z. Phys. **C62**, 521 (1994), hep-th/9311108.
- [80] A. Langnau and S. J. Brodsky, J. Comput. Phys. **109**, 84 (1993).
- [81] J. Schwinger, Phys. Rev. **125**, 397 (1962).
- [82] J. Schwinger, Phys. Rev. **128**, 2425 (1962).
- [83] N. S. Manton, Annals Phys. **159**, 220 (1985).
- [84] J. Kiskis, Phys. Rev. **D18**, 3690 (1978).
- [85] J. Curtis G. Callan, R. Dashen, and D. J. Gross, Phys. Rev. **D17**, 2717 (1978).
- [86] J. Ambjorn, J. Greensite, and C. Peterson, Nucl. Phys. **B221**, 381 (1983).
- [87] S. L. Adler, Phys. Rev. **177**, 2426 (1969).
- [88] J. S. Bell and R. Jackiw, Nuovo Cim. **60A**, 47 (1969).
- [89] K. Fujikawa, Phys. Rev. Lett. **42**, 1195 (1979).
- [90] S. Coleman, R. Jackiw, and L. Susskind, Ann. Phys. **93**, 267 (1975).
- [91] S. Coleman, Ann. Phys. **101**, 239 (1976).
- [92] N. N. Bogolubov, Nuovo Cim. **7**, 794 (1958).
- [93] N. N. Bogolubov, Zh. Eksp. Teor. Fiz. **34**, 58 (1958).
- [94] G. McCartor, Int. J. Mod. Phys. **A12**, 1091 (1997), hep-th/9609140.

- [95] C. J. Hamer, W.-H. Zheng, and J. Oitmaa, Phys. Rev. **D56**, 55 (1997), hep-lat/9701015.
- [96] X.-Y. Fang, X.-Q. Luo, G.-C. Xu, and Q.-Z. Chen, Z. Phys. **C54**, 587 (1992).
- [97] X.-Q. Luo and Q.-Z. Chen, J. Phys. G **G16**, 1181 (1990).
- [98] X.-Q. Luo, Z. Phys. **C48**, 283 (1990).
- [99] Q.-Z. Chen and X.-Q. Luo, Phys. Rev. **D42**, 1293 (1990).
- [100] H. Kröger and N. Scheu, Phys. Rev. **D56**, 1455 (1997), hep-lat/9607006.
- [101] H. Kröger and N. Scheu, Nucl. Phys. Proc. Suppl. **53**, 321 (1997), hep-lat/9607008.
- [102] J. P. Vary, T. J. Fields, and H. J. Pirner, Phys. Rev. **D53**, 7231 (1996).
- [103] J. P. Vary, T. J. Fields, and H. J. Pirner, (1994), hep-ph/9411263.
- [104] S. Elser, The Spectrum of $QED(1+1)$ in the Frame-Work of the DLCQ Method, Diploma thesis, Ruperto Carola University Heidelberg, 1994.
- [105] J. E. Hetrick, Y. Hosotani, and S. Iso, Phys. Lett. **B350**, 92 (1995), hep-th/9502113.
- [106] Y. Hosotani, (1995), hep-th/9505168.
- [107] H. Kröger and N. Scheu, Computation of Structure Functions from a Lattice Hamiltonian, in *Proceedings of the IX th International Conference on Recent Progress in Many-Body Theories*, edited by D.Neilson, World Scientific, Singapore, 1997, accepted for publication.

- [108] H. Kröger and N. Scheu, Accepted for publication in Phys.Lett **B** (1998), hep-lat/9804024, ULaval-Laphyt-March-98.

Index

- $\vdash$, 67
- $\hat{}$, 15
- θ -states, 115
- $\triangleq$, 66, 73
- $\top$, 15
- $\dagger$, 15
- $\therefore$, 15
- $\mathbb{I}$, 24
- A , 107
- $\mathbb{A}$, 124
- $\tilde{A}$, 122
- $\mathcal{A}$, 123
- A , 107
- a , 118
- a (lattice spacing), 62
- $\mathbf{a}$, 62
- a -vacuum, 112
- a -vacuum ($|\Omega_a : 0, A_3\rangle$), 112
- absolute velocity, 78
- α -, 97
- ambiguities, 72
- anti-periodic BCs, 86
- artificial Hamiltonian, 122
- aspect of simultaneity, 57
- at rest, 21
- axial anomaly, 115
- axial charge, 111
- axial current, 111
- Axial gauge, 117
- axial gauge, 36, 189
- axial transformations, 121
- axial vacuum, 112
- axiomatic field theory, 8
- b , 118
- BC, 54
- bd-vacuum, 118
- binding energy, 160, 162
- Bjorken limit, 45, 48
- Bjorken scaling, 47
- Bjorken scaling variable, 20
- bosonisation, 103
- boundary conditions, 54
- Breit frame, 21, 48

- C, 128
- $C(L)$, 177
- $\mathcal{C}$, 92
- c , 13
- causality region, 92
- chiral QED, 165
- chiral Schwinger model, 100
- chiral spinors, 109
- classically effective Hamiltonian, 84, 86
- closeness to the light-cone, 77
- closeness to the light-front, 72
- co-ordinate, 70
- co-ordinate boost, 70
- co-ordinate boosts, 71, 93
- co-ordinate frame, 56
- co-ordinate transformation, 56
- co-variant, 56
- condensate, 177
- constituent picture, 131
- constituent quark model, 124
- constraints, 59
- Continuum limit, 113
- continuum limit, 3, 48, 146, 147
- contra-gradient, 67
- contra-variant, 56
- $\otimes$, 46
- convolution, 46
- correlation length, 139
- Coulomb gauge, 108
- covariance limit, 146, 148, 149
- current operator, 18
- cyclic group, 128
- cyclic vacua, 129
- D : covariant derivative, 141
- d , 118
- Deep inelastic scattering, 17
- deep(ly) inelastic scattering, 1
- DGLAP, 46
- DIS, 1
- discrete Fermi level, 112
- discretised light-cone quantisation, 54
- distribution function
 - parton, 34, 35
 - raw, 30
- distribution functions, 2
- $\Delta\omega$, 81
- $d^R k$, 24
- $d^{[R]} k$, 24
- dual frame, 56
- dynamical, 41
- E , 17, 18
- E, 109

$e_{(\mu)}$, 56
 E_B , 22
 effective, 82
 effective lattice, 147
 effective lattice size, 79, 147
 ELGT, 2
 energy-momentum tensor, 58
 ephemeral stability group, 101
 e^{quant} , 59
 Euclidean lattice gauge theory, 2
 Euler number, 177
 exclusion on the classical level, 83
 exclusion on the quantum level, 83
 FF, 9, 51
 FF causality region, 97
 first class constraint, 108
 flow of artificial time, 60
 frame of reference, 55
 front form, 51
 front-form, 9
 f_s , 30
 fundamental modular domain, 108
 g , 14, 106
 g : coupling constant in the Schwinger
 model, 141
 $\mathbf{g}$, 35
 γ : Dirac matrices, 107
 γ_E , 177
 gauge transformation, 107
 gauge-transformations, 121
 Gauss' law, 108
 general co-ordinate transformation,
 55
 general co-ordinate transformations,
 56
 General co-ordinates, 56
 global Wilson loop, 108
 gluons, 122
 H : Hamiltonian
 of the Schwinger model, 110
 $\mathcal{H}$, 19
 $\hbar$, 13
 h , 133
 Hamiltonian, 4, 109
 Hamiltonian in axial gauge, 122
 heat-kernel regularisation, 113, 121
 Heisenberg picture, 15
 helicity, 14, 19
 helicity polarisation axis, 19
 helicity-direction, 25
 H_I , 82
 higher twist corrections, 47
 H_n , 126
 $H_{\text{N.O.}}$, 109

H_{raw} , 110
 H_{eff} , 86
 H_{raw} , 119
 ι , 107
 IA, 12, 23
 IF, 9, 51
 IMF, 6, 51
 impulse approximation, 12, 23, 45
 inclusive scattering, 17
 Inertial Frame, 55
 infinite momentum frame, 6, 51, 144
 initial surface, 49, 50
 instant form, 9, 51
 instant of time, 49, 50, 57
 $\hat{j}^\mu$, 18
 $\mathbf{J}$, 18
 $\tilde{j}^\mu$, 110
 j^μ , 110
 k , 17
 k' , 17
 kinematical, 41, 52, 59
 kinematical equivalence, 66
 kinematical equivalence of quantisation-frames, 12
 kinematically equivalent, 66
 kinetic energy, 119
 Kogut-Susskind Hamiltonian, 4
 L , 76, 104, 140, 188
 $\mathcal{L}$:
 Lagrangian density, Schwinger model, 107
 $\mathfrak{L}$, 62
 ℓ , 35
 Λ , 62
 large gauge transformations, 107, 115, 119, 120
 large momentum frame, 6, 139
 Lattice Gauge Theory, 2
 lattice rest frame, 55
 lattice size, 140
 lattice spacing, 72
 L_c , 76
 left, 105
 left-movers, 83, 140
 leptonic tensor, 34
 LGT, 2
 Λ_h , 113
 light-cone, 51
 light-cone co-ordinates, 13
 light-cone gauge, 36
 light-front, 51
 light-front co-ordinates, 13
 LMF, 6, 139

longitudinal, 18
 longitudinally polarised, 19
 $\mathcal{L}_R$, 86
 M , 17, 18, 139
 $\mathcal{M}$, 119, 174
 m :
 mass of a virtual fermion in the
 Schwinger model, 141
 $m_\perp$, 82
 massive Schwinger model, 7, 103
 mathematica, 15, 16
 M_B , 103
 M_B , 145
 metric aspect, 57
 MFF, 165
 modified front form, 165
 modified kinetic energy, 120
 modulo division, 128
 momentum transfer, 20
 M_S , 162
 M_V , 139
 N , 79
 naïve axial charge, 112
 naïve parton model, 44, 47
 Normal ordering, 15
 null-plane, 51
 n_W , 107
 $\Omega(p^3)$, 116
 $|\Omega \dots\rangle$, 132
 $|\Omega : \mathbf{A}_3, 0\rangle$, 118
 $|\Omega_a : \mathbf{A}_3\rangle$, 112
 $|\Omega : \tilde{\mathbf{A}}; \theta\rangle$, 125, 128
 $|\Omega_n : \mathcal{M}\rangle$, 126
 $|\Omega_n; \theta\rangle$, 126
 operator, 15
 operator product expansion, 46, 189
 ortho-normal frame, 56
 P , 17, 59, 139
 P' , 17
 $\mathbf{P}$, 18
 $\mathbf{P}(e^{\text{quant}})$, 59
 $\mathcal{P}$, 97
 $\mathcal{P}_{x_B}$, 151
 pairing, 81
 particle-hole representation, 109
 parton distribution, 31, 134
 parton distribution functions, 34,
 35
 parton helicity, 24
 parton model, 6
 path-integral quantisation, 115
 Pauli-Lubansky four-vector, 18
 P_B , 22
 perturbation theory, 115

perturbative vacuum, 118, 125
 $\varphi(0, \mathbf{x})$, 59
 physical boosts, 70
 physical Hamiltonian, 69
 physical subspace, 116
 $\pi(0, \mathbf{x})$, 59
 Π_0 , 84
 Π_l , 84
 Π_r , 84
 point-form, 51
 polarised parton distribution function, 35
 prescription, 97
 principal value prescription, 97
 probability, 151
 projection operators, 84
 proper time, 56, 57
 $|PS\rangle$, 18
 $|\Psi \dots\rangle$, 132
 $\psi(x)$: fermionic field, 109
 ψ_A , 25
 $|\Psi_n; \theta\rangle$, 132
 $|P', X\rangle$, 17

 Q , 20
 $\tilde{q}^\mu$, 35
 q , 17
 $\mathbf{q}$, 35

 $\mathbf{q}_s$, 35
 $\mathbf{q}_s$, 35
 $:\mathcal{Q}_A:$, 126
 $\mathcal{Q}_A$, 119
 $\mathcal{Q}_A^{\text{reg}}$, 126
 QCD, 1, 2
 QED(1 + 1), 9, 103
 QFT, 8
 Quantisation Frame, 55
 quantisation frame, 58, 60
 quantisation surface, 49
 Quantum Chromo-Dynamics, 2
 Quantum chromo-dynamics, 1
 quantum effective Hamiltonian, 86
 quantum electro-dynamics, 9
 quark distribution function, 2
 quarks, 122
 quenched approximation, 3

 R , 139
 $\mathcal{R}$, 81
 radius, 139
 raw distribution functions, 30, 134
 raw pairing functions, 30
 real time, 51
 regularisation, 2
 relative velocity, 37, 77
 renormalisation group equations, 45

renormalisation group improved par-
ton model, 47
renormalisation of the speed of light,
4
renormalised, 111
resolution, 20
rest frame of the observer, 56
 ρ_L , 110
 ρ_R , 110
right, 18, 105
right-movers, 83

 S , 18
 $\uparrow$
 s , 24
 $\uparrow$
 s_{spin} , 25
scaling window, 7
Schwinger bosons, 116
Schwinger model, 7, 100, 103
self-energy, 81
set of causally independent points,
50
 $\uparrow$
 S_H , 19
 σ : Pauli matrices, 107
size, 144
size of the Lattice, 62
smooth cut-off, 113
space-time volume, 38
spin quantisation axis, 25
spurious states, 123
 $\uparrow$
 $\hat{S}_T$, 19
stability group, 51
stability group proper, 101
strong exclusion, 83
strong IMF, 95, 137
strong infinite momentum frame, 78
structure functions, 1, 20
super-renormalisable, 111

 T , 82, 119
 T^A , 26
 T^B , 26
 T^C , 26
 $\mathbb{T}$, 4
 T^+ , 67
 $\mathfrak{T}$, 120
 $\mathcal{T}$, 19
 t , 49
 T^{AB} , 32
 τ , 56
 $T(\varepsilon)$, 72
thermodynamic limit, 5
 $\Theta^{\mu\nu}$, 58
 θ , 142
 $\theta(x)$, 134
time axis, 60
time-evolution operator, 23

topological glueball, 127
 topologically non-trivial gauge trans-
 formations, 107
 total angular momentum three-vector,
 18
 transfer matrix, 4
 Transitivity, 53
 transverse mass, 82
 transverse polarisation axis, 19
 transverse spin, 19
 transversely polarised., 19
 T_{raw} , 111
 triangle diagram, 115
 trivial, 125

 U , 23
 $\mathbf{u}$, 118
 U_A , 25
 $U_B(L)$, 132
 U_G , 115, 120
 $U_G^{(N_\theta)}$, 128
 ultra-relativistic region, 147
 unpolarised parton distribution func-
 tion, 35

 v , 139
 $\mathbf{v}$, 118
 vacuum, 104
 vector boson, 139, 143
 vector charge, 111
 vector current, 110, 111
 virtual particles, 140
 virtual particles associated with the
 vacuum, 140, 144

 weak exclusion, 83
 weak infinite momentum frame, 78
 weak interactions, 21
 Wilson action, 3
 winding number, 107

 w_L , 109
 $W^{\mu\nu}$, 18
 $W_A^{\mu\nu}$, 20
 $W_S^{\mu\nu}$, 19
 w_R , 109

 $\bar{x}^\mu(\varepsilon)$, 72
 $\check{x}^\mu$, 74
 $\tilde{x}^\mu$, 74
 x , 106
 x^μ , 13
 x_μ , 13
 x_B , 20
 ξ , 139
 Ξ_s , 30
 $\bar{x}^\mu$, 56

 zero modes, 83

zero volume basis, 124

zero-mode of the gauge field, 107